

Information Limits for Detecting a Subhypergraph

Mingao Yuan¹ and Zuofeng Shang²

¹*Department of Statistics, North Dakota State University, e-mail: mingao.yuan@ndsu.edu*

²*Department of Mathematical Sciences, New Jersey Institute of Technology, e-mail: zshang@njit.edu*

Abstract: We consider the problem of recovering a subhypergraph based on an observed adjacency tensor corresponding to a uniform hypergraph. The uniform hypergraph is assumed to contain a subset of vertices called as subhypergraph. The edges restricted to the subhypergraph are assumed to follow a different probability distribution than other edges. We consider both weak recovery and exact recovery of the subhypergraph, and establish information-theoretic limits in each case. Specifically, we establish sharp conditions for the possibility of weakly or exactly recovering the subhypergraph from an information-theoretic point of view. These conditions are fundamentally different from their counterparts derived in hypothesis testing literature.

MSC2020 subject classifications: Primary 62G10; secondary 05C80.

Keywords and phrases: sharp information-theoretic condition, uniform hypergraph, subhypergraph detection, weak recovery, exact recovery.

1. Introduction

An *undirected* m -uniform hypergraph is a pair $(\mathcal{V}, \mathcal{E})$ in which $\mathcal{V} = [N] := \{1, 2, \dots, N\}$ is a vertex set and \mathcal{E} is an edge set. Each edge in \mathcal{E} consists of exactly m vertices in \mathcal{V} . The corresponding adjacency tensor is an m -dimensional symmetric array $A \in (B^N)^{\otimes m}$ satisfying $A_{i_1 i_2 \dots i_m} \in B$ for $1 \leq i_1 < i_2 < \dots < i_m \leq N$, in which $B \subset \mathbb{R}$. Here, symmetry means that $A_{i_1 i_2 \dots i_m} = A_{j_1 j_2 \dots j_m}$ whenever i_1, i_2, \dots, i_m is a permutation of j_1, j_2, \dots, j_m . If $|\{i_1, i_2, \dots, i_m\}| \leq m - 1$, then $A_{i_1 i_2 \dots i_m} = 0$, i.e., no self-loops are allowed. In particular, $B = \{0, 1\}$ corresponds to binary hypergraphs. The general B corresponds to weighted hypergraphs.

Given probability distributions P and Q over B , let $\mathcal{H}_m(N, Q; n, P)$ denote a uniform hypergraph model including a subhypergraph of cardinality n defined as follows: for a uniformly and randomly drawn subset $S^* \subset [N]$ with $|S^*| = n$,

$$A_{i_1 i_2 \dots i_m} \sim \begin{cases} P, & i_1, \dots, i_m \in S^*, \\ Q, & \text{otherwise.} \end{cases} \quad (1)$$

Observing A , we are interested in recovering the subhypergraph S^* . When P and Q are both Bernoulli distributions, various algorithms have been developed by [3, 5, 9, 12, 14, 17, 6, 23] for $m = 2$, and by [16, 22, 21, 26, 11, 4, 20, 18, 21] for $m \geq 3$. For general P and Q with $m = 2$, [13] studied this problem systematically from information-theoretic viewpoint and derived sharp recovery boundaries. For $m \geq 3$, relevant literature only exist in block models including [1, 2, 7, 19]. Specifically, under stochastic block models, [7] obtained minimax detection bounds, [1] derived sufficient conditions for weak or exact recovery, [19] derived exact recovery information limits; under generalized censored block models, [2] derived sufficient and necessary conditions for weak recovery. To the best of our knowledge, information limits for detecting a subhypergraph under model (1) are nonexistent. In this paper, we will study this problem for arbitrary probability distributions P, Q and integer $m \geq 2$.

Our theoretical results include information limits for weak/exact recovery of the subhypergraph obtained in an asymptotic regime $N \rightarrow \infty$ and $\limsup n/N < 1$. Let $D(P||Q) = \mathbb{E}_P \left(\log \frac{dP}{dQ} \right)$ be the Kullback-Leibler

divergence from Q to P and both $D(P\|Q)$, $D(Q\|P)$ are supposed to be finite. We derive sharp regions characterized by N, n for weakly or exactly recovering the subhypergraph with computational complexity aside which provides a benchmark for developing polynomial time algorithms. Specifically, our main results are summarized in Table 1.

TABLE 1
Regions for recovering a subhypergraph.

Region	Detectability
(a) $n^{m-1}D(P\ Q) \rightarrow \infty$ and $\liminf_{N \rightarrow \infty} \frac{n^{(m)}D(P\ Q)}{n \log \frac{N}{n}} > 1$	Weak recovery of subhypergraph is possible
(b) $n^{m-1}D(P\ Q) = O(1)$ or $\limsup_{N \rightarrow \infty} \frac{n^{(m)}D(P\ Q)}{n \log \frac{N}{n}} < 1$	Weak recovery of subhypergraph is impossible
(a) and $\liminf_{N \rightarrow \infty} \frac{n^{(m-1)}E_Q\left(\frac{1}{n^{(m-1)} \log \frac{N}{n}}\right)}{\log N} > 1$	Exact recovery of subhypergraph is possible
(b) or $\limsup_{N \rightarrow \infty} \frac{n^{(m-1)}E_Q\left(\frac{1}{n^{(m-1)} \log \frac{N}{n}}\right)}{\log N} < 1$	Exact recovery of subhypergraph is impossible

In the special case where $P = \text{Bern}(p_1)$ and $Q = \text{Bern}(p_0)$ are both Bernoulli distributions, we are able to visualize the weak recovery region characterized by (p_1, p_0) for $m = 2, 3$ in Figure 1. Red (green) region indicates that weak recovery is impossible (possible). Clearly, the red region corresponds to $m = 3$ is smaller than the one corresponding to $m = 2$, indicating that detecting a subhypergraph is easier than detecting a graph. [24, 25] make a similar discovery in hypothesis testing scenario, i.e., testing the existence of subhypergraph is easier than testing the existence of a subgraph. Figure 2 displays the regions characterized by (p_1, p_0) for testing the existence of a subhypergraph for $m = 2, 3$; red (green) region indicates that a powerful test is impossible (possible). Interestingly, for fixed integer m , the red region in Figure 2 is smaller than the red region in Figure 1, indicating that hypothesis testing problem is generally easier to solve than detection problem.




FIGURE 1. Weak recovery regions in terms of (p_1, p_0) for $m = 2, 3$. Red: weak recovery is impossible; green: weak recovery is possible.

2. Main Results

Let $Z = (Z_1, \dots, Z_N)$ denote the membership vector of S^* , i.e., $Z_i = 1$ if $i \in S^*$ and $Z_i = 0$ otherwise. For \hat{S} an estimator of S^* , let \hat{Z} denote the membership vector of \hat{S} . We say that \hat{Z} is an *exact recovery* of Z if $\mathbb{P}(\hat{Z} \neq Z) = o(1)$; \hat{Z} is a *weak recovery* of Z if $\mathbb{P}(d_H(\hat{Z}, Z)/n) = o(1)$, where $d_H(\hat{Z}, Z)$ is the Hamming




FIGURE 2. Hypothesis testing regions in terms of (p_1, p_0) for $m = 2, 3$. Red: all tests are powerless; green: a powerful test exists.

distance between \widehat{Z} and Z . Obviously, when $n = O(1)$, weak recovery and exact recovery are equivalent. Let $L = \log \frac{dP}{dQ}$ and $\phi_Q(x) = \log \mathbb{E}_Q[e^{xL}]$. Moreover, assume that P and Q satisfy the following property:

$$\frac{d^2}{dx^2} \phi_Q(x) = O(\min \{D(P\|Q), D(Q\|P)\}), \quad x \in [-1, 1]. \quad (2)$$

This condition requires the log moment generating functions $\phi_Q(x)$ of L is bounded by the divergences on the interval $[-1, 1]$. The following theorem provides a sufficient condition under which weak recovery is possible.

Theorem 2.1 (Weak Recovery). *Suppose that P and Q satisfy (2). If*

$$n^{m-1}D(P\|Q) \rightarrow \infty, \quad \liminf_{N \rightarrow \infty} \frac{n^{(m)}D(P\|Q)}{n \log \frac{N}{n}} > 1, \quad (3)$$

then there exists an estimator \widehat{S} of S^* such that

$$\mathbb{P} \left[|\widehat{S} \Delta S^*| \leq 2n\epsilon_m \right] \geq 1 - e^{-\Omega(\frac{n}{\epsilon_m})}, \quad \text{where } \epsilon_m = \left(n^{(m-1)}D(P\|Q) \right)^{-\frac{1}{2}}. \quad (4)$$

Conversely, if there is an estimator \widehat{S} such that $\mathbb{E}[d_H(\widehat{Z}, Z)] = o(n)$, then

$$n^{m-1}D(P\|Q) \rightarrow \infty, \quad \liminf_{N \rightarrow \infty} \frac{n^{(m)}D(P\|Q)}{n \log \frac{N}{n}} \geq 1. \quad (5)$$

When $D(P\|Q)$ remains constant, condition (3) is weaker for $m = 3$ than $m = 2$. The later has been considered for weak recovery of subgraph by [13]. Besides, the error rate ϵ_m in (4) is smaller for $m = 3$ than $m = 2$. This indicates the significant difference between subhypergraph recovery and subgraph recovery.

Theorem 2.1 says that (3) are sufficient conditions for weak recovery. Meanwhile, (3) are almost necessary conditions for weak recovery, since if (5) fails, then weak recovery is impossible. These conditions are derived under model (1), and are dramatically different from the ones derived under stochastic block models ([7, 1, 19]) or generalized censored block models ([2]). The theorem holds for either $n = o(N)$ or $n \asymp N$, i.e., the cardinality of the underlying subhypergraph is either significantly smaller than N or of the same order as N .

Theorem 2.1 is established in the perspective of weak recovery, fundamentally different from the ones derived in hypothesis testing ([24]). To demonstrate the difference, consider unweighted hypergraphs with

$P = \text{Bern}(p_1)$ and $Q = \text{Bern}(p_0)$. Define $H_p(q) = q \log \frac{q}{p} + (1-q) \log \frac{1-q}{1-p}$ for $p, q \in (0, 1)$. Then it can be seen that $D(P\|Q) = H_{p_0}(p_1)$. By Theorem 2.1, we have the following corollary.

Corollary 2.2. *Suppose $P = \text{Bern}(p_1)$ and $Q = \text{Bern}(p_0)$, $\log \frac{p_1}{p_0}$ and $\log \frac{1-p_1}{1-p_0}$ are bounded. Then weak recovery is possible if*

$$n^{m-1} H_{p_0}(p_1) \rightarrow \infty, \quad \liminf_{N \rightarrow \infty} \frac{n^{(m)} H_{p_0}(p_1)}{n \log \frac{N}{n}} > 1. \quad (6)$$

Conversely, if weak recovery is possible, then

$$n^{m-1} H_{p_0}(p_1) \rightarrow \infty, \quad \liminf_{N \rightarrow \infty} \frac{n^{(m)} H_{p_0}(p_1)}{n \log \frac{N}{n}} \geq 1. \quad (7)$$

It might be interesting to compare the weak recovery condition (6) with the detection boundary condition in Theorem 3.1 of [24]. Assume $n = o(N)$ as in [24], then by Corollary 2.2, weak recovery is impossible if

$$\liminf_{N \rightarrow \infty} \frac{n^{(m)} H_{p_0}(p_1)}{n \log \frac{N}{n}} < 1. \quad (8)$$

Note that (8) is weaker than Condition (7) in [24]. Hence, the undetectable region derived in [24] is smaller than the weak recovery region for any integer $m \geq 2$. We visualize the regions in Figure 1 and Figure 2, which demonstrates the significant distinction between weak recovery limits and hypothesis testing.

The proof of sufficiency in Theorem 2.1 proceeds by showing the maximum likelihood estimator is a weak recovery estimator. Let

$$L_{i_1 \dots i_m} = \begin{cases} \frac{dP}{dQ}(A_{i_1 \dots i_m}), & \text{if } i_1, \dots, i_m \in \mathcal{V} \text{ are pairwise distinct,} \\ 0, & \text{otherwise.} \end{cases}$$

For any two subsets $S_1, S_2 \subset \mathcal{V}$, let

$$I_m(S_1, S_2) = \{(i_1, \dots, i_m) | i_1 < \dots < i_m, \{i_1, \dots, i_m\} \cap S_1 \neq \emptyset, \{i_1, \dots, i_m\} \cap S_2 \neq \emptyset, \{i_1, \dots, i_m\} \subset S_1 \cup S_2\}$$

and

$$L(S_1, S_2) = \sum_{(i_1, \dots, i_m) \in I_m(S_1, S_2)} L_{i_1 \dots i_m}.$$

Clearly, $I_m(S_1, S_2)$ includes the tuples (i_1, \dots, i_m) that intersects with both S_1 and S_2 whose entries are included in $S_1 \cup S_2$. Let \widehat{S}_{ML} be the maximum likelihood estimator of S^* defined as

$$\widehat{S}_{ML} = \arg \max_{S \subset [N], |S|=n} L(S, S). \quad (9)$$

Proof of Theorem 2.1. (Sufficiency). Let $\widehat{S} = \widehat{S}_{ML}$ and $R = |\widehat{S} \cap S^*|$. Then $|\widehat{S}| = |S^*| = n$ and $|\widehat{S} \Delta S^*| = 2(n - R)$. To prove (4), we only need to show $\mathbb{P}(R \leq (1 - \epsilon_m)n) \leq \exp\left(-\Omega\left(\frac{n}{\epsilon_m}\right)\right)$.

By condition (3), there exists a constant $\eta \in (0, 1)$ such that $K(m, n)D(P\|Q) \geq (1 - \eta) \log \frac{N}{n}$. Let $\theta = (1 - \eta)D(P\|Q)$. For $0 \leq r \leq n - 1$, we have

$$\begin{aligned} \{R = r\} &\subset \{\exists S \subset S^* : |S| = n - r, L(S, S^*) \leq (n^{(m)} - r^{(m)})\theta\} \\ &\cup \{\exists S \subset S^*, \exists T \subset (S^*)^c : |S| = |T| = n - r, L(T, T) + L(T, S^* \setminus S) \geq (n^{(m)} - r^{(m)})\theta\}. \end{aligned}$$

Note that $A_{i_1 \dots i_m} \sim P$ and $A_{i_1 \dots i_m} \sim Q$ for $\{i_1, \dots, i_m\} \subset S \subset S^*$ and $\{i_1, \dots, i_m\} \cap T \neq \emptyset$ respectively. Hence, we get

$$\begin{aligned}
& \mathbb{P}[R = r] \\
& \leq \binom{n}{n-r} \mathbb{P}\left(L(S, S^*) \leq (n^{(m)} - r^{(m)})\theta\right) \\
& \quad + \binom{n}{n-r} \binom{N-n}{n-r} \mathbb{P}\left(L(T, T) + L(T, S^* \setminus S) \geq (n^{(m)} - r^{(m)})\theta\right) \\
& \leq \left(\frac{ne}{n-r}\right)^{n-r} \exp\left(- (n^{(m)} - r^{(m)})E_P(\theta)\right) + \left(\frac{(N-n)ne^2}{(n-r)^2}\right)^{(n-r)} \exp\left(- (n^{(m)} - r^{(m)})E_Q(\theta)\right) \\
& \leq \exp\left(- (n-r) \left[K(m, n)E_P(\theta) - \log \frac{e}{\epsilon_m} \right]\right) + \exp\left(- (n-r) \left[K(m, n)E_Q(\theta) - \log \frac{(N-n)e^2}{N\epsilon_m^2} \right]\right),
\end{aligned}$$

here we used the fact that $\epsilon_m \leq \frac{n-r}{n}$ since $r \leq (1 - \epsilon_m)n$.

By Lemma 3.1, (13) and condition (3), for a constant c , we have

$$C_N := K(m, n)E_P(\theta) - \log \frac{e}{\epsilon_m} \geq c\eta^2 K(m, n)D(P\|Q) - \log \frac{e}{\epsilon_m} \asymp \frac{1}{\epsilon_m^2}.$$

By the fact that $E_Q(\theta) = E_P(\theta) + \theta$ and the definition of η , one has

$$\begin{aligned}
D_N & \equiv K(m, n)E_Q(\theta) - \log \frac{(N-n)e^2}{N\epsilon_m} \\
& \geq c\eta^2 K(m, n)D(P\|Q) - 2 \log \frac{e}{\epsilon_m} + (1 - \eta)K(m, n)D(P\|Q) - \log \frac{N-n}{n} \\
& \geq c\eta^2 K(m, n)D(P\|Q) - 2 \log \frac{e}{\epsilon_m} \asymp \frac{1}{\epsilon_m^2}.
\end{aligned}$$

Consequently,

$$\mathbb{P}[R \leq (1 - \epsilon_m)n] \leq \sum_{r=\epsilon_m n}^{\infty} \exp(-rC_N) + \sum_{r=\epsilon_m n}^{\infty} \exp(-rD_N) = \exp\left(-\Omega\left(\frac{n}{\epsilon_m}\right)\right).$$

(Necessity). For fixed indexes $i, j \in [N]$, let $Z_{ij} = \{Z_k : k \neq i, j\}$. If $Z_i = 0$, randomly and uniformly select a node from $\{j : Z_j = 1\}$ and denote it as J . If $Z_i = 1$, randomly and uniformly select a node from $\{j : Z_j = 0\}$ and denote it as J . Note that

$$\mathbb{P}(J = k | Z_i = 0) = \mathbb{P}(J = k | Z_i = 0, k \in \{j : Z_j = 1\}) \mathbb{P}(k \in \{j : Z_j = 1\} | Z_i = 0) = \frac{1}{N-1} = \mathbb{P}(J = k | Z_i = 1),$$

and $\mathbb{P}(Z_{iJ} | Z_i = 0, J) = \mathbb{P}(Z_{iJ} | Z_i = 1, J) = \binom{N-2}{n-1}^{-1}$. Then by the property of conditional probability, careful calculation yields

$$\begin{aligned}
& \frac{\mathbb{P}(J, Z_{iJ}, A | Z_i = 0)}{\mathbb{P}(J, Z_{iJ}, A | Z_i = 1)} = \frac{\mathbb{P}(A | Z_{iJ}, J, Z_i = 0)}{\mathbb{P}(A | Z_{iJ}, J, Z_i = 1)} \\
& = \frac{\prod_{\substack{i_2 < \dots < i_m \\ i, J \notin \{i_2, \dots, i_m\} \\ Z_{i_2} \dots Z_{i_m} = 1}} Q(A_{ii_2 \dots i_m}) \prod_{\substack{i_2 < \dots < i_m \\ i, J \notin \{i_2, \dots, i_m\} \\ Z_{i_2} \dots Z_{i_m} = 0}} Q(A_{ii_2 \dots i_m}) \prod_{\substack{i_2 < \dots < i_m \\ i, J \notin \{i_2, \dots, i_m\} \\ Z_{i_2} \dots Z_{i_m} = 1}} P(A_{Ji_2 \dots i_m}) \prod_{\substack{i_2 < \dots < i_m \\ i, J \notin \{i_2, \dots, i_m\} \\ Z_{i_2} \dots Z_{i_m} = 0}} Q(A_{Ji_2 \dots i_m})}{\prod_{\substack{i_2 < \dots < i_m \\ i, J \notin \{i_2, \dots, i_m\} \\ Z_{i_2} \dots Z_{i_m} = 1}} P(A_{ii_2 \dots i_m}) \prod_{\substack{i_2 < \dots < i_m \\ i, J \notin \{i_2, \dots, i_m\} \\ Z_{i_2} \dots Z_{i_m} = 0}} Q(A_{ii_2 \dots i_m}) \prod_{\substack{i_2 < \dots < i_m \\ i, J \notin \{i_2, \dots, i_m\} \\ Z_{i_2} \dots Z_{i_m} = 1}} Q(A_{Ji_2 \dots i_m}) \prod_{\substack{i_2 < \dots < i_m \\ i, J \notin \{i_2, \dots, i_m\} \\ Z_{i_2} \dots Z_{i_m} = 0}} Q(A_{Ji_2 \dots i_m})}
\end{aligned}$$

$$\begin{aligned}
& \frac{\prod_{\substack{i_1 < i_2 < \dots < i_m \\ i, J \notin \{i_1, i_2, \dots, i_m\} \\ Z_{i_1} \dots Z_{i_m} = 1}} P(A_{i_1 i_2 \dots i_m})}{\prod_{\substack{i_1 < i_2 < \dots < i_m \\ i, J \notin \{i_1, i_2, \dots, i_m\} \\ Z_{i_1} \dots Z_{i_m} = 1}} P(A_{i_1 i_2 \dots i_m})} \times \frac{\prod_{\substack{i_1 < i_2 < \dots < i_m \\ i, J \notin \{i_1, i_2, \dots, i_m\} \\ Z_{i_1} \dots Z_{i_m} = 0}} Q(A_{i_1 i_2 \dots i_m})}{\prod_{\substack{i_1 < i_2 < \dots < i_m \\ i, J \notin \{i_1, i_2, \dots, i_m\} \\ Z_{i_1} \dots Z_{i_m} = 0}} Q(A_{i_1 i_2 \dots i_m})} \frac{\prod_{i, J \in \{i_1, \dots, i_m\}} Q(A_{i_1 i_2 \dots i_m})}{\prod_{i, J \in \{i_1, \dots, i_m\}} Q(A_{i_1 i_2 \dots i_m})} \\
& = \prod_{\substack{i_2 < \dots < i_m \\ i, J \notin \{i_2, \dots, i_m\} \\ Z_{i_2} \dots Z_{i_m} = 1}} \frac{Q(A_{i_2 \dots i_m}) P(A_{J i_2 \dots i_m})}{P(A_{i_2 \dots i_m}) Q(A_{J i_2 \dots i_m})}.
\end{aligned}$$

Let $T_{iJ} = \{A_{i_2 \dots i_m}, A_{J i_2 \dots i_m} : i_2 < \dots < i_m, i, J \notin \{i_2, \dots, i_m\}, Z_{i_2} \dots Z_{i_m} = 1\}$. For $Z_i = 0$, $T_{iJ} \sim Q^{\otimes (n-1)^{(m-1)}} \otimes P^{\otimes (n-1)^{(m-1)}}$ and $T_{iJ} \sim P^{\otimes (n-1)^{(m-1)}} \otimes Q^{\otimes (n-1)^{(m-1)}}$ if $Z_i = 1$. Then by a similar argument as in proof of Theorem 1 in [13], we conclude $(n-1)^{(m-1)} D(P\|Q) \rightarrow \infty$ and $\liminf_{N \rightarrow \infty} \frac{n^{(m)} D(P\|Q)}{n \log \frac{N}{n}} \geq 1$. \square

Next, we derive sufficient and necessary conditions for exact recovery of a subhypergraph. Let $E_Q(t) = \sup_{\lambda \in \mathbb{R}} (\lambda t - \phi_Q(\lambda))$ be the Fenchel conjugate of ϕ_Q . Obviously, $E_P(t) = \sup_{\lambda \in \mathbb{R}} (\lambda t - \phi_P(\lambda)) = E_Q(t) - t$. The following theorem presents necessary and sufficient conditions for exact recovery of a subhypergraph.

Theorem 2.3 (Exact Recovery). *Suppose P and Q satisfy (2). If (3) and the following hold*

$$\liminf_{N \rightarrow \infty} \frac{n^{(m-1)} E_Q\left(\frac{1}{n^{(m-1)}} \log \frac{N}{n}\right)}{\log N} > 1, \quad (10)$$

then there exists an estimator \tilde{S} of S^* such that $\mathbb{P}(\tilde{S} = S^*) = 1 + o(1)$. Conversely, if there exists an estimator \tilde{S} of S^* such that $\mathbb{P}(\tilde{S} = S^*) = 1 + o(1)$, then (5) and the following hold

$$\liminf_{N \rightarrow \infty} \frac{n^{(m-1)} E_Q\left(\frac{1}{n^{(m-1)}} \log \frac{N}{n}\right)}{\log N} \geq 1, \quad (11)$$

Theorem 2.3 says that the conditions (3) and (10) are nearly sharp for exact recovery. Again, these conditions are fundamentally different from the ones derived in the literature of stochastic block models ([1, 19]) or generalized censored block models ([2]). In particular, exact recovery in [19] only requires a condition similar to (3) up to some constant. In contrast, our exact recovery conditions requires additionally (10). For $m = 2$, conditions (10) and (11) degenerate to Theorem 2 of [13].

The proof of Theorem 2.3 heavily relies on Theorem 2.1. Specifically, under (2) and (3), one can construct a provably exact recovery estimator \tilde{S} based on weak recovery. Specifically, one construct \tilde{S} as follows.

- For given n, N and P, Q, A , fix a small constant $\delta \in (0, 1)$ such that $\delta N, \frac{1}{\delta}$ are integers.
- Divide the vertex set $[N]$ into subsets C_k with $|C_k| = \delta N$ for $k = 1, 2, \dots, \frac{1}{\delta}$.
- For $k = 1, 2, \dots, \frac{1}{\delta}$, let A_k be the subhypergraph on $[N] \setminus C_k$ with $(1 - \delta)N$ nodes and subhypergraph of size $(1 - \delta)n$. Let \hat{S}_k be the weak recovery estimator based on A_k .
- Let $r_i = \sum_{\substack{i \notin \{i_2, \dots, i_m\} \subset \hat{S}_k \\ i_2 < \dots < i_m}} L_{i i_2 \dots i_m}$ for $i \in C_k$ and $k = 1, 2, \dots, \frac{1}{\delta}$. The exact recovery estimator \tilde{S} consists of n nodes with the largest r_i .

Proof of Theorem 2.3. (Sufficiency). When $n = O(1)$, exact recover is the same as weak recovery. Hence we assume $n \rightarrow \infty$. By Lemma 3.6, we only need to verify (15). Condition (3) implies that

$$\liminf_{N \rightarrow \infty} \frac{((1 - \delta)n)^{(m)} D(P\|Q)}{(1 - \delta)n \log \frac{(1 - \delta)N}{(1 - \delta)n}} = (1 + o(1))(1 - \delta)^{m-1} \liminf_{N \rightarrow \infty} \frac{n^{(m)} D(P\|Q)}{n \log \frac{N}{n}} > 1,$$

for small enough fixed constant $\delta \in (0, 1)$ and large n . Then as argued in the proof of Theorem 2 of [13], Lemma 3.5 holds. Since $\epsilon_m = o(1)$ in Lemma 3.5 holds. Then (15) holds.

(Necessity). Obviously (5) holds. Let $\gamma_m = \frac{1}{n^{(m-1)}} \log \frac{N}{n}$. Then $\gamma_m < \frac{D(P\|Q)}{m(1+\epsilon_0)}$ for a small constant $\epsilon_0 > 0$. In the following, we only need to prove (11) holds. We will prove (11) by contradiction. If (11) is not true, then

$$\limsup_{N \rightarrow \infty} \frac{n^{(m-1)} E_Q(\gamma_m)}{\log N} < 1. \quad (12)$$

By Lemma 3.8, we show there is no θ_n satisfies (16) and (17) by showing $\theta_n = \gamma_m$ doesn't satisfy (16) or (17). There exists a small constant $\tau > 0$ such that γ_m and $\gamma_m + \tau D(Q\|P)$ are in $[-D(Q\|P), D(P\|Q)]$. Note that for small τ , $(n-1)^{(m-1)} E_Q(\gamma_m + \tau D(Q\|P)) < (1-2\tau) \log N$ by (12) and Lemma 2 in [13]. Hence, by Lemma 3.3, we have

$$\begin{aligned} Q \left(\sum_{1 \leq i_2 < \dots < i_m \leq n-1} L_{i_1 i_2 \dots i_m} \geq (n-1)^{(m-1)} \gamma_m \right) &\geq \exp \left(- \frac{(n-1)^{(m-1)} E_Q(\gamma_m + \tau D(Q\|P)) + \log 2}{1 - \frac{C}{(n-1)^{(m-1)} \tau^2 \min\{D(Q\|P), D(P\|Q)\}}} \right) \\ &\geq \frac{1}{n^{1-\tau}}. \end{aligned}$$

Then γ_m does not satisfy (17).

Let $n_1 = \frac{n}{\log n}$ and

$$\tau_1 := \frac{(n_1 - 1)^{(m-1)} D(P\|Q) + [(n - n_1)^{(m-1)} - (n - 1)^{(m-1)}] \gamma_m + 6\sigma}{(n - n_1)^{(m-1)} D(P\|Q)} = o(1).$$

Take small $\tau > 0$ such that $\gamma_m - \tau_1 D(P\|Q)$ and $\gamma_m - (\tau_1 + \tau) D(P\|Q)$ are in $[-D(Q\|P), D(P\|Q)]$. By (12) and $E_P(\gamma_m) = E_Q(\gamma_m) - \gamma$, for small τ ,

$$(n - n_1)^{(m-1)} E_P(\gamma_m - (\tau_1 + \tau) D(Q\|P)) \leq (1 - 2\tau_2) \log n,$$

with a small constant $\tau_2 > 0$. Then by Lemma 3.3, it follows

$$\begin{aligned} &P \left(\sum_{1 \leq i_2 < \dots < i_m \leq n - n_1} L_{i_1 i_2 \dots i_m} \leq (n - 1)^{(m-1)} \gamma_m - (n_1 - 1)^{(m-1)} D(P\|Q) - 6\sigma \right) \\ &= P \left(\sum_{1 \leq i_2 < \dots < i_m \leq n - n_1} L_{i_1 i_2 \dots i_m} \leq (n - n_1)^{(m-1)} (\gamma_m - \tau_1 D(P\|Q)) \right) \\ &\geq \exp \left(- \frac{(n - n_1)^{(m-1)} E_P(\gamma_m - (\tau_1 + \tau) D(Q\|P)) + \log 2}{1 - \frac{C}{(n - n_1)^{(m-1)} \tau^2 \min\{D(Q\|P), D(P\|Q)\}}} \right) \geq \frac{1}{n^{1-\tau_2}}. \end{aligned}$$

Hence γ_m does not satisfy (16). □

3. Proof of Additional Lemmas

3.1. Lemmas for weak recovery

Lemma 3.1. *For any fixed integer $m \geq 2$, the following equations hold.*

$$K(m, n) := \min_{\text{integer } r: 0 \leq r \leq n-1} \frac{n^{(m)} - r^{(m)}}{n - r} = (1 + o(1)) \frac{n^{(m)}}{n},$$

$$\max_{\text{integer } r: 0 \leq r \leq n-1} \frac{n^{(m)} - r^{(m)}}{n - r} = (1 + o(1)) (n - 1)^{(m-1)}.$$

Proof. Define a function $f(x) = x(x-1)\dots(x-m+1)$. The derivative of $f(x)$ is equal to

$$f'(x) = \sum_{i=0}^{m-1} x(x-1)\dots(x-i)(x-i-2)\dots(x-m+1).$$

Then $f'(x) \geq 0$ for $x \geq m-1$ and $f'(x)$ is increasing in x . Note that $\frac{n^{(m)} - r^{(m)}}{n-r} = \frac{1}{m!} \frac{f(n) - f(r)}{n-r}$. Let $g(x) = \frac{1}{m!} \frac{f(n) - f(x)}{n-x}$. Then

$$g'(x) = \frac{1}{m!} \frac{f(n) - f(x) - (n-x)f'(x)}{(n-x)^2}, \quad x \leq n-1.$$

By the Lagrange mean value theorem, for $m-1 \leq x \leq n-1$, we have

$$\frac{f(n) - f(x)}{n-x} = f'(x^*) \geq f'(x) \geq 0, \quad x^* \in (x, n).$$

Hence, $g'(x) \geq 0$ for $x \geq m-1$ and hence $g(r)$ is increasing in $r \geq m-1$. The minimum can only be achieved at $r \in [0, m-1]$. In this case, $\min_{0 \leq r \leq m-1} g(r) = \frac{n^{(m)}}{n} (1 + o(1))$. The maximum is achieved at $r = n-1$. By Pascal's rule, we have $n^{(m)} = (n-1)^{(m-1)} + (n-1)^{(m)}$. Then the desired result follows. \square

In the following, we present several lemmas from [13].

Lemma 3.2 ([13]). *Condition (2) implies the following*

$$E_P((1-\eta)D(P\|Q)) \geq \frac{\eta^2}{2C} D(P\|Q), \quad (13)$$

$$E_Q(-(1-\eta)D(Q\|P)) \geq \frac{\eta^2}{2C} D(Q\|P), \quad (14)$$

for constant C and $\eta \in [0, 1]$ and $D(P\|Q) \asymp D(Q\|P)$.

Lemma 3.3 ([13]). *Suppose condition (2) holds. For $-D(Q\|P) \leq \gamma < \gamma + \delta \leq D(P\|Q)$,*

$$\exp(-nE_Q(\gamma)) \geq Q\left(\sum_{i=1}^n L_i > n\gamma\right) \geq \exp\left(-\frac{nE_Q(\gamma + \delta) + \log 2}{1 - \frac{C \min\{D(Q\|P), D(P\|Q)\}}{n\delta^2}}\right),$$

where $L_i = \frac{dP}{dQ}(X_i)$ with independent $X_i \sim Q$.

For later use, we assume $|S^*|$ is random and unknown and present the following lemmas.

Lemma 3.4. *If $|n_1 - n| \leq \frac{n}{\log n}$, then*

$$\min_{\text{integer } r: 0 \leq r \leq n_1-1} \frac{n_1^{(m)} - r^{(m)}}{n_1 - r} = (1 + o(1)) \frac{n^{(m)}}{n}, \quad \max_{0 \leq r \leq (1-\epsilon_m)n - \frac{n}{\log n}} \left| 1 - \frac{n_1 - r}{n - r} \right| = o(1),$$

$$\min_{\text{integer } r: 0 \leq r \leq n_1-1} \frac{n^{(m)} - r^{(m)}}{n_1^{(m)} - r^{(m)}} \geq \frac{1}{m} (1 + o(1)),$$

with $\epsilon_m = (\min\{\log n, n^{m-1}D(P\|D)\})^{-\frac{1}{2}}$.

Proof. By Lemma 3.1 and $|\frac{n_1}{n} - 1| \leq \frac{1}{\log n} = o(1)$, we have

$$\min_{\text{integer } r: 0 \leq r \leq n_1 - 1} \frac{n_1^{(m)} - r^{(m)}}{n_1 - r} = (1 + o(1)) \frac{n_1^{(m)}}{n_1} = (1 + o(1)) \frac{n}{n_1} \frac{n_1^{(m)}}{n^{(m)}} = (1 + o(1)) \frac{n^{(m)}}{n}.$$

Note that $0 \leq \frac{r}{n} \leq 1 - \epsilon_m - \frac{1}{\log n}$. Then

$$\begin{aligned} \max_{0 \leq r \leq (1 - \epsilon_m)n - \frac{n}{\log n}} \left| 1 - \frac{n_1 - r}{n - r} \right| &\leq \max_{0 \leq r \leq (1 - \epsilon_m)n - \frac{n}{\log n}} \frac{n}{n - r} \frac{1}{\log n} = \max_{0 \leq r \leq (1 - \epsilon_m)n - \frac{n}{\log n}} \frac{1}{1 - \frac{r}{n}} \frac{1}{\log n} \\ &\leq \frac{1}{\epsilon_m \log n + 1} = o(1). \end{aligned}$$

By Lemma 3.1, the following holds uniformly for all $r \leq (1 - \epsilon_m)n - \frac{n}{\log n}$,

$$\frac{n^{(m)} - r^{(m)}}{n_1^{(m)} - r^{(m)}} = \frac{n^{(m)} - r^{(m)}}{n - r} \left(\frac{n_1^{(m)} - r^{(m)}}{n_1 - r} \right)^{-1} (1 + o(1)) \geq (1 + o(1)) \frac{n^{(m)}}{n} \left((n_1 - 1)^{(m-1)} \right)^{-1} = (1 + o(1)) \frac{1}{m}.$$

□

Lemma 3.5. *Suppose Condition (2) holds, $n \rightarrow \infty$, $\limsup_{N \rightarrow \infty} n/N < 1$. If condition (3) holds and*

$$\mathbb{P} \left(\left| |S^*| - n \right| \leq \frac{n}{\log n} \right) = 1 - o(1),$$

then

$$\mathbb{P} \left(|\widehat{S}_{ML} \Delta S^*| \leq 2n\epsilon_m + 3\frac{n}{\log n} \right) = 1 - o(1), \quad \epsilon_m = (\min\{\log n, n^{m-1}D(P\|D)\})^{-\frac{1}{2}}.$$

Proof. Let $n_1 = |S^*|$. Then $|n_1 - n| \leq \frac{n}{\log n}$ with probability tending to 1. Let $\widehat{S} = \widehat{S}_{ML}$ and $R = |\widehat{S} \cap S^*|$. Then $|\widehat{S} \Delta S^*| = n_1 + n - 2R$. To prove (4), we only need to show $\mathbb{P} \left(R \leq (1 - \epsilon_m)n - \frac{n}{\log n} \right) = o(1)$.

By condition (3), there exists a constant $\eta \in (0, 1)$ such that $K(m, n)D(P\|Q) \geq (1 - \eta) \log \frac{N}{n}$. Let $\theta = (1 - \eta)D(P\|Q)$. For $0 \leq r \leq n - 1$, we have

$$\begin{aligned} \{R = r\} &\subset \{ \exists S \subset S^* : |S| = n_1 - r, L(S, S^*) \leq (n^{(m)} - r^{(m)})\theta \} \\ &\cup \{ \exists S \subset S^*, \exists T \subset (S^*)^c : |S| = n_1 - r, |T| = n - r, L(T, T) + L(T, S^* \setminus S) \geq (n^{(m)} - r^{(m)})\theta \}. \end{aligned}$$

Hence, by Lemma 3.4, we get

$$\begin{aligned} \mathbb{P}[R = r] &\leq \left(\frac{n_1 e}{n_1 - r} \right)^{n_1 - r} \exp \left(-(n_1^{(m)} - r^{(m)}) E_P \left(\frac{n^{(m)} - r^{(m)}}{n_1^{(m)} - r^{(m)}} \theta \right) \right) \\ &\quad + \left(\frac{n_1 e}{n_1 - r} \right)^{n_1 - r} \left(\frac{(N - n_1) e}{n - r} \right)^{n - r} \exp \left(-(n^{(m)} - r^{(m)}) E_Q(\theta) \right) \\ &\leq \exp \left(-(n - r)(1 + o(1)) \left[K(m, n) E_P \left(\frac{n^{(m)} - r^{(m)}}{n_1^{(m)} - r^{(m)}} \theta \right) - \log \frac{e}{\epsilon_m} \right] \right) \\ &\quad + \exp \left(-(n - r)(1 + o(1)) \left[K(m, n) E_Q(\theta) - \log \frac{(N - n_1) e^2}{n \epsilon_m^2} \right] \right), \end{aligned}$$

By Lemma 3.1, Lemma 3.4, (13) and condition (3), for a constant $C > 0$, we have

$$C_N := K(m, n) E_P \left(\frac{n^{(m)} - r^{(m)}}{n_1^{(m)} - r^{(m)}} \theta \right) - \log \frac{e}{\epsilon_m} \geq (1 + o(1)) \frac{C}{m} n^{m-1} D(P\|D) - \log \frac{e}{\epsilon_m} \rightarrow \infty.$$

By the fact that $E_Q(\theta) = E_P(\theta) + \theta$, one has

$$\begin{aligned} D_N &\equiv K(m, n)E_Q(\theta) - \log \frac{(N - n_1)e^2}{n\epsilon_m^2} \\ &\geq c\eta^2 K(m, n)D(P\|Q) - 2\log \frac{e}{\epsilon_m} + (1 - \eta)\frac{n^{(m)}}{n}D(P\|Q) - \log \frac{N - n_1}{n} \\ &\geq c\eta^2 n^{m-1}D(P\|Q) - 2\log \frac{e}{\epsilon_m} \rightarrow \infty. \end{aligned}$$

Then the proof is complete as in the proof of Theorem 2.1. \square

3.2. Lemmas for exact recovery

Lemma 3.6. Let $S_k^* = ([N] \setminus C_k) \cap S^*$ for $1 \leq k \leq \delta^{-1}$. Suppose condition (2) holds and

$$\mathbb{P} \left[|\widehat{S}_k \Delta S_k^*| \leq \delta n, 1 \leq k \leq \frac{1}{\delta} \right] = 1 + o(1). \quad (15)$$

Then $\mathbb{P}(\widetilde{S} = S^*) = 1 + o(1)$.

Proof. Fix $i_1 > (1 - \delta)n$ and let $X_{i_1 i_2 \dots i_m} = \frac{dP}{dQ}(A_{i_1 i_2 \dots i_m})$ and $Y_{i_1 i_2 \dots i_m} = \frac{dP}{dQ}(A_{i_1 i_2 \dots i_m})$ be random variables under P and Q respectively for $i_2 < \dots < i_m$. By a similar argument as in the proof of Theorem 3 in [13], r_i is stochastically greater than or equal to $\sum_{1 \leq i_2 < \dots < i_m \leq (1-2\delta)n} X_{i_1 i_2 \dots i_m} + \sum_{1 \leq i_2 < \dots < i_m \leq \delta n} Y_{i_1 i_2 \dots i_m}$ for $i \in S^*$. For $i \in [N] \setminus S^*$, r_i is identically distributed as $\sum_{1 \leq i_2 < \dots < i_m \leq (1-\delta)n} Y_{i_1 \dots i_m}$. By Lemma 3.7 below and union bound, $r_i \geq ((1 - \delta)n)^{(m-1)}\gamma_m$, $i \in S^*$ and $r_i < ((1 - \delta)n)^{(m-1)}\gamma_m$, $i \in [N] \setminus S^*$ with probability $1 + o(1)$. Then $\mathbb{P}(\widetilde{S} = S^*) = 1 + o(1)$. \square

Lemma 3.7. Suppose Condition (2) holds and let $\gamma_m = \frac{1}{n^{(m-1)}} \log \frac{N}{n}$. Fix $i_1 > (1 - \delta)n$ and let $X_{i_1 \dots i_m} = \frac{dP}{dQ}(A_{i_1 \dots i_m})$ and $Y_{i_1 \dots i_m} = \frac{dP}{dQ}(A_{i_1 \dots i_m})$ be random variables under P and Q respectively for $i_2 < \dots < i_m$. Then for small $\delta \in (0, 1)$, we have

$$\begin{aligned} \mathbb{P} \left(\sum_{1 \leq i_2 < \dots < i_m \leq (1-2\delta)n} X_{i_1 \dots i_m} + \sum_{1 \leq i_2 < \dots < i_m \leq \delta n} Y_{i_1 \dots i_m} \leq ((1 - \delta)n)^{(m-1)}\gamma_m \right) &= o\left(\frac{1}{n}\right), \\ \mathbb{P} \left(\sum_{1 \leq i_2 < \dots < i_m \leq (1-\delta)n} Y_{i_1 \dots i_m} \geq ((1 - \delta)n)^{(m-1)}\gamma_m \right) &= o\left(\frac{1}{N - n}\right). \end{aligned}$$

Proof. Condition (10) implies $n^{(m-1)}E_Q(\gamma_m) \geq (1 + \epsilon) \log N$ for a small constant $\epsilon > 0$. Then

$$\begin{aligned} \mathbb{P} \left(\sum_{1 \leq i_2 < \dots < i_m \leq (1-\delta)n} Y_{i_1 \dots i_m} \geq ((1 - \delta)n)^{(m-1)}\gamma_m \right) &\leq \exp\left(- (1 - \delta)n^{(m-1)}E_Q(\gamma_m)\right) \\ &\leq \exp\left(- (1 + o(1))(1 - \delta)^{m-1}(1 + \epsilon) \log N\right) = N^{-(1-\delta)^{m-1}(1+\epsilon)(1+o(1))} = o\left(\frac{1}{N - n}\right), \end{aligned}$$

for $\delta < 1 - (1 + \epsilon)^{-\frac{1}{m-1}}$.

By a similar argument as in the proof of Lemma 5 in [13] and for a constant $C > 0$, we have

$$\begin{aligned}
& \mathbb{P} \left(\sum_{1 \leq i_2 < \dots < i_m \leq (1-2\delta)n} X_{i_1 \dots i_m} + \sum_{1 \leq i_2 < \dots < i_m \leq \delta n} Y_{i_1 \dots i_m} \leq ((1-\delta)n)^{(m-1)} \gamma_m \right) \\
& \leq \exp \left(-(1-2\delta)n^{(m-1)} E_P(\gamma_m) + C(\delta n)^{(m-1)} E_Q(\gamma_m) + (\delta n)^{(m-1)} \gamma_m \right) \\
& \leq \exp \left\{ \left[-n^{(m-1)} E_P(\gamma_m) \left[(1-2\delta)^{m-1} - C\delta^{m-1} \right] + (1+C)\delta^{m-1} \log N \right] (1+o(1)) \right\} \\
& \leq \exp \left\{ \left[-(\log n + \epsilon \log N) \left[(1-2\delta)^{m-1} - C\delta^{m-1} \right] + (1+C)\delta^{m-1} \log N \right] (1+o(1)) \right\} \\
& = \frac{1}{n^{(1-2\delta)^{m-1} - C\delta^{m-1}}} \frac{1}{N^{\epsilon((1-2\delta)^{m-1} - C\delta^{m-1}) - (1+C)\delta^{m-1}}} = o\left(\frac{1}{n}\right),
\end{aligned}$$

if δ satisfies $\frac{1+(1+C)\delta^{m-1}}{(1-2\delta)^{m-1} - C\delta^{m-1}} - 1 < \epsilon$. This is possible since

$$\frac{1+(1+C)\delta^{m-1}}{(1-2\delta)^{m-1} - C\delta^{m-1}} > 1 \quad \text{and} \quad \lim_{\delta \rightarrow 0^+} \frac{1+(1+C)\delta^{m-1}}{(1-2\delta)^{m-1} - C\delta^{m-1}} = 1.$$

□

Lemma 3.8. *Suppose $n \rightarrow \infty$ and $\limsup n/N < 1$. Fix $i_1 > n$ and let $L_{i_1 \dots i_m} = \frac{dP}{dQ}(A_{i_1 \dots i_m})$. If there exists an estimator \hat{S} such that $\mathbb{P}(\hat{S} = S^*) = 1 + o(1)$, then for $n_1 \rightarrow \infty$ and $n_1 = o(n)$, there exists a sequence θ_n such that*

$$P \left(\sum_{1 \leq i_2 < \dots < i_m \leq n-n_1} L_{i_1 i_2 \dots i_m} \leq (n-1)^{(m-1)} \theta_n - (n_1-1)^{(m-1)} D(P\|Q) - 6\sigma \right) \leq \frac{2}{n_1}, \quad (16)$$

$$Q \left(\sum_{1 \leq i_2 < \dots < i_m \leq n-1} L_{i_1 i_2 \dots i_m} \geq (n-1)^{(m-1)} \theta_n \right) \leq \frac{1}{N-n}. \quad (17)$$

Here $\sigma^2 = n_1^{(m-1)} \text{Var}_P(L_{1 \dots m})$ with $\text{Var}_P(L_{1 \dots m})$ the variance of $L_{1 \dots m}$ under P .

Proof. Let \hat{S} be the MLE and $S^* = [n]$. For $i \in S^*$, let $i_0 = \arg \min_{i \in S^*} L(i, S^*)$. Let $F = \{\min_{i \in S^*} L(i, S^*) \leq \max_{i \notin S^*} L(i, S^* \setminus \{i_0\})\}$,

$$\theta_{1n} = \inf_{x \in \mathbb{R}} \left\{ P \left(\sum_{1 \leq i_2 < \dots < i_m \leq n-n_1} L_{i_1 i_2 \dots i_m} \leq (n-1)^{(m-1)} x - (n_1-1)^{(m-1)} D(P\|Q) - 6\sigma \right) \geq \frac{2}{n_1} \right\},$$

$$\theta_{2n} = \sup_{x \in \mathbb{R}} \left\{ Q \left(\sum_{1 \leq i_2 < \dots < i_m \leq n-1} L_{i_1 i_2 \dots i_m} \geq (n-1)^{(m-1)} x \right) \geq \frac{1}{N-n} \right\}.$$

Define $E_1 = \{\min_{i \in S^*} L(i, S^*) \leq (n-1)^{(m-1)} \theta_{1n}\}$ and $E_2 = \{\max_{i \notin S^*} L(i, S^* \setminus \{i_0\}) \geq (n-1)^{(m-1)} \theta_{2n}\}$.

Suppose $\mathbb{P}(E_1) \geq c_1 > 0$ and $\mathbb{P}(E_2) \geq c_2 > 0$ for two constants $c_1, c_2 \in (0, 1)$. Note that E_1 and E_2 are independent. Hence, as argued in Lemma 6 of [13], we have

$$\mathbb{P}(\theta_{1n} > \theta_{2n}) \geq \mathbb{P}(E_1 \cap E_2 \cap F^c) \geq \mathbb{P}(E_1)\mathbb{P}(E_2) - o(1) \geq c_1 c_2 - o(1) > 0.$$

Since θ_{1n} and θ_{2n} are deterministic, $\theta_{1n} > \theta_{2n}$ holds for large n . Let $\theta_n = \frac{\theta_{1n} + \theta_{2n}}{2}$. Then (16) and (17) hold.

In the following, we prove $\mathbb{P}(E_1) \geq c_1 > 0$ and $\mathbb{P}(E_2) \geq c_2 > 0$ for two constants $c_1, c_2 \in (0, 1)$. By the right-continuity of Q , we have

$$Q \left(\sum_{1 \leq i_2 < \dots < i_m \leq n-1} L_{i_1 i_2 \dots i_m} \geq (n-1)^{(m-1)} \theta_{2n} \right) \geq \frac{1}{N-n}.$$

Then the fact that $1 - x \leq e^{-x}$ yields

$$\begin{aligned} \mathbb{P}(E_2) &= 1 - \prod_{i \notin S^*} \mathbb{P}\left(L(i, S^* \setminus \{i_0\}) < (n-1)^{(m-1)}\theta_{2n}\right) \\ &= 1 - \left(1 - Q\left(\sum_{1 \leq i_2 < \dots < i_m \leq n-1} L_{i_1 i_2 \dots i_m} \geq (n-1)^{(m-1)}\theta_{2n}\right)\right)^{N-n} \geq 1 - \frac{1}{e}. \end{aligned}$$

Hence $\mathbb{P}(E_2) \geq c_2 > 0$ for constant $c_2 \in (0, 1)$. By the right-continuity of P , we have

$$P\left(\sum_{1 \leq i_2 < \dots < i_m \leq n-n_1} L_{i_1 i_2 \dots i_m} \leq (n-1)^{(m-1)}\theta_{1n} - (n_1-1)^{(m-1)}D(P\|Q) - 6\sigma\right) \geq \frac{2}{n_1}.$$

Let $T = [n_1]$ and $T_1 = \{i \in T : L(i, T) \leq (n_1-1)^{(m-1)}D(P\|Q) + 6\sigma\}$. Then $\mathbb{P}(|T_1| \geq \frac{n_1}{2}) = 1 + o(1)$ by a similar argument as in Lemma 6 of [13]. Hence,

$$\begin{aligned} \mathbb{P}(E_1) &\geq \mathbb{P}\left\{\min_{i \in T_1} L(i, S^* \setminus T) \leq (n-1)^{(m-1)}\theta_{1n} - (n_1-1)^{(m-1)}D(P\|Q) - 6\sigma\right\} \\ &\geq 1 - \mathbb{P}\left(|T_1| \geq \frac{n_1}{2}\right) \mathbb{P}\left(\prod_{i \in T_1} L(i, S^* \setminus T) \geq (n-1)^{(m-1)}\theta_{1n} - (n_1-1)^{(m-1)}D(P\|Q) - 6\sigma \mid |T_1| \geq \frac{n_1}{2}\right) \\ &\quad - \mathbb{P}\left(|T_1| < \frac{n_1}{2}\right) \\ &\geq 1 - \exp\left(-\mathbb{P}\left(\sum_{1 \leq i_2 < \dots < i_m \leq n-n_1} L_{i_1 i_2 \dots i_m} < (n-1)^{(m-1)}\theta_{1n} - (n_1-1)^{(m-1)}D(P\|Q) - 6\sigma\right) \frac{n_1}{2}\right) \\ &\quad - o(1) \geq 1 - \frac{1}{e} - o(1). \end{aligned}$$

Then proof is completed. \square

References

- [1] Ahn, K., Lee, K. and Suh, C. (2018). Hypergraph Spectral Clustering in the Weighted Stochastic Block Model. *IEEE Journal of Selected Topics in Signal Processing*, 12(5), 2018.
- [2] Ahn, K., Lee, K. and Suh, C. (2019). Community Recovery in Hypergraphs. *IEEE Transactions on Information Theory*, 12(5), 6561-6578.
- [3] Anderson, R. and Chellapilla, K. 2009. Finding dense subgraphs with size bounds. *WAW 2009*, 25-37.
- [4] Bhangale, A., Gandhi, R. and Kortsarz, G. (2017). Improved approximation algorithm for the dense-3-subhypergraph problem. <https://arxiv.org/abs/1704.08620>
- [5] Charikar, M. 2000. Greedy approximation algorithms for finding dense components in a graph. *APPROX*, 84-95.
- [6] Chiplunkar, A., Kapralov, M., Khanna, S., Mousavifar, A. and Peres, Y.(2018). Testing graph clusterability: algorithms and lower bounds. *2018 IEEE 59th Annual Symposium on Foundations of Computer Science*, 497-508.
- [7] Chien, I., Lin, C. and Wang, I.(2018). Community Detection in Hypergraphs: Optimal Statistical Limit and Efficient Algorithms. *Proceedings of the Twenty-First International Conference on Artificial Intelligence and Statistics*, 84:871-879.

- [8] Chen, J. and B. Yuan. 2006. Detecting functional modules in the yeast proteinprotein interaction network. *Bioinformatics*, **22(18)**, 2283-2290.
- [9] Chen, J. and Y. Saad. 2010. Dense subgraph extraction with application to community detection, *IEEE Transactions on Knowledge and Data Engineering* 24,7: 1216-1230.
- [10] Chertok, M. and Keller, Y. (2010). Efficient high order matching. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, **32(12)**, 2205-2215.
- [11] Chlamtac, E., M. Dinitz, C. Konrad, G. Kortsarz, and G. Rabanca. The Densest k-Subhypergraph problem. *SIAM Journal on Discrete Mathematics*. 32 (2):1458–77.
- [12] Gibson, D., R. Kumar, and A. Tomkins. 2005. Discovering large dense subgraphs in massive graphs. *VLDB 05*,721–732.
- [13] Hajek, B., Wu, Y. and Xu, J.(2017). Information limits for recovering a hidden community. *IEEE Transaction on Information Theory*, 63(8): 4729-4745.
- [14] Hajek, B., Wu, Y. and Xu, J. 2018. Recovering a hidden community beyond the Kesten Stigum threshold in $O(|E| \log |V|)$ time. *Journal of Applied Probability*, 55, 2: 325-352.
- [15] Hooi,B., H.A., Song, A. Beutel, N. Shah, K. Shin,and C. Faloutsos. 2016. Fraudar:bounding graph fraud in the face of camouflage, *KDD,ACM*,895-904.
- [16] Hu, S., Wu, X. and Chan, T-H. (2017). Maintaining densest subsets efficiently in evolving hypergraphs *Proceedings of the 2017 ACM on Conference on Information and Knowledge Management*,929-938,Singapore.
- [17] Khuller, S. and Saha, B. 2009. On finding dense subgraphs, *ICALP 2009*: 597-608.
- [18] Kim, S. 2011. Higher-order correlation clustering for image segmentation. *Advances in Neural Information Processing Systems* 1530–8.
- [19] Liang, J., Ke C. and Honorio, J.(2021). Information theoretic limits of exact recovery in subhypergraph models for community detection. <https://arxiv.org/pdf/2101.12369.pdf>
- [20] Luo, Y. and Zhang, A. (2020). Open problem: average-case hardness of hypergraphic planted clique detection. *Proceedings of Machine Learning Research*, 1-4, 2020.
- [21] Liu, H., Jan, L. and Yan, S.(2015). Dense subgraph partition of positive hypergraphs. *IEEE Transactions on Pattern Analysis and Machine Intelligence*,37,3:541-554.
- [22] Tsourakakis, C.(2015). The K-clique densest subgraph problem, *Proceedings of the 24th International Conference on World Wide Web*,1122-1132,Florence, Italy.
- [23] Wu, M., Li, X., Kwok, C.K. and Ng. S.K. 2009. A coreattachment based method to detect protein complexes in ppi networks, *BMC bioinformatics*, 10: 169.
- [24] Yuan, M. and Shang, Z. (2021). Sharp detection boundaries on testing dense subhypergraph. <https://arxiv.org/abs/2101.04584>
- [25] Yuan, M. and Shang, Z. (2021). Heterogeneous Dense Subhypergraph Detection. <https://arxiv.org/pdf/2104.04047.pdf>
- [26] Zhou, D., Huang,J. and Scholkopf, B. (2006). Learning with hyper graphs: Clustering, classification, and embedding. *Advances in Neural Information Processing Systems*. 6:1601–8.