DISENTANGLED REPRESENTATION LEARNING WITH THE GROMOV-MONGE GAP

Theo Uscidda 1,2* Luca Eyring 2,3,4,* Karsten Roth 2,4,5 Fabian Theis 2,3,4 Zeynep Akata 2,3,4,† Marco Cuturi 1,6,†

 1 CREST-ENSAE 2 Helmholtz Munich 3 TU Munich 4 MCML 5 Tubingen AI Center 6 Apple theo.uscidda@ensae.fr luca.eyring@tum.de

ABSTRACT

Learning disentangled representations from unlabelled data is a fundamental challenge in machine learning. Solving it may unlock other problems, such as generalization, interpretability, or fairness. Although remarkably challenging to solve in theory, disentanglement is often achieved in practice through prior matching. Furthermore, recent works have shown that prior matching approaches can be enhanced by leveraging geometrical considerations, e.g., by learning representations that preserve geometric features of the data, such as distances or angles between points. However, matching the prior while preserving geometric features is challenging, as a mapping that *fully* preserves these features while aligning the data distribution with the prior does not exist in general. To address these challenges, we introduce a novel approach to disentangled representation learning based on quadratic optimal transport. We formulate the problem using Gromov-Monge maps that transport one distribution onto another with minimal distortion of predefined geometric features, preserving them as much as can be achieved. To compute such maps, we propose the Gromov-Monge-Gap (GMG), a regularizer quantifying whether a map moves a reference distribution with minimal geometry distortion. We demonstrate the effectiveness of our approach for disentanglement across four standard benchmarks, outperforming other methods leveraging geometric considerations.

1 Introduction

Learning low-dimensional representations of high-dimensional data is a fundamental challenge in unsupervised deep learning (Bengio et al., 2014). Emphasis is put on learning representations that allow for efficient and robust adaptation across a wide range of tasks (Higgins et al., 2018; Locatello et al., 2019a). The fundamental property of *disentanglement* has shown significant promise to improve generalization (Locatello et al., 2020; Roth et al., 2023; Hsu et al., 2023; Barin-Pacela et al., 2024), interpretability and fairness (Locatello et al., 2019b; Träuble et al., 2021). Most works regard disentanglement as a one-to-one map between learned representations and ground-truth latent factors, effectively seeking to recover these factors from data alone in an unsupervised fashion. While unsupervised disentanglement is theoretically impossible (Locatello et al., 2019a), the inductive biases of autoencoder architectures ensure effective disentanglement in practice (Rolinek et al., 2019; Zietlow et al., 2021). Most approaches operate using variational autoencoder (VAE) frameworks (Kingma and Welling, 2014), using objectives that match latent VAE posteriors to factorized priors (Higgins et al., 2017; Kim and Mnih, 2018; Kumar et al., 2018; Burgess et al., 2018; Chen et al., 2018).

More recently, studies such as Gropp et al. (2020); Chen et al. (2020a); Lee et al. (2022); Horan et al. (2021); Nakagawa et al. (2023); Huh et al. (2023); Hahm et al. (2024) have provided a new perspective, showing that geometric constraints on representation spaces may also enable disentanglement. Typically, latent representations are encouraged to preserve key geometric features of the data distribution, such as (scaled) distances or angles between samples. Horan et al. (2021) even demonstrate that unsupervised disentanglement is *always* possible provided that the latent space

^{*}Equal contribution

[†]Equal advising†

is locally isometric to the data, further supporting the geometric desiderata. However, combining prior matching with these geometric aspects is challenging. In general, a mapping that perfectly aligns the data distribution with the prior while *fully* preserving the geometric features of interest may not exist. This leads to an *inherent trade-off*: Practitioners must carefully fine-tune regularization terms, either by altering prior matching to prioritize geometry preservation, or vice-versa.

In this work, we demonstrate how to effectively combine geometric desiderata with prior matching within the VAE framework, using optimal transport (OT) theory (Santambrogio, 2015; Peyré and Cuturi, 2019). By treating mappings from the data manifold to the latent space (encoders) or vice versa (decoders) as transport maps $T: \mathcal{X} \to \mathcal{Y}$, we can leverage the Gromov-{Monge, Wasserstein} paradigm (Sturm, 2020; Mémoli, 2011), which aligns two distributions by finding a mapping that minimizes the distortion between intra-domain cost functions defined on their supports. Specifically, we consider cost functions $c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}')$ on \mathcal{X} and $c_{\mathcal{Y}}(\mathbf{y}, \mathbf{y}')$ on \mathcal{Y} that encode geometric features such as scaled distances or angles. Consequently, the resulting mapping transforms one distribution onto the other while preserving these geometric features as much as possible.

Our Contribution: A novel OT-based approach to disentanglement through geometric considerations.

- (i) We address the challenge of learning disentangled representations using geometric constraints by leveraging Gromov-Monge mappings between the data and prior distributions. Since *fully* preserving geometric features—such as (scaled) distances or angles between points—during the alignment of these two distributions is generally impossible, we aim to find an alignment that, instead, minimizes the distortion of these features, thereby preserving them *as much as possible*.
- (ii) Inspired by (Uscidda and Cuturi, 2023), we introduce the *Gromov-Monge Gap* (GMG), a regularizer that measures how closely a map T approximates a Gromov-Monge map for costs $c_{\mathcal{X}}, c_{\mathcal{Y}}$. GMG measures whether T transports distributions with minimal distortion w.r.t. $c_{\mathcal{X}}, c_{\mathcal{Y}}$. We propose an efficient procedure to compute GMG and describe how to integrate it within the VAE framework.
- (iii) We show that when $c_{\mathcal{X}}$ and $c_{\mathcal{Y}}$ encode scaled distances or angles, the GMG and its finite-sample counterpart are weakly convex functions. In both cases, we precisely characterize the weak convexity constants and analyze their practical implications for practitioners.
- (iv) Across four standard disentangled representation learning benchmarks, we show that incorporating geometry-preserving desiderata via the GMG significantly enhances disentanglement across various methods—from the standard β -VAE to the combination of β -TCVAE with HFS (Roth et al., 2023)—notably outperforming other methods that leverage geometric constraints.

2 BACKGROUND AND RELATED WORKS

2.1 DISENTANGLED REPRESENTATION LEARNING

The Disentanglement Formalism. Disentanglement has varying operational definitions. In this work, we follow the common understanding (Locatello et al., 2019a; 2020; Roth et al., 2023; Träuble et al., 2021; Higgins et al., 2017) where data \mathbf{x} is generated by a process $p(\mathbf{x}|\mathbf{z})$ operating on ground-truth latent factors $\mathbf{z} \sim p(\mathbf{z})$, modeling underlying source of variations (s.a. object shape, color, background...). Given a dataset $\mathcal{D} = \{\mathbf{x}_i\}_{i=1}^N, \mathbf{x}_i \sim p_{\text{data}},$ unsupervised disentangled representation learning aims to find a mapping e_{ϕ} s.t. $e_{\phi}(\mathbf{x}_i) \approx \mathbb{E}[\mathbf{z}|\mathbf{x}_i]$, up to element-wise transformations. Notably, this is to be achieved without prior information on $p(\mathbf{z})$ and $p(\mathbf{x}|\mathbf{z})$.

Unsupervised Disentanglement through Prior Matching. Most unsupervised disentanglement methods operate on variational autoencoders (VAEs, Kingma and Welling (2014)), which define a generative model of the form $p_{\theta}(\mathbf{x}, \mathbf{z}) = p(\mathbf{z})p_{\theta}(\mathbf{x}|\mathbf{z})$. Here, $p_{\theta}(\mathbf{x}|\mathbf{z})$ is a product of exponential family distributions with parameters computed by a decoder $d_{\theta}(\mathbf{z})$. The latent prior $p(\mathbf{z})$ is usually chosen as a standard Gaussian $\mathcal{N}(\mathbf{z}|\mathbf{0}_d,\mathbf{I}_d)$, and the probabilistic encoder $q_{\phi}(\mathbf{z}|\mathbf{x})$ is implemented through neural networks $e_{\phi}(\mathbf{x})$, $\sigma_{\phi}(\mathbf{x})$ that predicts the latent parameters so that $q_{\phi}(\mathbf{z}|\mathbf{x}) = \mathcal{N}(\mathbf{z}|e_{\phi}(\mathbf{x}), \sigma_{\phi}^2(\mathbf{x}))$. The β -VAE (Higgins et al., 2017) achieves disentanglement by minimizing

$$\min_{\theta,\phi} \mathbb{E}_{\mathbf{x} \sim p_{\text{data}}, \mathbf{z} \sim q_{\phi}(\mathbf{z}|\mathbf{x})} \underbrace{\left[-\log p_{\theta}(\mathbf{x}|\mathbf{z}) + \underbrace{\beta D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x})||p(\mathbf{z}))}_{\text{(ii) prior matching}} \right]}, \tag{1}$$

which enforces β -weighted prior matching on top of the reconstruction loss, assuming statistical factor independence (Roth et al., 2023). Several follow-ups refine latent prior matching through

different losses or prior choices (Rolinek et al., 2019; Kim and Mnih, 2018; Kumar et al., 2018; Burgess et al., 2018; Chen et al., 2018; Moor et al., 2021; Balabin et al., 2024).

Disentanglement through a Geometric Lens. Recent research suggests that disentanglement can arise by imposing geometric constraints on representation spaces (Horan et al., 2021). Most works (Huh et al., 2023; Karras et al., 2020) promote latent representations to preserve scaled distances between data points. Closest to our work, Lee et al. (2022) use a Jacobian-based regularization within a VAE, while Nakagawa et al. (2023) employ a distortion-based regularization, directly comparing pairwise squared Euclidean distances, within a Wasserstein AE (WAE, Tolstikhin et al. (2019)).

2.2 QUADRATIC OPTIMAL TRANSPORT

Gromov-{Monge, Wasserstein} Formulations. OT (Peyré and Cuturi, 2019) involves transferring a probability distribution to another with inductive biases. When these distributions lie on incomparable domains, one relies on the Gromov-Monge (GM) and GW problems, also known as quadratic formulations of OT. Consider two compact $\mathcal{X} \subset \mathbb{R}^{d_{\mathcal{X}}}$, $\mathcal{Y} \subset \mathbb{R}^{d_{\mathcal{Y}}}$, each of them equipped with an *intra-domain* cost $c_{\mathcal{X}}: \mathcal{X} \times \mathcal{X} \to \mathbb{R}$ and $c_{\mathcal{Y}}: \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$. For technical reasons, we assume that $c_{\mathcal{X}}$ and $c_{\mathcal{Y}}$ (or $-c_{\mathcal{X}}$ and $-c_{\mathcal{Y}}$) are CPD kernels (see Def. (A.1)). For $p \in \mathcal{P}(\mathcal{X})$ and $q \in \mathcal{P}(\mathcal{Y})$ —two distributions supported on each domain—the GM problem (Mémoli and Needham, 2022) seeks a map $T: \mathcal{X} \to \mathcal{Y}$ that push-forwards p onto q, while minimizing the distortion of the costs:

$$\inf_{T:T \nmid p=q} \int_{\mathcal{X} \times \mathcal{X}} \frac{1}{2} |c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}') - c_{\mathcal{Y}}(T(\mathbf{x}), T(\mathbf{x}'))|^2 \, \mathrm{d}p(\mathbf{x}) \, \mathrm{d}p(\mathbf{x}') \,. \tag{GMP}$$

When it exists, we call a solution T^* to (GMP) a *Gromov-Monge map* for costs $c_{\mathcal{X}}, c_{\mathcal{Y}}$. However, this formulation is ill-suited for discrete distributions p, q, as the constraint set might be empty in that case. Replacing maps by coupling $\pi \in \Pi(p,q)$, i.e. distributions on $\mathcal{X} \times \mathcal{Y}$ with marginals p and q, we obtain the GW problem (Sturm, 2020; Mémoli, 2011)

$$GW(p,q) := \min_{\pi \in \Pi(p,q)} \int_{(\mathcal{X} \times \mathcal{Y})^2} \frac{1}{2} |c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}') - c_{\mathcal{Y}}(\mathbf{y}, \mathbf{y}')|^2 d\pi(\mathbf{x}, \mathbf{y}) d\pi(\mathbf{x}', \mathbf{y}').$$
 (GWP)

A solution π^* to (GWP) always exists, making $\mathrm{GW}(p,q)$ a well-defined quantity. It quantifies the minimal distortion of the geometries induced by $c_{\mathcal{X}}$ and $c_{\mathcal{Y}}$ achievable when coupling p and q.

Discrete Solvers. When both p and q are instantiated as samples, GW Prob. (GWP) translates to a quadratic assignment problem, whose objective can be regularized using entropy (Cuturi, 2013; Peyré et al., 2016). For empirical measures $p_n = \frac{1}{n} \sum_{i=1}^n \delta_{\mathbf{x}_i}$, $q_n = \frac{1}{n} \sum_{j=1}^n \delta_{\mathbf{y}_j}$ and $\varepsilon \ge 0$, we set:

$$GW_{\varepsilon}(p_n, q_n) := \min_{\mathbf{P} \in U_n} \sum_{i, j, i', j'=1}^{n} (\mathbf{C}_{\mathcal{X}_{i, i'}} - \mathbf{C}_{\mathcal{Y}_{j, j'}})^2 \mathbf{P}_{i, j} \mathbf{P}_{i', j'} - \varepsilon H(\mathbf{P}), \qquad (EGWP)$$

with $\mathbf{C}_{\mathcal{X}} = [c_{\mathcal{X}}(\mathbf{x}_i, \mathbf{x}_{i'})]_{i,i'}, \mathbf{C}_{\mathcal{Y}} = [c_{\mathcal{Y}}(\mathbf{y}_j, \mathbf{y}_{j'})]_{j,j'} \in \mathbb{R}^{n \times n}, U_n = \{\mathbf{P} \in \mathbb{R}_+^{n \times n}, \mathbf{P}\mathbf{1}_n = \mathbf{P}^T\mathbf{1}_n = \frac{1}{n}\mathbf{1}_n\}$ and $H(\mathbf{P}) = -\sum_{i,j=1}^n \mathbf{P}_{i,j} \log(\mathbf{P}_{i,j})$. As $\varepsilon \to 0$, we recover $\mathrm{GW}_0^{c_{\mathcal{X}},c_{\mathcal{Y}}} = \mathrm{GW}^{c_{\mathcal{X}},c_{\mathcal{Y}}}$. In addition to yielding greater convexity (Rioux et al., 2023) and better statistical (Zhang et al., 2023) and properties, entropic regularization improves computational performance. We can solve (EGWP) using a scheme that iterates the Sinkhorn algorithm (Peyré et al., 2016). This solver has $\mathcal{O}(n^2)$ memory complexity. Its time complexity depends on $c_{\mathcal{X}}, c_{\mathcal{Y}}$. For general $c_{\mathcal{X}}, c_{\mathcal{Y}}$, it runs in $\mathcal{O}(n^3)$. However, for the most common practical choices of $c_{\mathcal{X}} = c_{\mathcal{Y}} = \langle \cdot, \cdot \rangle$ or $c_{\mathcal{X}} = c_{\mathcal{Y}} = \|\cdot - \cdot\|_2^2$, it can also be reduced to $\mathcal{O}(n^2(d_{\mathcal{X}} + d_{\mathcal{Y}}))$, as detailed by Scetbon et al. (2022, §3 & Alg. 2).

Neural GW. While for classical OT, numerous neural methods have been proposed (Makkuva et al., 2020; Korotin et al., 2022; Tong et al., 2023; Eyring et al., 2024), the GW setting has received less attention. Carrasco et al. (2024) propose an approach for solving GM with inner product costs, that involves a min-max-min optimization procedure. Klein et al. (2024) propose a framework to computeentropic GW couplings. Yet, these formulations are end-to-end, and not based on a regularizer that can be flexibly added to a loss function. Therefore, they cannot be combined with other models, as we do in this work with the β -VAE. In a concurrent work, Sotiropoulou and Alvarez-Melis (2024) introduced the Gromov-Monge gap. However, (i) they do not apply it for disentangled representational learning, and (ii) they do not investigate its properties, as we do in this work.

3 DISENTANGLEMENT WITH THE GROMOV-MONGE GAP

This section outlines our novel OT-based approach to disentanglement through geometric considerations within the VAE framework. First, we explore how an arbitrary map $T:\mathcal{X}\to\mathcal{Y}$ between two domains can be encouraged to preserve predefined geometric features. In a VAE, T may represent either the encoder e_ϕ , which generates latent codes from the data, or the decoder d_θ , which reconstructs the data from these codes. In the case of the encoder, the source domain \mathcal{X} corresponds to the data, and the target domain \mathcal{Y} is the latent space, while for the decoder, these roles are swapped. Assuming that d_θ perfectly reconstructs the data from the latents produced by e_ϕ , i.e., $e_\phi \circ d_\theta = \mathrm{I} d$, the preservation of geometric features by either e_ϕ or d_θ becomes equivalent. Therefore, in the following sections, we refer to T as either the encoder or the decoder without loss of generality.

Outline. Building on this perspective, this section begins by defining cost functions to encode geometric features and the notion of distortions in §3.1. We leverage this concept in §3.2 to introduce the Gromov-Monge Gap (GMG), a regularizer that measures whether a map moves distributions while preserving geometric features as much as possible, i.e., minimizing distortion. §3.3 then shows how the GMG can be efficiently computed from samples to be practically applicable in the VAE framework. This transitions into §3.4 studying (weak) convexity properties of the GMG, as an operator. Finally, in §3.5, we describe how to integrate the GMG with disentangled representation learning objectives, effectively combining prior matching with geometric constraints.

3.1 From the distortion...

We encode the geometric features of interest using two cost functions, one for each domain: $c_{\mathcal{X}}: \mathcal{X} \times \mathcal{X} \to \mathbb{R}$ and $c_{\mathcal{Y}}: \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$. We aim for T to preserve the geometric feature, which means that T preserves these costs, that is, $c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}') \approx c_{\mathcal{Y}}(T(\mathbf{x}), T(\mathbf{x}'))$ for $\mathbf{x}, \mathbf{x}' \in \mathcal{X}$. In practice, two types of cost function from this family are particularly relevant:

- [i] (Scaled) squared L2 distance: $c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}') = \|\mathbf{x} \mathbf{x}'\|_2^2$ and $c_{\mathcal{Y}}(\mathbf{y}, \mathbf{y}') = \alpha^2 \|\mathbf{y} \mathbf{y}\|_2^2$, with $\alpha > 0$. A map T preserving $c_{\mathcal{X}}, c_{\mathcal{Y}}$ preserves the scaled distances between the points, i.e. it is a scaled isometry. When $\alpha = 1$, we recover the standard definition of an isometry.
- [ii] Cosine-Similarity: $c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}') = \operatorname{cos-sim}(\mathbf{x}, \mathbf{x}') := \langle \frac{\mathbf{x}}{\|\mathbf{x}\|_2}, \frac{\mathbf{x}'}{\|\mathbf{x}'\|_2} \rangle$ and $c_{\mathcal{Y}}(\mathbf{y}, \mathbf{y}') = \operatorname{cos-sim}(\mathbf{y}, \mathbf{y}')$ similarly. On has $\operatorname{cos-sim}(\mathbf{x}, \mathbf{x}') = \operatorname{cos}(\theta_{\mathbf{x}, \mathbf{x}'})$ where $\theta_{\mathbf{x}, \mathbf{x}'}$ is the angle between \mathbf{x} and \mathbf{x}' . A map T preserving $c_{\mathcal{X}}, c_{\mathcal{Y}}$ then preserves the angles between the points, i.e. it is a conformal map. Note that if T is (scaled) isometry (see above), it is a conformal map.

We refer to these costs via $L2^2$ for [i] with $\alpha = 1$, $ScL2^2$ for [i] with $\alpha \neq 1$ and Cos for [ii]. Introducing a reference distribution $r \in \mathcal{P}(\mathcal{X})$, weighting the areas of \mathcal{X} where we penalize deviations of $c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}')$ from $c_{\mathcal{Y}}(T(\mathbf{x}), T(\mathbf{x}'))$, we can quantify this property using the following criterion:

Definition 3.1 (Distortion). The distortion (DST) of a map T, for cost functions $c_{\mathcal{X}}$, $c_{\mathcal{Y}}$ and reference distribution r, is defined as:

$$DST_r(T) := \int_{\mathcal{X} \times \mathcal{X}} \frac{1}{2} (c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}') - c_{\mathcal{Y}}(T(\mathbf{x}), T(\mathbf{x}')))^2 dr(\mathbf{x}) dr(\mathbf{x}').$$
 (DST)

 $\mathrm{DST}_r(T)$ quantifies how much T distorts geometric features encoded by $c_{\mathcal{X}}, c_{\mathcal{Y}}$ on the support of r, that is, when $\mathrm{DST}_r(T) = 0$, one has $c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}') = c_{\mathcal{Y}}(T(\mathbf{x}), T(\mathbf{x}'))$ for $\mathbf{x}, \mathbf{x}' \in \mathrm{Spt}(r)$. Nakagawa et al. (2023) suggest promoting geometry preservation by regularizing the encoder d_{θ} using the DST , with $\mathrm{ScL2}^2$ as costs, and the latent representation as reference distribution, namely $r = e_{\phi} \sharp p_{\mathrm{data}}$. While they use it within a WAE, translated in our VAE setting, this corresponds to minimizing

$$\min_{\theta,\phi} \mathbb{E}_{\mathbf{x} \sim p_{\text{data}}, \mathbf{z} \sim q_{\phi}(\mathbf{z}|\mathbf{x})} \underbrace{\left[-\log p_{\theta}(\mathbf{x}|\mathbf{z}) + \underbrace{\beta D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x})||p(\mathbf{z}))}_{\text{(ii) prior matching}} \right] + \underbrace{\lambda DST_r(d_{\theta})}_{\text{(iii) geom. preservation}}, \tag{2}$$

where $\lambda > 0$. Given the choice of the costs, d_{θ} is distortion-free (i.e., $\mathrm{DST}_r(d_{\theta}) = 0$), if it is a scaled isometry. Since a scaled isometry that maps the prior onto the data distribution may not exist, there is an inherent trade-off between minimizing terms (ii) and (iii), which are responsible for achieving practical disentanglement. As these terms cannot be simultaneously minimized to 0, the DST loss will move away from accurately matching the prior, which will negatively impact




Figure 1: Sketch of $\mathrm{GMG}_{r_n}(T)$ for two different maps T. We use a discrete reference distribution r_n on 3 points, and $c_{\mathcal{X}} = c_{\mathcal{Y}} = \|\cdot - \cdot\|_2$, hence measuring if T minimally distorts the distances. On the left, T is the optimal map, T^\star , and maps the three points with minimal (yet non-zero) distortion, which is measured as the sum of the squared lengths of the orange segments. This results in $\mathrm{GMG}_{r_n}(T) = 0$. On the right, T swaps two points compared to T^\star —specifically, $\mathbf{x_2}$ and $\mathbf{x_3}$ —causing a higher distortion than the minimal one, and measured as the sum of the squared lengths of the red segments. This results in $\mathrm{GMG}_{r_n}(T) > 0$, equal to the gap between these distortions.

the quality of the learned latent representations. This naturally raises the question of how to avoid this over-penalization. Instead of seeking a distortion-free decoder, we should seek a decoder that transports the prior to the data distribution with *minimal distortion* of the costs $c_{\mathcal{X}}, c_{\mathcal{Y}}$. In other words, the decoder should be a Gromov-Monge map between the prior and the data distributions for costs $c_{\mathcal{X}}, c_{\mathcal{Y}}$ (see 2.2). Conversely, if we choose to regularize the encoder, the same reasoning applies by swapping the roles of the prior and the data distribution. In the next section, we introduce a regularizer to fit Gromov-Monge maps, which we will use as a replacement to the DST.

3.2 ... TO THE GROMOV-MONGE GAP

Recently, Uscidda and Cuturi (2023) introduced the Monge gap, a regularizer that measures whether a map T transports a reference distribution at the minimal displacement cost. Building on this concept, we replace "displacement" with "distortion" to introduce the Gromov-Monge gap, a regularizer that assesses whether a map T transports a reference distribution at the minimal distortion cost.

Definition 3.2 (Gromov-Monge gap). The Gromov-Monge gap (GMG) of a map T, for cost functions $c_{\mathcal{X}}, c_{\mathcal{Y}}$ and reference distribution r, is defined as:

$$GMG_r(T) := DST_r(T) - GW(r, T\sharp r)$$
(GMG)

We recall from Eq. (GWP) that $\mathrm{GW}(r,T\sharp r)$ represents the minimal distortion of $c_{\mathcal{X}},c_{\mathcal{Y}}$ achievable when transporting r to $T\sharp r$. Thus, the GMG quantifies the difference between the distortion incurred when transporting r to $T\sharp r$ via T, and this minimal distortion. Formally, when Prob. (GMP) and Prob. (GWP) between r and $T\sharp r$ are equivalent, the the GMG is the suboptimality gap of T in Prob. (GMP). This is the case, for example, when r is a density and $c_{\mathcal{X}}=c_{\mathcal{Y}}=\langle\cdot,\cdot\rangle$ (Dumont et al., 2022). Otherwise, the GMG is the suboptimality gap of $\pi=(\mathrm{I}_d,T)\sharp r$ in Prob. (GWP) between r and $T\sharp r$. In light of this, it is a well-defined quantity and:

- The GMG measures how close T is to be a Gromov-Monge map for costs $c_{\mathcal{X}}, c_{\mathcal{Y}}$. Indeed, $\mathrm{GMG}_r(T) \geq 0$ with equality if T is a Gromov-Monge map solution of Prob. (GMP) between r and $T\sharp r$, i.e., T moves r with minimal (but eventually non zero) distortion; see App. B.1.
- When transport without distortion is possible, the GMG coincides with the distortion. When there exists another map $U: \mathcal{X} \to \mathcal{Y}$ transporting r to $T\sharp r$ with zero distortion, i.e., $U\sharp r = T\sharp r$ and $\mathrm{DST}_r(U) = 0$, then $\mathrm{GMG}_r(T) = \mathrm{DST}_r(T)$. Indeed, $\mathrm{GW}(r,T\sharp r) = 0$ in that case, as the coupling $\pi = (\mathrm{Id},U)\sharp r$ sets the GW objective to zero, thereby minimizing it.

The last point illustrates how the GMG functions as a *debiased distortion*. It compares the distortion induced by T to a baseline distortion, defined as the minimal achievable distortion when transforming

r into $T \sharp r$. Thus, when transformation without distortion is achievable, this baseline becomes zero, and the GMG equals the distortion. Consequently, the GMG offers the optimal compromise: it avoids the over-penalization induced by the distortion when fully preserving $c_{\mathcal{X}}, c_{\mathcal{V}}$ is not feasible, yet it coincides with it when such full preservation is feasible. See Fig. 1 for a simple illustration.

The Influence of the Reference Distribution. A crucial property of DST_r is that if T transforms r without distortion, it will also apply distortion-free to any distribution s whose support is contained within that of r. Formally, if $DST_r(T) = 0$ and $s \in \mathcal{P}(\mathcal{X})$ with $Spt(s) \subseteq Spt(r)$, then $DST_s(T) = 0$ 0. This raises a question for the GMG: If T maps r with minimal distortion, does it similarly map s with minimal distortion? We answer this question positively with Prop. (3.3).

Proposition 3.3. If $GMG_r(T) = 0$, $\forall s \in \mathcal{P}(\mathcal{X})$ s.t. $Spt(s) \subseteq Spt(r)$, one has $GMG_s(T) = 0$.

3.3 ESTIMATION AND COMPUTATION FROM SAMPLES

Plug-In Estimation. In practice, we estimate Eq. (DST) and Eq. (GMG) from samples $\mathbf{x}_1,...,\mathbf{x}_n \sim r$. We consider the empirical version $r_n := \frac{1}{n} \sum_{i=1}^n \delta_{\mathbf{x}_i}$ of r and use plug-in estimators, i.e.

$$DST_{r_n}(T) = \frac{1}{n^2} \sum_{i,j=1}^n (c_{\mathcal{X}}(\mathbf{x}_i, \mathbf{x}_j) - c_{\mathcal{Y}}(T(\mathbf{x}_i), T(\mathbf{x}_j)))^2,$$
(3)

and $\mathrm{GMG}_{r_n}(T) = \mathrm{DST}_{r_n}(T) - \mathrm{GW}(r_n, T\sharp r_n)$, where $T\sharp r_n = \frac{1}{n}\sum_{i=1}^n \delta_{T(\mathbf{x}_i)}$.

Efficient and Stable Computation. Computing the GMG requires solving a discrete GW problem between r_n and $T\sharp r_n$ to get $\mathrm{GW}(r_n,T\sharp r_n)$. We compute this term using an entropic regularization $\varepsilon \geq 0$, as in Eq. (EGWP):

$$GMG_{r_n,\varepsilon}(T) := DST_{r_n}(T)$$
 (4)
- $GW_{\varepsilon}(r_n, T \sharp r_n)$.

Choosing $\varepsilon = 0$, we recover $GMG_{r_n,0}(T) = GMG_{r_n}(T)$. Moreover, the entropic estimator preserves

Algorithm 1 GMG($\mathbf{x}_1, \dots \mathbf{x}_n, T, \varepsilon$).

- 1: **Require:** $\mathbf{x}_1, \dots, \mathbf{x}_n \sim r$; map T; entropic solver GW, entropic regularization scale ε_0 (default = 0.1), statistic operator on cost matrix stat (default = mean).
- 2: $\mathbf{t}_1, \dots \mathbf{t}_n \leftarrow T(\mathbf{x}_1), \dots, T(\mathbf{x}_n)$.
- 3: $\mathbf{C}_{\mathcal{X}} \leftarrow [c_{\mathcal{X}}(\mathbf{x}_{i}, \mathbf{x}_{i'})]_{1 \leq i, i' \leq n}$ \triangleright usually $\mathcal{O}(n^{2}d_{\mathcal{X}})$ 4: $\mathbf{C}_{\mathcal{Y}} \leftarrow [c_{\mathcal{Y}}(\mathbf{t}_{j}, \mathbf{t}_{j'})]_{1 \leq j, j' \leq n}$ \triangleright usually $\mathcal{O}(n^{2}d_{\mathcal{Y}})$ 5: DST \leftarrow mean $((\mathbf{C}_{\mathcal{X}} \mathbf{C}_{\mathcal{Y}})^{2})$

- 6: $\varepsilon \leftarrow \varepsilon_0 \cdot \operatorname{stat}(\mathbf{C}_{\mathcal{X}}) \cdot \operatorname{stat}(\mathbf{C}_{\mathcal{Y}}) \quad \triangleright \text{ usually } \mathcal{O}(n^2)$
- 7: GW \leftarrow GW($\mathbf{C}_{\mathcal{X}}, \mathbf{C}_{\mathcal{Y}}, \varepsilon$) $\triangleright \mathcal{O}(n^3)$ or $\mathcal{O}(n^2(d_{\mathcal{X}} + d_{\mathcal{Y}}))$
- 8: return DST GW

positivity, as for $\varepsilon \geq 0$, $\mathrm{GMG}_{r_n,\varepsilon}(T) \geq 0$ (see B.3). We compute $\mathrm{GW}_{\varepsilon}(r_n,T\sharp r_n)$ using Peyré et al. (2016)'s solver introduced in 2.2. We use the implementation provided by ott-jax (Cuturi et al., 2022). In practice, we select ε based on (positive) statistics from the cost matrices $C_{\mathcal{X}}, C_{\mathcal{Y}}$. We define a scale ε_0 and set $\varepsilon = \varepsilon_0 \cdot \text{stat}(\mathbf{C}_{\mathcal{X}}) \cdot \text{stat}(\mathbf{C}_{\mathcal{Y}})$. Standard options for the statistic include $\mathtt{stat} \in \{\mathtt{mean}, \mathtt{max}, \mathtt{std}\}.$ This procedure is equivalent to running the entropic GW solver on the re-scaled cost matrices $C_{\mathcal{X}}/\text{stat}(C_{\mathcal{X}})$ and $C_{\mathcal{Y}}/\text{stat}(C_{\mathcal{Y}})$ with $\varepsilon = \varepsilon_0$; see App. B.2. It is a common practical trick, initially suggested for stabilizing the Sinkhorn algorithm Cuturi (2013), then extended to the entropic GW solver (Cuturi et al., 2022). Additionally, as suggested by Scetbon et al. (2022), to promote the solver's stability, we always initialize it with $\mathbf{P}_{\text{init}} = \frac{1}{n^2} \mathbf{1}_n \mathbf{1}_n^{\top}$.

Computational Complexity. For usual costs, including inner products, ℓ_p^q distances, and standard CPD kernels, the computation of $c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}')$ (resp. $c_{\mathcal{Y}}(\mathbf{y}, \mathbf{y}')$) can be done in $\mathcal{O}(d_{\mathcal{X}})$ time (resp. $\mathcal{O}(d_{\mathcal{V}})$). Consequently, the DST can be computed in $\mathcal{O}(n^2(d_{\mathcal{X}}+d_{\mathcal{V}}))$ time. Furthermore, as discussed in § 2.2, the time complexity of the entropic GW solver is $\mathcal{O}(n^3)$ in general, but can be reduced to $\mathcal{O}(n^2(d_{\mathcal{X}}+d_{\mathcal{Y}}))$ when $c_{\mathcal{X}}=c_{\mathcal{Y}}=\langle\cdot,\cdot\rangle$, or $c_{\mathcal{X}}=\|\cdot-\cdot\|_2^2, c_{\mathcal{Y}}=\alpha\|\cdot-\cdot\|_2^2$. Therefore, since the cosine similarity is equivalent to the inner product, up to pre-normalization of \mathbf{x}_i and $T(\mathbf{x}_i)$, this solver runs in $\mathcal{O}(n^2(d_{\mathcal{X}}+d_{\mathcal{Y}}))$ for the costs of interest $(\mathbf{Sc})\mathbf{L2^2}$ and \mathbf{Cos} . The complete algorithm, along with a time complexity analysis of each step, is described in Alg. 1.

(WEAK) CONVEXITY OF THE GROMOV-MONGE GAP

As laid out, the GMG can be used as a regularization loss to push any model T to move distributions with minimal distortion. A natural question arises: is this regularizer convex? In the following, we study the convexity of $T\mapsto \mathrm{GMG}_r(T)$ and its finite-sample counterpart $T\mapsto \mathrm{GMG}_{r_n}(T)$. We focus on the costs $\mathbf{L2}$ and \mathbf{Cos} . For simplicity, we replace \mathbf{Cos} with $\langle\cdot,\cdot\rangle$, as these costs are equivalent, up to normalization of r and T. We respectively denote by GMG_r^2 and $\mathrm{GMG}_r^{\langle\cdot,\cdot\rangle}$ the GMG for these costs. We start by introducing a weaker notion of convexity, previously defined on \mathbb{R}^d (Davis et al., 2018), which we extend here to $L_2(r) := \{T \mid \|T\|_{L_2(r)}^2 := \int_{\mathcal{X}} \|T(\mathbf{x})\|_2^2 \, \mathrm{d}r(\mathbf{x}) < +\infty\}$.

Definition 3.4 (Weak convexity.). With $\gamma > 0$, a functional $\mathcal{F} : L_2(r) \to \mathbb{R}$ is γ -weakly convex if $\mathcal{F}_{\gamma} : T \mapsto \mathcal{F}(T) + \frac{\gamma}{2} \|T\|_{L_2(r)}^2$ is convex.

A weakly convex functional is convex up to an additive quadratic perturbation. The weak convexity constant γ quantifies the magnitude of this perturbation and indicates a degree of non-convexity of \mathcal{F} . A lower γ suggests that \mathcal{F} is closer to being convex, while a higher γ indicates greater non-convexity.

Theorem 3.5. Both GMG_r^2 and $GMG_r^{\langle \cdot, \cdot \rangle}$, as well as their finite sample versions, are weakly convex.

- Finite sample. We note $\mathbf{X} \in \mathbb{R}^{n \times d}$ the matrix that stores the \mathbf{x}_i , i.e. the support of r_n , as rows. Then, (i) $\mathrm{GMG}^2_{r_n}$ and (ii) $\mathrm{GMG}^{\langle \cdot, \cdot \rangle}_{r_n}$ are respectively (i) $\gamma_{2,n}$ and (ii) $\gamma_{inner,n}$ -weakly convex, where: $\gamma_{inner,n} = \lambda_{\max}(\frac{1}{n}\mathbf{X}\mathbf{X}^\top) \lambda_{\min}(\frac{1}{n}\mathbf{X}\mathbf{X}^\top)$ and $\gamma_{2,n} = \gamma_{inner,n} + \max_{i=1...n} \|\mathbf{x}_i\|_2^2$.
- Asymptotic. (i) GMG_r^2 and (ii) $\mathrm{GMG}_r^{\langle \cdot, \cdot \rangle}$ are respectively (i) γ_2 and (ii) γ_{inner} -weakly convex, where: $\gamma_{inner} = \lambda_{\max}(\mathbb{E}_{\mathbf{x} \sim r}[\mathbf{x}\mathbf{x}^\top])$ and $\gamma_{2,n} = \gamma_{inner} + \max_{\mathbf{x} \in \mathrm{Spt}(r)} \|\mathbf{x}\|_2^2$.

From a practitioner's perspective, we analyze the insights provided by Thm. (3.5) in three parts.

- First, we have $\gamma_2 \geq \gamma_{\text{inner}}$. Therefore, GMG_r^2 is less convex than $\mathrm{GMG}_r^{\langle \cdot, \cdot \rangle}$, making it harder to optimize, and the same argument holds for their estimator. In other words, we provably recover that, in practice, preserving the (scaled) distances is harder than simply preserving the angles.
- Second, as $\gamma_{\text{inner}} = \lambda_{\max}(\mathbb{E}_{\mathbf{x} \sim r}[\mathbf{x}\mathbf{x}^{\top}]) \geq \lambda_{\max}(\text{Cov}_{\mathbf{x} \sim r}[\mathbf{x}])$, this exhibits a tradeoff w.r.t. Prop. (3.3): by choosing a bigger reference distribution r, we trade the convexity of the GMG. For γ_2 , the dependency in r is even worse. In practice, we then choose r with support as small as possible, precisely where we want T to move points with minimal distortion.
- Third, and most surprising, the finite sample GMG is more convex in high dimension. Indeed, $\gamma_{\text{inner},n}$ is the spectral width of $\frac{1}{n}\mathbf{X}\mathbf{X}^{\top}$, which contains the (rescaled) inner products between $\mathbf{x}_i \sim r$. When n > d, $\lambda_{\min}(\mathbf{X}\mathbf{X}^{\top}) = 0$ as $\operatorname{rank}(\mathbf{X}\mathbf{X}^{\top}) = d$. Then, $\gamma_{\text{inner},n}$ increases, which in turn decreases the GMG's convexity. However, when d > n, $\lambda_{\min}(\mathbf{X}\mathbf{X}^{\top}) > 0$ if \mathbf{X} is full rank. Intuitively, $\operatorname{GMG}_{r_n}^{\langle \cdot, \cdot \rangle}$ is nearly convex when $\mathbf{X}\mathbf{X}^{\top}$ is well conditioned. Assuming that the \mathbf{x}_i are normalized, this might happen in high dimension, as they will be orthogonal with high probability. This suggests that, contrary to the insights provided by the statistical OT literature (Zhang et al., 2023; Weed and Bach, 2017), the GMG might not benefit a large sample size.

3.5 LEARNING WITH THE GROMOV-MONGE GAP

General Learning Procedure. Given a source p and a target distribution q, we can use the GMG to guide a parameterized map T_{θ} towards approximating a Gromov-Monge map between p and q. We handle the marginal constraint $T_{\theta}\sharp p=q$ separately through a fitting loss $\Delta(T_{\theta},p,q)$. Provided any reference r s.t. $\operatorname{Spt}(p)\subset\operatorname{Spt}(r)$, we minimize

$$\min_{\theta} \Delta(T_{\theta}, p, q) + \lambda GMG_r(T_{\theta})$$
 (5)

 Δ can operate on paired (e.g., in VAE, the reconstruction loss), or unpaired (e.g., in VAE, the KL loss in VAE), samples of p and q. Note that, in theory and as stated in Prop. (3.3), we can select any reference r such that $\operatorname{Spt}(p) \subset \operatorname{Spt}(r)$. However, based on the insights from Thm. (3.4), we typically choose r with minimal support size and, in practice, set r=p. This learning procedure is illustrated in Fig. 2, where we also explore the effect of replacing $\operatorname{GMG}_r(T_\theta)$ by $\operatorname{DST}_r(T_\theta)$ in Eq. (5).

VAE Learning Procedure. In the VAE setting, we can use the GMG promote the (i) encoder e_{ϕ} or the (ii) decoder d_{θ} to mimic a Gromov-Monge map. In (i) we use $r_e = p_{\text{data}}$ the data distribution as reference r, while in (ii) we use the latent distribution $r_d = e_{\phi} \sharp p_{\text{data}}$. Introducing weightings




Figure 2: Learning of geometry-preserving maps with the DST and the GMG. Provided a mixture of Gaussian source distribution p and a uniform or circular target distribution q, we minimize a fitting loss together with a geometry-preserving regularization. Each line correspond to a different data setup and cost. As costs on the top line, we use $\mathbf{L2^2} = \|\cdot - \cdot\|_2^2$, to preserve distances between points while on the bottom line, we use $\mathbf{Cos} = \mathbf{cos\text{-}sim}(\cdot, \cdot)$, to preserve angles.

 $\lambda_e, \lambda_d \geq 0$, determining which mapping we regularize, this remains to minimize

$$\min_{\theta,\phi} \mathbb{E}_{\mathbf{x} \sim p_{\text{data}}, \mathbf{z} \sim q_{\phi}(\mathbf{z}|\mathbf{x})} \underbrace{\left[-\log p_{\theta}(\mathbf{x}|\mathbf{z}) + \underbrace{\beta D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x})||p(\mathbf{z}))}_{\text{(ii) prior matching}} \right] + \underbrace{\lambda_e \text{GMG}_{r_e}(e_{\phi}) + \lambda_d \text{GMG}_{r_d}(d_{\theta})}_{\text{(iii) geom. preservation}},$$
(6)

With this loss, prior matching and geometric desiderata can be efficiently combined, as terms (ii) and (iii) can simultaneously be 0. Note that this loss can be easily extended to more advanced prior matching objectives, such as the β -TCVAE loss (Chen et al., 2018), and can be combined with other regularizers, such as the HFS (Roth et al., 2023). We explore this strategy in experiments § 4.2.

4 EXPERIMENTS

4.1 GMG vs. DST: Illustrative Example

In Fig. 2, we start by illustrating the difference between the DST and GMG on synthetic data. We train T_{θ} to map from a 3D source p to a 2D target distribution q (first and second column) by minimizing $\mathcal{L}(\theta) := S_{\varepsilon}(T_{\theta}\sharp p,q) + R(T_{\theta})$, where S_{ε} is the Sinkhorn divergence (Feydy et al., 2019) and R a geometry-preserving regularizer. We compare three settings, [i] no regularization R = 0, [ii] $R = DST_{p}$ as regularizer, and [iii] $R = GMG_{p}$. For all cases, we plot the transported distribution $T_{\theta}\sharp p$ after training. Without regularization (third column), we fit the marginal constraint $T_{\theta}\sharp p = q$ but do not preserve the geometric features. With the DST (fourth column), we preserve geometric features but do not fit the marginal constraint, i.e., $T_{\theta}\sharp p \neq q$. On the other hand, with the GMG (fifth column), we get the best compromise by approximating a Gromov-Monge map: we fit $T_{\theta}\sharp p = q$, while preserving the geometric features as fully as possible.

4.2 LEVERAGING THE GROMOV-MONGE GAP FOR DISENTANGLEMENT

Experimental Setup. Having demonstrated how the DST enables (i) fitting a marginal constraint while (ii) preserving geometric features as much as possible, we now apply it to disentangled representation learning. Our primary goal is to investigate whether the GMG results in enhanced disentanglement compared to the DST by efficiently combining (i) prior matching with (ii) geometric constraints on the representation space. Moreover, we aim to determine which costs $c_{\mathcal{X}}, c_{\mathcal{Y}}$ to choose in the geometric regularization and what part of the pipeline should be regularized, the encoder e_{ϕ} , or the decoder d_{θ} . The experimental setup is described below.

• Baselines. We use the standard β -VAE and β -TCVAE as our starting models, with the option to apply the recent HFS regularization (Roth et al., 2023) to each, resulting in a total of four base configurations. Note that the latter does not leverage geometric constraints; it is only used to enhance prior matching. We then investigate the effect of various geometry-preserving

Table 1: Effect of different regularization on disentanglement (DCI-D on Shapes3D). We highlight the best method per regularization type (L2², ScL2², or Cos), and the best/second best per column.

	β-VAE	β -TCVAE	β -VAE + HFS	β -TCVAE + HFS
Base	65.8 ± 15.6	75.0 ± 3.4	88.1 ± 7.4	90.2 ± 7.5
	L2	$\mathbf{c}_{\mathcal{X}} = c_{\mathcal{Y}} = c_{Y$	$=\ \cdot-\cdot\ _2^2$	
+ Enc-DST + Enc-GMG + Dec-DST + Dec-GMG	59.6 ± 6.9 62.3 ± 8.4 71.5 ± 3.6 72.0 ± 8.5	75.7 ± 3.0 75.4 ± 5.3 75.8 ± 6.6 78.9 ± 5.0	88.7 ± 7.1 88.4 ± 7.7 92.1 ± 9.7 92.5 ± 4.4	90.3 ± 7.9 90.1 ± 4.3 90.9 ± 7.6 91.7 ± 6.0
$\mathbf{ScL2^2}: c_{\mathcal{X}} = \ \cdot - \cdot\ _2^2, c_{\mathcal{Y}} = \alpha\ \cdot - \cdot\ _2^2, \alpha > 0$ learnable				
+ Jac + Enc-DST + Enc-GMG + Dec-DST + Dec-GMG	$\begin{array}{c} \textbf{61.4} \pm 12.8 \\ \textbf{65.8} \pm 11.9 \\ \textbf{65.1} \pm 5.5 \\ \textbf{67.4} \pm 7.1 \\ \textbf{70.0} \pm 5.9 \end{array}$	76.7 ± 4.5 73.0 ± 7.9 76.1 ± 7.7 77.9 ± 4.5 81.0 ± 3.2	$\begin{array}{c} 90.5 \pm 3.8 \\ 92.4 \pm 3.7 \\ 90.8 \pm 9.2 \\ 93.2 \pm 9.7 \\ \underline{93.3} \pm 8.6 \end{array}$	$\begin{array}{c} 91.5 \pm 5.6 \\ 89.2 \pm 3.8 \\ 92.0 \pm 5.3 \\ 94.5 \pm 6.9 \\ \underline{96.1} \pm 3.8 \end{array}$
$\mathbf{Cos}: c_{\mathcal{X}} = c_{\mathcal{Y}} = \mathbf{cos\text{-}sim}(\cdot, \cdot)$				
+ Enc-DST + Enc-GMG + Dec-DST + Dec-GMG	$69.2 \pm 9.1 70.9 \pm 9.5 \underline{76.8} \pm 4.1 82.1 \pm 4.5$	77.2 ± 7.5 79.6 ± 6.6 81.3 ± 4.7 83.7 ± 8.8	87.7 ± 7.7 92.5 ± 5.9 87.5 ± 3.3 95.7 ± 5.8	90.5 ± 5.9 93.5 ± 6.9 91.9 ± 9.4 96.9 ± 4.9

regularizations on disentanglement on top of these four base configurations. We consider GMG, DST and the Jacobian-based (Jac) regularization (Lee et al., 2022) discussed in 2.1. Given the inclusion of the DST, we naturally consider Nakagawa et al. (2023) as a baseline.

- Metrics. We evaluate the learned representations using the DCI-D (Eastwood and Williams, 2018) as it was found that it is the metric most suitable for measuring the disentanglement (Locatello et al., 2020; Dittadi et al., 2021). We report mean and standard deviation over 5 seeds.
- Datasets. We benchmark over four standard datasets: Shapes3D (Kim and Mnih, 2018), DSprites Higgins et al. (2017), SmallNORB LeCun et al. (2004), and Cars3D Reed et al. (2015).
- Hyperparameters. To ensure a fair experimental comparison, we follow recent works (Locatello et al., 2019a; 2020; Roth et al., 2023) by using the same architecture and hyperparameters. We perform a similar small grid search for the weighting terms: β for the KL loss and γ for HFS. Additionally, we include the weighting λ for the geometry-preserving regularizer in the grid search. Note that we search over the same loss weightings λ for DST, GMG, and Jac. For all experiments with the GMG, we compute it with Alg. 1, and systematically use $\varepsilon_0 = 0.1$ and stat = mean. As a result, ε is *not* included in the grid search. See Appendix C for full details.

Which costs $c_{\mathcal{X}}, c_{\mathcal{Y}}$ should we choose? The first question that naturally arises when using a geometry-preserving regularizer is: Which geometric features should be preserved? Previous works (Lee et al., 2022; Nakagawa et al., 2023; Huh et al., 2023) focused on preserving scaled distances between points, with the scale being learnable. We follow and extend this approach by also investigating plain distances and angles. This leads to three choices for $c_{\mathcal{X}}, c_{\mathcal{Y}}$ (L2², ScL2², Cos), as introduced in §3.1, following the hierarchy of geometry-preserving mappings proposed in Lee et al. (2022). We benchmark these on Shapes3D across various settings including DST, GMG, and Jac. Table 1 shows that angle preservation (Cos), previously unconsidered for disentangled representation learning, consistently outperforms (scaled) distance preservation. This result is intuitive, as preserving angles imposes a weaker constraint, allowing for greater latent space expressiveness. In practice, preserving scaled distances seems to overly restrict the expressiveness of the latent space.

Should we regularize the encoder e_{ϕ} , or the decoder d_{θ} ? The next question we aim to answer is whether the decoder or encoder should be regularized. Therefore, we follow the previous setup on Shapes3D and benchmark all geometry-preserving regularizers on e_{ϕ} and d_{θ} as reported in Table 1. We find that regularizing the decoder is beneficial over regularizing the encoder. We hypothesize this is due to the regularization of d_{θ} offering a stronger signal as its gradients impact both ϕ and θ , as in this case, the reference $r = e_{\phi} \sharp p_{\text{data}}$ is the distribution of encoded images. Our findings align

Table 2: Effect of GMG and DST with Cos as costs on disentanglement, as measured by DCI-D,
over three datasets. We highlight the best , and <u>second best</u> result for each dataset and method.

With Cos costs	β -VAE	β -TCVAE	β -VAE + HFS	β -TCVAE + HFS
DSprites Higgins et al. (2017)				
Base	26.2 ± 18.5	32.3 ± 19.3	33.6 ± 17.9	48.7 ± 10.2
+ Dec-DST	28.6 ± 19.3	32.4 ± 8.5	39.3 ± 18.1	49.0 ± 11.2
+ Dec-GMG	39.5 ± 15.2	42.2 ± 3.6	46.7 ± 2.0	50.1 ± 8.5
SmallNORB LeCun et al. (2004)				
Base	26.8 ± 0.2	29.8 ± 0.4	26.8 ± 0.2	29.8 ± 0.4
+ Dec-DST	28.2 ± 0.3	29.9 ± 0.4	28.2 ± 0.3	29.9 ± 0.4
+ Dec-GMG	28.3 ± 0.6	29.9 ± 0.5	28.3 ± 0.6	29.9 ± 0.5
Cars3D Reed et al. (2015)				
Base	29.6 ± 5.7	32.3 ± 4.6	29.6 ± 5.7	32.3 ± 4.6
+ Dec-DST	$\overline{26.8} \pm 3.6$	33.7 ± 4.2	$\overline{26.8} \pm 3.6$	33.7 ± 4.2
+ Dec-GMG	30.1 ± 5.6	36.4 ± 5.7	30.1 ± 5.6	36.4 ± 5.7

with prior works (Lee et al., 2022; Nakagawa et al., 2023), which focus on regularizing d_{θ} yet do not offer this type of analysis. Additionally, we find that the GMG consistently outperforms the DST over all costs. Overall, the GMG on d_{θ} with Cos achieves best DCI-D results over all baselines. Consequently, moving forward we regularize the decoder for angle preservation.

GMG Consistently Enhances Disentanglement. To further validate our findings, we benchmark the GMG for decoder regularization with angle preservation (Cos) against its distortion counterpart across three more datasets. We report full results in Table 2. Again we observe that the GMG outperforms or performs equally well to its distortion equivalent, confirming the benefits of accounting for the optimal possible mapping in the regularization. Note that for SmallNORB and Cars3D, we found no benefits with respect to DCI-D in adding an HFS regularization and obtained the best results without it. We emphasize that using the GMG with Cos regularization significantly improves results for all datasets versus not using any isometric regularization. This establishes the GMG as an effective tool for enhanced disentanglement.

Stability of the GMG. Furthermore, we investigate the stability of the GMG compared to the DST. Computing the GMG using Algorithm 1 involves solving an optimization problem through a GW solver. This experiment aims to demonstrate that our proposed method for solving the GW problem leads to a stable loss in the GMG. We assess the stability by measuring the alignment of GMG gradients $\nabla_{\theta} \mathrm{GMG}_{r_n}(T_{\theta})$ across 5 randomly sampled batches from each dataset $r_n \sim \mathcal{D}$, for a fixed neural map T_{θ} . We repeat this procedure for each of the four datasets \mathcal{D} considered and apply the same methodology to the DST. Figure 4 presents results for our best-performing configuration, which regularizes the decoder (i.e., $T_{\theta} = d_{\theta}$) and employs the Cos cost. Results for encoder regularization (i.e., $T_{\theta} = e_{\phi}$) are reported




Figure 3: Stability analysis for the DST and the GMG applied to the decoder d_{θ} with cost Cos. For each regularizer, we assess stability by measuring the alignment of its gradients across 5 batches. We compute the cosine similarity between all pairs of gradients. Values closer to 1 indicates better alignment, hence better stability.

in Appendix D.1. We observe that the GMG's gradients exhibit significantly higher alignment compared to those of the DST, demonstrating greater stability of our proposed regularizer. This suggests that the GMG 's formulation, which accounts for the minimal distortion, effectively mitigates the inherent variability inherent of the distortion, leading to more stable gradient computations.

5 Conclusion

In this work, we introduce an OT perspective on unsupervised disentangled representation learning to incorporate general latent geometrical constraints. We derive the GMG, a provably weakly convex regularizer that measures whether a map T transports a fixed reference distribution with minimal distortion of some predefined geometric features. By formulating disentangled representation learning as a transport problem, we integrate the GMG into standard training objectives, allowing for incorporating and studying various geometric constraints on the learned representation spaces. We show significant performance benefits of our approach on four standard disentanglement benchmarks.

6 REPRODUCIBLITY

In this work, we introduce the GMG, computed as detailed in Algorithm 1. To facilitate reproducibility, we provide the implementation code for computing the GMG on source and target batches in Appendix E. Comprehensive proofs, including all underlying assumptions, are presented in Appendix B. For our experiments on disentanglement benchmarks, we adhere to standard practices, employing streamlined preprocessing across all datasets. Detailed descriptions of these procedures are available in Appendix C. All experiments described in this paper can be conducted using a single RTX 2080TI GPU, ensuring accessibility and replicability of our results.

ACKNOWLEDGEMENTS

Co-funded by the European Union (ERC, DeepCell - 101054957). Views and opinions expressed are, however, those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them. Luca Eyring and Karsten Roth thank the European Laboratory for Learning and Intelligent Systems (ELLIS) PhD program for support. Karsten Roth also thanks the International Max Planck Research School for Intelligent Systems (IMPRS-IS) for support. Zeynep Akata was supported by BMBF FKZ: 01IS18039A, by the ERC (853489 - DEXIM), by EXC number 2064/1 – project number 390727645. Fabian J. Theis consults for Immunai Inc., Singularity Bio B.V., CytoReason Ltd, Cellarity, and has ownership interest in Dermagnostix GmbH and Cellarity.

REFERENCES

Yoshua Bengio, Aaron Courville, and Pascal Vincent. Representation learning: A review and new perspectives, 2014. 1

Irina Higgins, David Amos, David Pfau, Sebastien Racaniere, Loic Matthey, Danilo Rezende, and Alexander Lerchner. Towards a definition of disentangled representations, 2018. 1

Francesco Locatello, Stefan Bauer, Mario Lucic, Gunnar Raetsch, Sylvain Gelly, Bernhard Schölkopf, and Olivier Bachem. Challenging common assumptions in the unsupervised learning of disentangled representations. In Kamalika Chaudhuri and Ruslan Salakhutdinov, editors, *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pages 4114–4124. PMLR, 09–15 Jun 2019a. URL https://proceedings.mlr.press/v97/locatello19a.html. 1, 2, 9, 26

Francesco Locatello, Ben Poole, Gunnar Raetsch, Bernhard Schölkopf, Olivier Bachem, and Michael Tschannen. Weakly-supervised disentanglement without compromises. In Hal Daumé III and Aarti Singh, editors, *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 6348–6359. PMLR, 13–18 Jul 2020. URL https://proceedings.mlr.press/v119/locatello20a.html. 1, 2, 9, 26

Karsten Roth, Mark Ibrahim, Zeynep Akata, Pascal Vincent, and Diane Bouchacourt. Disentanglement of correlated factors via hausdorff factorized support. In *The Eleventh International Conference on Learning Representations*, 2023. URL https://openreview.net/forum?id=OKcJhpQiGiX. 1, 2, 8, 9, 26

- Kyle Hsu, Will Dorrell, James C. R. Whittington, Jiajun Wu, and Chelsea Finn. Disentanglement via latent quantization. In *Thirty-seventh Conference on Neural Information Processing Systems*, 2023. URL https://openreview.net/forum?id=LLETO26Ga2. 1
- Vitória Barin-Pacela, Kartik Ahuja, Simon Lacoste-Julien, and Pascal Vincent. On the identifiability of quantized factors, 2024. 1
- F. Locatello, G. Abbati, T. Rainforth, S. Bauer, B. Schölkopf, and O. Bachem. On the fairness of disentangled representations. In *Advances in Neural Information Processing Systems 32 (NeurIPS 2019)*, pages 14584–14597. Curran Associates, Inc., December 2019b. URL https://papers.nips.cc/paper/9603-on-the-fairness-of-disentangled-representations. 1
- Frederik Träuble, Elliot Creager, Niki Kilbertus, Francesco Locatello, Andrea Dittadi, Anirudh Goyal, Bernhard Schölkopf, and Stefan Bauer. On disentangled representations learned from correlated data. In Marina Meila and Tong Zhang, editors, *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pages 10401–10412. PMLR, 18–24 Jul 2021. URL https://proceedings.mlr.press/v139/trauble21a.html. 1, 2
- Michal Rolinek, Dominik Zietlow, and Georg Martius. Variational autoencoders pursue pca directions (by accident). In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, June 2019. 1, 3
- Dominik Zietlow, Michal Rolinek, and Georg Martius. Demystifying inductive biases for (beta-)vae based architectures. In Marina Meila and Tong Zhang, editors, *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pages 12945–12954. PMLR, 18–24 Jul 2021. URL https://proceedings.mlr.press/v139/zietlow21a.html. 1
- Diederik P. Kingma and Max Welling. Auto-Encoding Variational Bayes. In 2nd International Conference on Learning Representations, ICLR 2014, Banff, AB, Canada, April 14-16, 2014, Conference Track Proceedings, 2014. 1, 2
- Irina Higgins, Loic Matthey, Arka Pal, Christopher Burgess, Xavier Glorot, Matthew Botvinick, Shakir Mohamed, and Alexander Lerchner. beta-VAE: Learning basic visual concepts with a constrained variational framework. In *International Conference on Learning Representations*, 2017. URL https://openreview.net/forum?id=Sy2fzU9gl. 1, 2, 9, 10, 27
- Hyunjik Kim and Andriy Mnih. Disentangling by factorising. In Jennifer Dy and Andreas Krause, editors, *Proceedings of the 35th International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, pages 2649–2658. PMLR, 10–15 Jul 2018. URL https://proceedings.mlr.press/v80/kim18b.html. 1, 3, 9
- Abhishek Kumar, Prasanna Sattigeri, and Avinash Balakrishnan. VARIATIONAL INFERENCE OF DISENTANGLED LATENT CONCEPTS FROM UNLABELED OBSERVATIONS. In *International Conference on Learning Representations*, 2018. URL https://openreview.net/forum?id=H1kG7GZAW. 1, 3
- Christopher P. Burgess, Irina Higgins, Arka Pal, Loïc Matthey, Nick Watters, Guillaume Desjardins, and Alexander Lerchner. Understanding disentangling in beta-vae. *CoRR*, abs/1804.03599, 2018. URL http://arxiv.org/abs/1804.03599. 1, 3
- Ricky T. Q. Chen, Xuechen Li, Roger B Grosse, and David K Duvenaud. Isolating sources of disentanglement in variational autoencoders. In S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 31. Curran Associates, Inc., 2018. URL https://proceedings.neurips.cc/paper/2018/file/lee3dfcd8a0645a25a35977997223d22-Paper.pdf. 1, 3, 8
- Amos Gropp, Matan Atzmon, and Yaron Lipman. Isometric autoencoders, 2020. 1
- Nutan Chen, Alexej Klushyn, Francesco Ferroni, Justin Bayer, and Patrick van der Smagt. Learning flat latent manifolds with vaes, 2020a. 1

- Yonghyeon Lee, Sangwoong Yoon, MinJun Son, and Frank C. Park. Regularized autoencoders for isometric representation learning. In *International Conference on Learning Representations*, 2022. URL https://openreview.net/forum?id=mQxt817JL04. 1, 3, 9, 10
- Daniella Horan, Eitan Richardson, and Yair Weiss. When is unsupervised disentanglement possible? In A. Beygelzimer, Y. Dauphin, P. Liang, and J. Wortman Vaughan, editors, *Advances in Neural Information Processing Systems*, 2021. URL https://openreview.net/forum?id=XqEF9riB93S. 1, 3
- Nao Nakagawa, Ren Togo, Takahiro Ogawa, and Miki Haseyama. Gromov-wasserstein autoencoders, 2023. 1, 3, 4, 9, 10
- In Huh, changwook jeong, Jae Myung Choe, Young-Gu Kim, and Dae Sin Kim. Isometric quotient variational auto-encoders for structure-preserving representation learning. In *Thirty-seventh Conference on Neural Information Processing Systems*, 2023. URL https://openreview.net/forum?id=EdgPb3ngR4. 1, 3, 9
- Jaehoon Hahm, Junho Lee, Sunghyun Kim, and Joonseok Lee. Isometric representation learning for disentangled latent space of diffusion models, 2024. URL https://arxiv.org/abs/2407. 11451. 1
- Filippo Santambrogio. Optimal Transport for Applied Mathematicians. *Birkhäuser, NY*, 55(58-63): 94, 2015. 2, 19
- Gabriel Peyré and Marco Cuturi. Computational Optimal Transport. Foundations and Trends in Machine Learning, 11(5-6), 2019. ISSN 1935-8245. 2, 3
- Karl-Theodor Sturm. The space of spaces: curvature bounds and gradient flows on the space of metric measure spaces, 2020. 2, 3
- Facundo Mémoli. Gromov-wasserstein distances and the metric approach to object matching. *Foundations of computational mathematics*, 11:417–487, 2011. 2, 3, 21
- Théo Uscidda and Marco Cuturi. The monge gap: A regularizer to learn all transport maps, 2023. 2, 5
- Michael Moor, Max Horn, Bastian Rieck, and Karsten Borgwardt. Topological autoencoders, 2021. URL https://arxiv.org/abs/1906.00722.3
- Nikita Balabin, Daria Voronkova, Ilya Trofimov, Evgeny Burnaev, and Serguei Barannikov. Disentanglement learning via topology, 2024. URL https://arxiv.org/abs/2308.12696.3
- Tero Karras, Samuli Laine, Miika Aittala, Janne Hellsten, Jaakko Lehtinen, and Timo Aila. Analyzing and improving the image quality of stylegan, 2020. URL https://arxiv.org/abs/1912.04958.3
- Ilya Tolstikhin, Olivier Bousquet, Sylvain Gelly, and Bernhard Schoelkopf. Wasserstein autoencoders, 2019. URL https://arxiv.org/abs/1711.01558.3
- Facundo Mémoli and Tom Needham. Comparison results for gromov-wasserstein and gromov-monge distances, 2022. 3
- Marco Cuturi. Sinkhorn Distances: Lightspeed Computation of Optimal Transport. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 26, 2013. 3, 6
- Gabriel Peyré, Marco Cuturi, and Justin Solomon. Gromov-wasserstein averaging of kernel and distance matrices. In *International Conference on Machine Learning*, pages 2664–2672, 2016. 3, 6
- Gabriel Rioux, Ziv Goldfeld, and Kengo Kato. Entropic gromov-wasserstein distances: Stability, algorithms, and distributional limits, 2023. 3, 21
- Zhengxin Zhang, Ziv Goldfeld, Youssef Mroueh, and Bharath K. Sriperumbudur. Gromov-wasserstein distances: Entropic regularization, duality, and sample complexity, 2023. 3, 7

- Meyer Scetbon, Gabriel Peyré, and Marco Cuturi. Linear-time gromov wasserstein distances using low rank couplings and costs. In *International Conference on Machine Learning*, pages 19347–19365. PMLR, 2022. 3, 6
- Ashok Makkuva, Amirhossein Taghvaei, Sewoong Oh, and Jason Lee. Optimal transport mapping via input convex neural networks. In *International Conference on Machine Learning (ICML)*, volume 37, 2020. 3
- Alexander Korotin, Daniil Selikhanovych, and Evgeny Burnaev. Neural optimal transport. 2022. doi: 10.48550/ARXIV.2201.12220. URL https://arxiv.org/abs/2201.12220. 3
- Alexander Tong, Nikolay Malkin, Guillaume Huguet, Yanlei Zhang, Jarrid Rector-Brooks, Kilian Fatras, Guy Wolf, and Yoshua Bengio. Improving and generalizing flow-based generative models with minibatch optimal transport, 2023. 3
- Luca Eyring, Dominik Klein, Théo Uscidda, Giovanni Palla, Niki Kilbertus, Zeynep Akata, and Fabian Theis. Unbalancedness in neural monge maps improves unpaired domain translation, 2024. 3
- Xavier Aramayo Carrasco, Maksim Nekrashevich, Petr Mokrov, Evgeny Burnaev, and Alexander Korotin. Uncovering challenges of solving the continuous gromov-wasserstein problem, 2024. URL https://arxiv.org/abs/2303.05978.3
- Dominik Klein, Théo Uscidda, Fabian Theis, and Marco Cuturi. Entropic (gromov) wasserstein flow matching with genot, 2024. 3
- Athina Sotiropoulou and David Alvarez-Melis. Strongly isomorphic neural optimal transport across incomparable spaces, 2024. URL https://arxiv.org/abs/2407.14957. 3
- Théo Dumont, Théo Lacombe, and François-Xavier Vialard. On the existence of monge maps for the gromov-wasserstein problem. 2022. 5
- Marco Cuturi, Laetitia Meng-Papaxanthos, Yingtao Tian, Charlotte Bunne, Geoff Davis, and Olivier Teboul. Optimal Transport Tools (OTT): A JAX Toolbox for all things Wasserstein. arXiv Preprint arXiv:2201.12324, 2022. 6
- Damek Davis, Dmitriy Drusvyatskiy, Kellie J. MacPhee, and Courtney Paquette. Subgradient methods for sharp weakly convex functions, 2018. 7
- Jonathan Weed and Francis Bach. Sharp asymptotic and finite-sample rates of convergence of empirical measures in wasserstein distance, 2017. URL https://arxiv.org/abs/1707.00087.7
- Jean Feydy, Thibault Séjourné, François-Xavier Vialard, Shun-Ichi Amari, Alain Trouvé, and Gabriel Peyré. Interpolating between Optimal Transport and MMD using Sinkhorn Divergences. In *International Conference on Artificial Intelligence and Statistics (AISTATS)*, volume 22, 2019. 8
- Cian Eastwood and Christopher K. I. Williams. A framework for the quantitative evaluation of disentangled representations. In *International Conference on Learning Representations*, 2018. URL https://openreview.net/forum?id=By-7dz-AZ.9
- Andrea Dittadi, Frederik Träuble, Francesco Locatello, Manuel Wuthrich, Vaibhav Agrawal, Ole Winther, Stefan Bauer, and Bernhard Schölkopf. On the transfer of disentangled representations in realistic settings. In *International Conference on Learning Representations*, 2021. URL https://openreview.net/forum?id=8VXvj1QNR11.9
- Yann LeCun, Fu Jie Huang, and Léon Bottou. Learning methods for generic object recognition with invariance to pose and lighting. *Proceedings of the IEEE Computer Society Conference on Computer Vision and Pattern Recognition*, 2:II97–II104, 2004. ISSN 1063-6919. Proceedings of the 2004 IEEE Computer Society Conference on Computer Vision and Pattern Recognition, CVPR 2004; Conference date: 27-06-2004 Through 02-07-2004. 9, 10

- Scott E Reed, Yi Zhang, Yuting Zhang, and Honglak Lee. Deep visual analogy-making. In C. Cortes, N. Lawrence, D. Lee, M. Sugiyama, and R. Garnett, editors, Advances in Neural Information Processing Systems, volume 28. Curran Associates, Inc., 2015. URL https://proceedings.neurips.cc/paper/2015/file/e07413354875be01a996dc560274708e-Paper.pdf. 9, 10
- Gaspard Monge. Mémoire sur la théorie des déblais et des remblais. *Histoire de l'Académie Royale des Sciences*, pages 666–704, 1781. 17
- L Kantorovich. On the transfer of masses (in russian). In *Doklady Akademii Nauk*, volume 37, page 227, 1942. 17
- Thibault Séjourné, François-Xavier Vialard, and Gabriel Peyré. The unbalanced gromov wasserstein distance: Conic formulation and relaxation, 2023. 19, 20
- Garrett Birkhoff. Tres observaciones sobre el algebra lineal. *Universidad Nacional de Tucumán Revista Series A*, 5:147–151, 1946. 20
- D. Bertsimas and J.N. Tsitsiklis. Introduction to linear optimization. Athena Scientific, 1997. 20
- Jean-François Le Gall. Intégration, Probabilités et Processus Aléatoires. 21
- Tudor Manole and Jonathan Niles-Weed. Sharp convergence rates for empirical optimal transport with smooth costs. *The Annals of Applied Probability*, 34(1B), February 2024. ISSN 1050-5164. doi: 10.1214/23-aap1986. URL http://dx.doi.org/10.1214/23-AAP1986. 22
- Stephen Boyd and Lieven Vandenberghe. *Convex Optimization*. Cambridge University Press, March 2004. ISBN 0521833787. URL http://www.amazon.com/exec/obidos/redirect?tag=citeulike-20&path=ASIN/0521833787. 22
- K. B. Petersen and M. S. Pedersen. The matrix cookbook, October 2008. URL http://www2.imm.dtu.dk/pubdb/p.php?3274. Version 20081110. 23, 25
- Igor Babuschkin, Kate Baumli, Alison Bell, Surya Bhupatiraju, Jake Bruce, Peter Buchlovsky, David Budden, Trevor Cai, Aidan Clark, Ivo Danihelka, Claudio Fantacci, Jonathan Godwin, Chris Jones, Ross Hemsley, Tom Hennigan, Matteo Hessel, Shaobo Hou, Steven Kapturowski, Thomas Keck, Iurii Kemaev, Michael King, Markus Kunesch, Lena Martens, Hamza Merzic, Vladimir Mikulik, Tamara Norman, John Quan, George Papamakarios, Roman Ring, Francisco Ruiz, Alvaro Sanchez, Rosalia Schneider, Eren Sezener, Stephen Spencer, Srivatsan Srinivasan, Luyu Wang, Wojciech Stokowiec, and Fabio Viola. The DeepMind JAX Ecosystem, 2020. URL http://github.com/deepmind.25
- Diederik P Kingma and Jimmy Ba. Adam: A Method for Stochastic Optimization. In *International Conference on Learning Representations (ICLR)*, 2014. 26
- Chris Burgess and Hyunjik Kim. 3d shapes dataset. https://github.com/deepmind/3dshapes-dataset/, 2018. 28
- Andrea Burns, Aaron Sarna, Dilip Krishnan, and Aaron Maschinot. Unsupervised disentanglement without autoencoding: Pitfalls and future directions. *arXiv preprint arXiv:2108.06613*, 2021. 28
- Julius von Kügelgen, Yash Sharma, Luigi Gresele, Wieland Brendel, Bernhard Schölkopf, Michel Besserve, and Francesco Locatello. Self-supervised learning with data augmentations provably isolates content from style. In *Advances in Neural Information Processing Systems*, 2021. 28
- Cian Eastwood, Julius von Kügelgen, Linus Ericsson, Diane Bouchacourt, Pascal Vincent, Mark Ibrahim, and Bernhard Schölkopf. Self-supervised disentanglement by leveraging structure in data augmentations. In *Causal Representation Learning Workshop at NeurIPS 2023*, 2023. URL https://openreview.net/forum?id=JoISqbH8vl. 28
- Stefan Matthes, Zhiwei Han, and Hao Shen. Towards a unified framework of contrastive learning for disentangled representations. In *Thirty-seventh Conference on Neural Information Processing Systems*, 2023. URL https://openreview.net/forum?id=QrB38MAAEP. 28

- Laurence Aitchison and Stoil Krasimirov Ganev. InfoNCE is variational inference in a recognition parameterised model. *Transactions on Machine Learning Research*, 2024. ISSN 2835-8856. URL https://openreview.net/forum?id=chbRsWwjax. 28
- Ting Chen, Simon Kornblith, Mohammad Norouzi, and Geoffrey Hinton. A simple framework for contrastive learning of visual representations. In Hal Daumé III and Aarti Singh, editors, *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 1597–1607. PMLR, 13–18 Jul 2020b. URL https://proceedings.mlr.press/v119/chen20j.html. 28
- Jure Zbontar, Li Jing, Ishan Misra, Yann LeCun, and Stephane Deny. Barlow twins: Self-supervised learning via redundancy reduction. In Marina Meila and Tong Zhang, editors, *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pages 12310–12320. PMLR, 18–24 Jul 2021. URL https://proceedings.mlr.press/v139/zbontar21a.html. 28
- Adrien Bardes, Jean Ponce, and Yann LeCun. VICReg: Variance-invariance-covariance regularization for self-supervised learning. In *International Conference on Learning Representations*, 2022. URL https://openreview.net/forum?id=xm6YD62D1Ub. 28
- Quentin Garrido, Yubei Chen, Adrien Bardes, Laurent Najman, and Yann LeCun. On the duality between contrastive and non-contrastive self-supervised learning. In *The Eleventh International Conference on Learning Representations*, 2023. URL https://openreview.net/forum?id=kDEL91Dufpa. 28

APPENDIX

The Appendix is organized as follows:

- Section A provides additional background information to supplement the main text.
- Section B presents all theoretical proofs, including detailed assumptions.
- Section C outlines comprehensive experimental details, ensuring reproducibility.
- Section D offers supplementary empirical results that further support our findings.
- Section E includes the implementation code for computing the GMG.

A ADDITIONAL BACKGROUND

A.1 REMINDERS ON MONGE AND KANTOROVICH OT

In this section, we recall the Monge and Kantorovich formulations of OT, which we will use to prove various results. These are the classical formulations of OT. Although we introduce them here after discussing the Gromov-Monge and Gromov-Wasserstein formulations, it should be noted that they are generally introduced beforehand. Indeed, the Gromov-Monge and Gromov-Wasserstein formulations were historically developed to derive OT formulations for comparing measures supported on incomparable spaces.

Monge Formulation. Instead of intra-domain cost functions, we consider here an *inter-domain* continuous cost function $c: \mathcal{X} \times \mathcal{Y} \to \mathbb{R}$. This assumes that we have a meaningful way to compare elements \mathbf{x}, \mathbf{y} from the source and target domains. The Monge (1781) problem (MP) between $p \in \mathcal{P}(\mathcal{X})$ and $p \in \mathcal{P}(\mathcal{Y})$ consists of finding a map $T: \mathcal{X} \to \mathcal{Y}$ that push-forwards p onto p, while minimizing the average displacement cost quantified by c

$$\inf_{T:T \sharp p = p} \int_{\mathcal{X}} c(\mathbf{x}, T(\mathbf{x})) \, \mathrm{d}p(\mathbf{x}) \,. \tag{MP}$$

We call any solution T^* to this problem a Monge map between p and q for cost c. Similarly to the Gromov-Monge Problem (GMP), solving the Monge Problem (MP) is difficult, as the constraint set is not convex and might be empty, especially when p, q are discrete.

Kantorovich Formulation. Instead of transport maps, the Kantorovich problem (KP) seeks a couplings $\pi \in \Pi(p,q)$:

$$W(p,q) := \min_{\pi \in \Pi(p,q)} \int_{\mathcal{X} \times \mathcal{Y}} c(\mathbf{x}, \mathbf{y}) \, d\pi(\mathbf{x}, \mathbf{y}).$$
 (KP)

An optimal coupling π^* solution of (KP), always exists. Studying the equivalence between (MP) and (KP) is easier than in the Gromov-Monge and Gromov-Wasserstein cases. Indeed, when (MP) is feasible, the Monge and Kantorovich formulations coincide and $\pi^* = (\operatorname{Id}, T^*) \sharp p$.

A.2 CONDITIONALLY POSITIVE KERNELS

In this section, we recall the definition of a conditionally positive kernel, which is involved in multiple proofs relying on the linearization of the Gromov-Wasserstein problem as a Kantorovich problem.

Definition A.1. A kernel $k : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ is CPD, i.e., conditionally positive, if it is symmetric and for any $\mathbf{x}_1, ..., \mathbf{x}_n \in \mathbb{R}^d$ and $\mathbf{a} \in \mathbb{R}^n$ s.t. $\mathbf{a}^\top \mathbf{1}_n = 0$, one has

$$\sum_{i,j=1}^{n} \mathbf{a}_i \mathbf{a}_j \, k(\mathbf{x}_i, \mathbf{x}_j) \ge 0$$

CPD include all positive kernels, such as the inner-product $k(\mathbf{x}, \mathbf{x}') = \langle \mathbf{x}, \mathbf{x}' \rangle$, or the cosine similarity $k(\mathbf{x}, \mathbf{x}') = \operatorname{cos-sim}(\mathbf{x}, \mathbf{x}') = \langle \frac{\mathbf{x}}{\|\mathbf{x}\|_2}, \frac{\mathbf{x}'}{\|\mathbf{x}'\|_2} \rangle$, but also the (scaled) negative squared Euclidean distance $k(\mathbf{x}, \mathbf{x}') = -\alpha \|\mathbf{x} - \mathbf{x}'\|_2^2$, $\alpha > 0$. Therefore, each of the costs of interest is either a conditionally positive kernel - for the inner product and the cosine distance - or its opposite is - for the squared Euclidean distance. Additionally, CPD kernels also include more exotic cost functions, such as:

- The RBF kernel $k(\mathbf{x}, \mathbf{x}') = \exp(-\|\mathbf{x} \mathbf{y}\|_2^2/\gamma), \gamma > 0.$
- The power kernel $k(\mathbf{x}, \mathbf{x}') = -\|\mathbf{x} \mathbf{x}'\|_2^p, 0$
- The thin plate spline kernel $k(\mathbf{x}, \mathbf{x}') = \|\mathbf{x} \mathbf{x}'\|_2^2 \log(\|\mathbf{x} \mathbf{x}')$
- The inverse multi-quadratic kernel $k(\mathbf{x}, \mathbf{x}') = 1/\sqrt{\|\mathbf{x} \mathbf{x}'\|_2^2 + c^2}, c \in \mathbb{R}$.

As a result, the family of CPD kernels includes a large variety of cost functions, which can be used to define various GMGs.

B PROOFS

B.1 THE GMG CHARACTERIZES GROMOV-MONGE OPTIMALITY

We show here that if $GMG_r(T) = 0$, then T is a Gromov-Monge map between r and $T\sharp r$ for costs $c_{\mathcal{X}}, c_{\mathcal{Y}}$. As the set of deterministic couplings $\{\pi_F := (I_d, F)\sharp r | F : \mathcal{X} \to \mathcal{Y}, F\sharp r = T\sharp r\} \subset \Pi(r, T\sharp r)$, we immediately get that

$$\inf_{F \sharp r = T \sharp r} \int_{\mathcal{X} \times \mathcal{X}} (c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}') - c_{\mathcal{Y}}(F(\mathbf{x}), F(\mathbf{x}')))^2 \, \mathrm{d}r(\mathbf{x}) \, \mathrm{d}r(\mathbf{x}') \ge \mathrm{GW}(r, T \sharp r) \tag{7}$$

On the other hand, if $GMG_r(T) = 0$, one has

$$GW(r, T \sharp r) = \int_{\mathcal{X} \times \mathcal{X}} (c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}') - c_{\mathcal{Y}}(T(\mathbf{x}), T(\mathbf{x}')))^{2} dr(\mathbf{x}) dr(\mathbf{x}')$$
(8)

Therefore, combining Eq. (7) and Eq. (8), we get

$$\inf_{F \sharp r = T \sharp r} \int_{\mathcal{X} \times \mathcal{X}} (c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}') - c_{\mathcal{Y}}(F(\mathbf{x}), F(\mathbf{x}')))^{2} dr(\mathbf{x}) dr(\mathbf{x}')$$

$$= \int_{\mathcal{X} \times \mathcal{X}} (c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}') - c_{\mathcal{Y}}(T(\mathbf{x}), T(\mathbf{x}')))^{2} dr(\mathbf{x}) dr(\mathbf{x}')$$
(9)

Finally, as T naturally satisfies the marginal constraint, we conclude that T is a Gromov-Monge map between r and $T \sharp r$ for costs $c_{\mathcal{X}}, c_{\mathcal{Y}}$.

B.2 ON RESCALING THE COSTS MATRICES IN THE ENTROPIC GW SOLVER

We remind, from Eq. (EGWP), that

$$GW_{\varepsilon}(p_n, q_n) = \min_{\mathbf{P} \in U_n} \sum_{i,j,i',j'=1}^{n} (\mathbf{C}_{\mathcal{X}_{i,i'}} - \mathbf{C}_{\mathcal{Y}_{j,j'}})^2 \mathbf{P}_{i,j} \mathbf{P}_{i',j'} - \varepsilon H(\mathbf{P}).$$
 (10)

By developping each terms, and using the fact that $P \in U_n$, we get

$$GW_{\varepsilon}(p_{n}, q_{n}) = \min_{\mathbf{P} \in U_{n}} \frac{1}{n^{2}} \langle \mathbf{C}_{\mathcal{X}}^{\odot 2} \mathbf{1}_{n}, \mathbf{1}_{n} \rangle + \frac{1}{n^{2}} \langle \mathbf{C}_{\mathcal{Y}}^{\odot 2} \mathbf{1}_{n}, \mathbf{1}_{n} \rangle - 2 \langle \mathbf{C}_{\mathcal{X}} \mathbf{P} \mathbf{C}_{\mathcal{Y}}, \mathbf{P} \rangle - \varepsilon H(\mathbf{P})$$

$$= \frac{1}{n^{2}} \langle \mathbf{C}_{\mathcal{X}}^{\odot 2} \mathbf{1}_{n}, \mathbf{1}_{n} \rangle + \frac{1}{n^{2}} \langle \mathbf{C}_{\mathcal{Y}}^{\odot 2} \mathbf{1}_{n}, \mathbf{1}_{n} \rangle + \min_{\mathbf{P} \in U_{n}} -2 \langle \mathbf{C}_{\mathcal{X}} \mathbf{P} \mathbf{C}_{\mathcal{Y}}, \mathbf{P} \rangle - \varepsilon H(\mathbf{P})$$

$$(11)$$

where $\mathbf{C}_{\mathcal{X}}^{\odot 2} = \mathbf{C}_{\mathcal{X}} \odot \mathbf{C}_{\mathcal{X}}$, with \odot the Hadamard (i.e., elementwise) product, and similarly for $\mathbf{C}_{\mathcal{Y}}^{\odot 2}$. As we can see that the two terms on the left do not depend on \mathbf{P} , they do not impact the minimization, an OT coupling \mathbf{P}^{\star} solving the problem satisfies:

$$\mathbf{P}^{\star} \in \underset{\mathbf{P} \in U_n}{\operatorname{arg \, min}} -2\langle \mathbf{C}_{\mathcal{X}} \mathbf{P} \mathbf{C}_{\mathcal{Y}}, \mathbf{P} \rangle - \varepsilon H(\mathbf{P}). \tag{12}$$

As a result, if we now replace $C_{\mathcal{X}}$ and $C_{\mathcal{Y}}$ by $C_{\mathcal{X}}/\text{stat}(C_{\mathcal{X}})$ and $C_{\mathcal{Y}}/\text{stat}(C_{\mathcal{Y}})$, respectively, the new OT coupling $\tilde{\mathbf{P}}^*$ solving the problem satisfies

$$\tilde{\mathbf{P}}^{\star} \in \underset{\mathbf{P} \in U_{n}}{\operatorname{arg \, min}} - 2\langle \frac{\mathbf{C}_{\mathcal{X}}}{\operatorname{stat}(\mathbf{C}_{\mathcal{X}})} \mathbf{P} \frac{\mathbf{C}_{\mathcal{Y}}}{\operatorname{stat}(\mathbf{C}_{\mathcal{Y}})}, \mathbf{P} \rangle - \varepsilon H(\mathbf{P})$$

$$\Leftrightarrow \quad \tilde{\mathbf{P}}^{\star} \in \underset{\mathbf{P} \in U_{n}}{\operatorname{arg \, min}} - \frac{2}{\operatorname{stat}(\mathbf{C}_{\mathcal{X}}) \cdot \operatorname{stat}(\mathbf{C}_{\mathcal{Y}})} \langle \mathbf{C}_{\mathcal{X}} \mathbf{P} \mathbf{C}_{\mathcal{Y}}, \mathbf{P} \rangle - \varepsilon H(\mathbf{P})$$

$$\Leftrightarrow \quad \tilde{\mathbf{P}}^{\star} \in \underset{\mathbf{P} \in U_{n}}{\operatorname{arg \, min}} - 2\langle \mathbf{C}_{\mathcal{X}} \mathbf{P} \mathbf{C}_{\mathcal{Y}}, \mathbf{P} \rangle - \operatorname{stat}(\mathbf{C}_{\mathcal{X}}) \cdot \operatorname{stat}(\mathbf{C}_{\mathcal{Y}}) \cdot \varepsilon H(\mathbf{P}),$$

$$(13)$$

where in the last line, we use the fact that $stat(\mathbf{C}_{\mathcal{X}})$, $stat(\mathbf{C}_{\mathcal{Y}}) > 0$. This yields the desired equivalence on scaling the cost matrix and adapting the entropic regularization strength.

B.3 Positivity of the Entropic GMG estimator

Recall that

$$\begin{aligned} \mathrm{GMG}_{r_n,\varepsilon}(T) &= \mathrm{DST}_{r_n}(T) - \mathrm{GW}_{\varepsilon}(r_n, T \sharp r_n) \\ &= \mathrm{DST}_{r_n}(T) - \min_{\mathbf{P} \in U_n} \sum_{i,j,i',j'=1}^n (c_{\mathcal{X}}(\mathbf{x}_i, \mathbf{x}_j) - c_{\mathcal{Y}}(\mathbf{y}_i, \mathbf{y}_j))^2 \mathbf{P}_{ij} \mathbf{P}_{i'j'} - \varepsilon H(\mathbf{P}) \,, \end{aligned}$$

For any coupling $\mathbf{P} \in U_n$, since $-\varepsilon H(\mathbf{P}) = -\varepsilon \sum_{i,j=1}^n \mathbf{P}_{ij} \log(\mathbf{P}_{ij}) < 0$, one has:

$$\sum_{i,j,i',j'=1}^{n}(c_{\mathcal{X}}(\mathbf{x}_i,\mathbf{x}_j)-c_{\mathcal{Y}}(\mathbf{y}_i,\mathbf{y}_j))^2\mathbf{P}_{ij}\mathbf{P}_{i'j'}-\varepsilon H(\mathbf{P})<\sum_{i,j,i',j'=1}^{n}(c_{\mathcal{X}}(\mathbf{x}_i,\mathbf{x}_j)-c_{\mathcal{Y}}(\mathbf{y}_i,\mathbf{y}_j))^2\mathbf{P}_{ij}\mathbf{P}_{i'j'}$$

As a result, applying minimization on both sides yields that $GW_{\varepsilon}(r_n, T\sharp r_n) < GW_0(r_n, T\sharp r_n) = GW(r_n, T\sharp r_n)$, and therefore:

$$\mathrm{GMG}_{r_n,\varepsilon}(T) > \mathrm{GMG}_{r_n,0}(T) = \mathrm{GMG}_{r_n}(T) \ge 0.$$

B.4 Proofs of Prop. 3.3

Proposition 3.3. If
$$GMG_r(T) = 0$$
, $\forall s \in \mathcal{P}(\mathcal{X})$ s.t. $Spt(s) \subseteq Spt(r)$, one has $GMG_s(T) = 0$.

Proof. Let T, r, s as described and suppose that $\mathcal{GM}_r^c(T) = 0$. Then, $\pi^r := (\mathrm{Id}, T) \sharp r$ is an optimal Gromov-Wasserstein coupling, solution of Problem (GWP) between r and $T \sharp r$ for costs $c_{\mathcal{X}}$ and $c_{\mathcal{Y}}$. Therefore, from (Séjourné et al., 2023, Theorem. 3), π^r is an optimal Kantorvich coupling, solution of Problem (KP) between r and $T \sharp r$ for the linearized cost:

$$\tilde{c}: (\mathbf{x}, \mathbf{y}) \in \mathcal{X} \times \mathcal{Y} \mapsto \int_{\mathcal{X} \times \mathcal{V}} \frac{1}{2} |c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}') - c_{\mathcal{Y}}(\mathbf{y}, \mathbf{y}')|^2 d\pi^r(\mathbf{x}', \mathbf{y}')$$
 (14)

Additionally, $\mathcal{X} \times \mathcal{Y}$ is a compact set as a product of compact sets, so since $(\mathbf{x}, \mathbf{y}) \mapsto |c_{\mathcal{X}}(\mathbf{x}, \mathbf{x}') - c_{\mathcal{Y}}(\mathbf{y}, \mathbf{y}')|^2$ is continuous as $c_{\mathcal{X}}$ and $c_{\mathcal{Y}}$ are continuous, it is bounded on $\mathcal{X} \times \mathcal{Y}$. Afterward, since π^r has finite mass, by Lebesgue's dominated convergence Theorem, it follows that \tilde{c} is continuous, and hence uniformly continuous, again since $\mathcal{X} \times \mathcal{Y}$ is compact.

Afterwards, by virtue of (Santambrogio, 2015, Theorem 1.38), $\operatorname{Spt}(\pi^r)$ is a \tilde{c} -cyclically monotone (CM) set (see (Santambrogio, 2015, Definition. 1.36)). From the definition of cyclical monotonicity, this property translates to subsets. Then, by defining $\pi^s = (\operatorname{Id}, T)\sharp s$, as $\operatorname{Spt}(p) \subset \operatorname{Spt}(r)$, one has $\operatorname{Spt}(\pi^s) = \operatorname{Spt}((\operatorname{Id}, T)\sharp s) \subset \operatorname{Spt}((\operatorname{Id}, T)\sharp r) = \operatorname{Spt}(\pi^r)$, so $\operatorname{Spt}(\pi^s)$ is \tilde{c} -CM. Finally, since $\mathcal X$ and $\mathcal Y$ are compact, and \tilde{c} is uniformly continuous, the \tilde{c} -cyclical monotonicity of its support implies that the coupling π^p is a Kantorovich optimal coupling between its marginals for cost \tilde{c} , thanks to (Santambrogio, 2015, Theorem 1.49). By re-applying (Séjourné et al., 2023, Theorem. 3), we get that π^s solves the Gromov-Wasserstein problem between its marginals for costs $c_{\mathcal X}$ and $c_{\mathcal Y}$. In other words, $\pi^s = (\operatorname{Id}, T)\sharp s$ is Gromov-Wasserstein optimal coupling between s and $T\sharp s$ so T is a Gromov-Monge map between s and $T\sharp s$ and $GMG_s(T)=0$.

B.5 Proofs of Thm. 3.5

Theorem 3.5. Both GMG_r^2 and $GMG_r^{\langle \cdot, \cdot \rangle}$, as well as their finite sample versions, are weakly convex.

- Finite sample. We note $\mathbf{X} \in \mathbb{R}^{n \times d}$ the matrix that stores the \mathbf{x}_i , i.e. the support of r_n , as rows. Then, (i) $\mathrm{GMG}_{r_n}^2$ and (ii) $\mathrm{GMG}_{r_n}^{\langle \cdot, \cdot \rangle}$ are respectively (i) $\gamma_{2,n}$ and (ii) $\gamma_{inner,n}$ -weakly convex, where: $\gamma_{inner,n} = \lambda_{\max}(\frac{1}{n}\mathbf{X}\mathbf{X}^\top) \lambda_{\min}(\frac{1}{n}\mathbf{X}\mathbf{X}^\top)$ and $\gamma_{2,n} = \gamma_{inner,n} + \max_{i=1...n} \|\mathbf{x}_i\|_2^2$.
- Asymptotic. (i) GMG_r^2 and (ii) $\mathrm{GMG}_r^{\langle \cdot, \cdot \rangle}$ are respectively (i) γ_2 and (ii) γ_{inner} -weakly convex, where: $\gamma_{inner} = \lambda_{\max}(\mathbb{E}_{\mathbf{x} \sim r}[\mathbf{x}\mathbf{x}^\top])$ and $\gamma_{2,n} = \gamma_{inner} + \max_{\mathbf{x} \in \mathrm{Spt}(r)} \|\mathbf{x}\|_2^2$.

Before proving Thm. 3.5, we first demonstrate some technical results that will be useful later.

Reformulation of the empirical GMG using permutations. We start by showing that $GMG_{r_n}(T)$ is always the sub-optimality gap of T in Prob. (GMP) between r_n and $T\sharp r_n$. This occurs because Prob. (GMP) and Prob. (GWP) coincide when applied between empirical measures on the same number of points. In other words, we can reformulate Prob. (GWP) between r_n and $T\sharp r_n$ using permutation matrices, instead of (plain) couplings.

Proposition B.1. The empirical GMG reads

$$GMG_{r_n}(T) = DST_{r_n}(T) - \min_{\sigma \in \mathcal{S}_n} \frac{1}{n^2} \sum_{i,j=1}^n \left(c_{\mathcal{X}}(\mathbf{x}_i, \mathbf{x}_j) - c_{\mathcal{Y}}(T(\mathbf{x}_{\sigma(i)}), T(\mathbf{x}_{\sigma(j)})) \right)^2$$
(15)

Proof. We first show a more general results, stating that when $c_{\mathcal{X}}, c_{\mathcal{Y}}$ are conditionally positive kernels (see A.1), the discrete GW couplings between uniform, empirical distributions supported on the same number of points, ae permutation matrices.

Proposition B.2 (Equivalence between Gromov-Monge and Gromov-Wasserstein problems in the discrete case.). Let $p_n = \frac{1}{n} \sum_{i=1}^n \delta_{\mathbf{x}_i}$ and $q_n = \frac{1}{n} \sum_{i=1}^n \delta_{\mathbf{y}_i}$ two uniform, empirical measures, supported on the same number of points. We denote by $P_n = \{\mathbf{P} \in \mathbb{R}^{n \times n}, \exists \sigma \in \mathcal{S}_n, \mathbf{P}_{ij} := \delta_{j,\sigma(i)}\}$ the set set of permutation matrices. Assume that $c_{\mathcal{X}}$ and $c_{\mathcal{Y}}$ (or $-c_{\mathcal{X}}$ and $-c_{\mathcal{Y}}$) are conditionally positive kernels (see A.1). Then, the GM and GW formulations coincide, in the sense that we can restrict the GW problem to permutations, namely

$$GW(p_n, p_n) = \min_{\mathbf{P} \in U_n} \sum_{i,j,i',j'=1}^{n} (c_{\mathcal{X}}(\mathbf{x}_i, \mathbf{x}_{i'}) - c_{\mathcal{Y}}(\mathbf{y}_j, \mathbf{y}_{j'}))^2 \mathbf{P}_{ij} \mathbf{P}_{i'j'}$$

$$= \frac{1}{n^2} \min_{\mathbf{P} \in P_n} \sum_{i,j,i',j'=1}^{n} (c_{\mathcal{X}}(\mathbf{x}_i, \mathbf{x}_{i'}) - c_{\mathcal{Y}}(\mathbf{y}_j, \mathbf{y}_{j'}))^2 \mathbf{P}_{ij} \mathbf{P}_{i'j'}$$

$$= \frac{1}{n^2} \min_{\sigma \in S_n} \sum_{i,j=1}^{n} (c_{\mathcal{X}}(\mathbf{x}_i, \mathbf{x}_j) - c_{\mathcal{Y}}(\mathbf{y}_{\sigma(i)}, \mathbf{y}_{\sigma(j)}))^2$$
(16)

Proof. Let $\mathbf{P}^{\star} \in U_n$ solution of the Gromov-Wasserstein between p_n and p_n , i.e.

$$\mathbf{P}^{\star} \in \operatorname*{arg\,min}_{\mathbf{P} \in U_n} \sum_{i,j,i',j'=1}^{n} (c_{\mathcal{X}}(\mathbf{x}_i, \mathbf{x}_{i'}) - c_{\mathcal{Y}}(\mathbf{y}_j, \mathbf{y}_{j'}))^2 \mathbf{P}_{ij} \mathbf{P}_{i'j'}$$

that always exists by continuity of the GW objective function on the compact U_n . We show that \mathbf{P}^* can be chosen as a (rescaled) permutation matrix without loss of generality.

As we assume that $c_{\mathcal{X}}$ and $c_{\mathcal{Y}}$ (or $-c_{\mathcal{X}}$ and $-c_{\mathcal{Y}}$) are conditionally positive kernels, from (Séjourné et al., 2023, Theorem. 3), \mathbf{P}^* also solves:

$$\mathbf{P}^{\star} \in \underset{\mathbf{Q} \in U_n}{\operatorname{arg \, min}} \sum_{i,j,i',j'=1}^{n} (c_{\mathcal{X}}(\mathbf{x}_i, \mathbf{x}_{i'}) - c_{\mathcal{Y}}(\mathbf{y}_j, \mathbf{y}_{j'}))^2 \mathbf{P}_{ij}^{\star} \mathbf{Q}_{i'j'}$$
(17)

We then define the linearized cost matrix $\tilde{C} \in \mathbb{R}^{n \times n}$, s.t.

$$\tilde{\mathbf{C}}_{ij} = \sum_{i',j'=1}^{n} (c_{\mathcal{X}}(\mathbf{x}_i, \mathbf{x}_{i'}) - c_{\mathcal{Y}}(\mathbf{y}_j, \mathbf{y}_{j'}))^2 \mathbf{P}_{ij}^{\star}$$

which allows us to reformulate Eq. (17) as

$$\mathbf{P}^{\star} \in \underset{\mathbf{Q} \in U_n}{\operatorname{arg\,min}} \langle \tilde{\mathbf{C}}, \mathbf{Q} \rangle \tag{18}$$

Birkhoff's theorem states that the extremal points of U_n are the permutation matrices P_n . Moreover, a seminal theorem of linear programming (Bertsimas and Tsitsiklis, 1997, Theorem 2.7) states that the minimum of a linear objective on a bounded polytope, if finite, is reached at an extremal point of the polyhedron. Therefore, as \mathbf{P}^* solves Eq. (18), it is an extremal point of U_n , so it can always be chosen as a permutation matrix. Therefore, the equivalence between GW and GM follows.

To conclude the proof of Prop. B.1, we simply remark that $r_n = \frac{1}{n} \sum_{i=1}^n \delta_{\mathbf{x}_i}$ and $T \sharp r_n = \frac{1}{n} \sum_{i=1}^n \delta_{T(\mathbf{x}_i)}$ are uniform, empirical distribution, and supported on the same number of points.

Consistency of the empirical GMG. We continue by proving a consistency result for the empirical GMG, which we will later use to deduce the asymptotic weak convexity constant from the finite-sample case.

Proposition B.3. For both $c_{\mathcal{X}} = c_{\mathcal{Y}} = \|\cdot - \cdot\|_2^2$ and $c_{\mathcal{X}} = c_{\mathcal{Y}} = \langle\cdot,\cdot\rangle$, one has $\mathrm{GMG}_{r_n}(T) \to \mathrm{GMG}_r(T)$ almost surely.

Proof. We first note that the empirical estimator of the distortion is consistent, as both costs are continuous, and \mathcal{X} is compact. We then need to study, in both cases, the convergence of $GW(r_n, T\sharp r_n)$ to $GW(r_n, T\sharp r)$.

To that end, we first remark that as, almost surely, $r_n \to r$ in distribution, one also has that, almost surely, $T\sharp r_n \to T\sharp r$ in distribution. Indeed, since $\mathcal Y$ is compact, T is bounded so for any bounded and continuous $f:\mathcal Y\to\mathbb R$ and $X\sim r,$ $f\circ T(X)$ is well defined and bounded so integrable. Afterwards, one can simply adapt the proof of the almost sure weak convergence of empirical measure based on the strong law of large numbers to show that, almost surely, $T\sharp r_n\to T\sharp r$ in distribution. See for instance (Le Gall, Theorem 10.4.1).

We start with the squared Euclidean distance. As, almost surely, both $r_n \to r$ and $T \sharp r_n \to T \sharp r$ in distribution, the results follows from (Mémoli, 2011, Thm 5.1, (e)).

We continue with the inner product. As noticed by Rioux et al. (2023, Lemma 2)—in the first version of the paper – the GW for inner product costs can be reformulated as:

$$GW^{\langle \cdot, \cdot \rangle}(p, q) = \int_{\mathcal{X} \times \mathcal{X}} \langle \mathbf{x}, \mathbf{x}' \rangle \, dp(\mathbf{x}) \, dp(\mathbf{x}') + \int_{\mathcal{Y} \times \mathcal{Y}} \langle \mathbf{y}, \mathbf{y}' \rangle \, dq(\mathbf{y}) \, dq(\mathbf{y}')$$

$$+ \min_{\mathbf{M} \in \mathcal{M}} \min_{\pi \in \Pi(p, q)} \int_{\mathcal{X} \times \mathcal{Y}} -4 \langle \mathbf{M} \mathbf{x}, \mathbf{y} \rangle \, d\pi(\mathbf{x}, \mathbf{y}) + 4 \|\mathbf{M}\|_{2}^{2},$$
(19)

where we define $\mathcal{M}=[-M/2,M/2]^{d_{\mathcal{X}}\times d_{\mathcal{Y}}}$ with $M=\sqrt{\int_{\mathcal{X}}\|\mathbf{x}\|_2^2\,\mathrm{d}p(\mathbf{x})\int_{\mathcal{Y}}\|\mathbf{y}\|_2^2\,\mathrm{d}q(\mathbf{y})}$. In particular, they show this result for the entropic GW problem with $\varepsilon>0$, but their proof is also valid for $\varepsilon=0$. The above terms only involving the marginal, i.e., not involved in the minimization, are naturally stable under convergence in distribution, as \mathcal{X} and \mathcal{Y} are compact, so as $\mathcal{X}\times\mathcal{X}$ and $\mathcal{Y}\times\mathcal{Y}$. As a result, we only need to study the stability of this quantity under the convergence in distribution of the following functional:

$$\mathcal{F}(p,q) = \min_{\mathbf{M} \in \mathcal{M}} \min_{\pi \in \Pi(p,q)} \int_{\mathcal{X} \times \mathcal{Y}} -4\langle \mathbf{M} \mathbf{x}, \mathbf{y} \rangle \, d\pi(\mathbf{x}, \mathbf{y}) + 4 \|\mathbf{M}\|_{2}^{2},$$
 (20)

We first remark that:

$$|\mathcal{F}(p,q) - \mathcal{F}(p_{n},q_{n})|$$

$$\leq \sup_{M \in \mathcal{M}} |\min_{\pi \in \Pi(p,q)} \int_{\mathcal{X} \times \mathcal{Y}} -4\langle \mathbf{M}\mathbf{x}, \mathbf{y} \rangle \, d\pi(\mathbf{x}, \mathbf{y}) - \min_{\pi \in \Pi(p,q)} \int_{\mathcal{X} \times \mathcal{Y}} -4\langle \mathbf{M}\mathbf{x}, \mathbf{y} \rangle \, d\pi(\mathbf{x}, \mathbf{y})|$$

$$\leq \sup_{M \in \mathcal{M}} |\min_{\pi \in \Pi(p,q)} \int_{\mathcal{X} \times \mathcal{Y}} 2\|\mathbf{M}\mathbf{x} - \mathbf{y}\|_{2}^{2} \, d\pi(\mathbf{x}, \mathbf{y}) - \min_{\pi \in \Pi(p_{n},q_{n})} \int_{\mathcal{X} \times \mathcal{Y}} 2\|\mathbf{M}\mathbf{x} - \mathbf{y}\|_{2}^{2} \, d\pi(\mathbf{x}, \mathbf{y})2|$$

$$+2 \cdot \sup_{M \in \mathcal{M}} |\int_{\mathcal{X}} \|\mathbf{M}\mathbf{x}\|_{2}^{2} \, dp(\mathbf{x}) - \int_{\mathcal{X}} \|\mathbf{M}\mathbf{x}\|_{2}^{2} \, dp_{n}(\mathbf{x})|$$

$$+2 \cdot |\int_{\mathcal{Y}} \|\mathbf{y}\|_{2}^{2} \, dq(\mathbf{y}) - \int_{\mathcal{Y}} \|\mathbf{y}\|_{2}^{2} \, dq_{n}(\mathbf{y})|$$

$$(21)$$

Then, we show the convergence of each term separately.

• For the first term, we remark that (up to a constant factor) it can be reformulated:

$$\sup_{M \in \mathcal{M}} |\mathbf{W}_2^2(\mathbf{M} \sharp p, q) - \mathbf{W}_2^2(\mathbf{M} \sharp p_n, q_n)|$$

where we remind that that W_2^2 is the (squared) Wasserstein distance, solution of Eq. (KP) induced by $c(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|_2^2$. By virtue of (Manole and Niles-Weed, 2024, Theorem 2), there exists a constant C > 0, s.t. we can uniformly bound

$$\sup_{M \in \mathcal{M}} |\mathbf{W}_2^2(\mathbf{M} \sharp p, q) - \mathbf{W}_2^2(\mathbf{M} \sharp p_n, q_n)| \le C n^{-1/d}$$

and the convergence follows.

- For the second one, this follows from from the convergence in distribution of p_n to p along with the Ascoli-Arzela theorem, since both \mathcal{M} and \mathcal{X} are compact sets, so the $\{f_{\mathbf{M}} \mid f_{\mathbf{M}} : \mathbf{x} \mapsto \|\mathbf{M}\mathbf{x}\|_2^2\}$ are uniformly bounded and equi-continuous.
- For the third one, this follows from the convergence in distribution of q_n to q.

As a result, we finally get $GW^{\langle \cdot, \cdot \rangle}(p_n, q_n) \to GW^{\langle \cdot, \cdot \rangle}(p, q)$.

Weak convexity. Finally, we demonstrate some useful results on weakly convex functions on \mathbb{R}^d . **Definition B.4.** A function $f: \mathbb{R}^d \to \mathbb{R}$ is γ -weakly convex if $f + \gamma \| \cdot \|_2^2$ is convex.

From the definition, we see that if f is γ -weakly convex, than f is also γ' weakly convex for any $\gamma' \geq \gamma$. This naturally extends to weakly convex functionals \mathcal{F} on $L_2(r)$.

Lemma B.5. Let $\mathbf{A} \in S_d(\mathbb{R})$ a symmetric matrix and define the quadratic form $f_{\mathbf{A}} : \mathbf{x} \in \mathbb{R}^d \mapsto \mathbf{x}^{\top} \mathbf{A} \mathbf{x}$. Then, $f_{\mathbf{A}}$ is $\max(0, -\lambda_{\min}(\mathbf{A}))$ -weakly convex.

Proof. We use the fact that a twice continuously differentiable function is convex i.f.f. its hessian is positive semi-definite (Boyd and Vandenberghe, 2004, §(3.1.4)). Therefore, $f_{\mathbf{A}}$ is convex i.f.f. $\nabla^2 f_{\mathbf{A}} = \mathbf{A} \geq 0$. If $\lambda_{\min}(\mathbf{A}) \geq 0$, then $\mathbf{A} \geq 0$ so $f_{\mathbf{A}}$ is convex, i.e. 0-weakly convex. Otherwise, $f_{\mathbf{A}} - \frac{1}{2}\lambda_{\min}(\mathbf{A})\|\cdot\|_2^2$ has hessian $A - \lambda_{\min}(\mathbf{A}) \geq 0$, so it is convex, which yields that $f_{\mathbf{A}}$ is $-\lambda_{\min}(\mathbf{A})$ -weakly convex.

Lemma B.6. Let $(f_i)_{i \in I}$ a family of γ -weakly convex functions, with potentially infinite I. Then, $f: \mathbf{x} \in \mathbb{R}^d \mapsto \sup_{i \in I} f_i(\mathbf{x})$ is γ -weakly convex.

Proof. As the f_i are γ -weakly convex, $f_i + \frac{1}{2}\gamma$ is convex, so $\mathbf{x} \mapsto \sup_{i \in I} f_i(\mathbf{x}) + \frac{1}{2}\gamma \|x\|_2^2 = (\sup_{i \in I} f_i(\mathbf{x})) + \frac{1}{2}\gamma \|x\|_2^2$ is convex (Boyd and Vandenberghe, 2004, Eq. (3.7)). Therefore, the γ -weak convexity of f follows

Let's now proceed to prove the main Thm. (3.5).

Proof of Thm. (3.5). **Finite sample**. We first study the weak convexity of $\mathcal{GM}_{r_n}^{\langle\cdot,\cdot\rangle}$, i.e. the Gromov-Monge gap for the inner product. For a map $T \in L_2(r)$, it reads

$$GMG_{r_n}^{\langle \cdot, \cdot \rangle}(T) = \frac{1}{n^2} \sum_{i,j=1}^{n} \frac{1}{2} |\langle \mathbf{x}_i, \mathbf{x}_j \rangle - \langle T(\mathbf{x}_i), T(\mathbf{x}_j) \rangle|^2$$
$$- \min_{\mathbf{P} \in U_n} \sum_{i,j,i',j'=1}^{n} \frac{1}{2} |\langle \mathbf{x}_i, \mathbf{x}_{i'} \rangle - \langle T(\mathbf{x}_j), T(\mathbf{x}_{j'}) \rangle|^2 \mathbf{P}_{ij} \mathbf{P}_{i'j'}$$

As r_n and $T \sharp r_n$ are uniform empirical supported on the same number of points, using Prop. B.2, we can reformulate the RHS with permutation matrices, which yields

$$GMG_{r_n}^{\langle \cdot, \cdot \rangle}(T) = \frac{1}{n^2} \sum_{i,j=1}^{n} \frac{1}{2} |\langle \mathbf{x}_i, \mathbf{x}_j \rangle - \langle T(\mathbf{x}_i), T(\mathbf{x}_j) \rangle|^2$$
$$- \frac{1}{n^2} \min_{\mathbf{P} \in P_n} \sum_{i,j,i',j'=1}^{n} \frac{1}{2} |\langle \mathbf{x}_i, \mathbf{x}_{i'} \rangle - \langle T(\mathbf{x}_j), T(\mathbf{x}_{j'}) \rangle|^2 \mathbf{P}_{ij} \mathbf{P}_{i'j'}$$

From this expression, $\mathrm{GMG}_{r_n}^{\langle \cdot, \cdot \rangle}$ can be reformulated as a matrix input function. Indeed, it only depends on the map T via its values on the support of r_n , namely $\mathbf{x}_1, ..., \mathbf{x}_n$. Therefore, we write $\mathbf{t}_i := T(\mathbf{x}_i)$, and define $\mathbf{X}, \mathbf{T} \in \mathbb{R}^{n \times d}$ which contain observations \mathbf{x}_i and \mathbf{t}_i respectively, stored as rows. Then, studying $\mathrm{GMG}_r^{\langle \cdot, \cdot \rangle}$ remains to study

$$f(\mathbf{T}) := \frac{1}{n^2} \sum_{i,j=1}^n \frac{1}{2} |\langle \mathbf{x}_i, \mathbf{x}_j \rangle - \langle \mathbf{t}_i, \mathbf{t}_j \rangle|^2 - \frac{1}{n^2} \min_{\mathbf{P} \in P_n} \sum_{i,j,i',j'=1}^n \frac{1}{2} |\langle \mathbf{x}_i, \mathbf{x}_{i'} \rangle - \langle \mathbf{t}_j, \mathbf{t}_{j'} \rangle|^2 \mathbf{P}_{ij} \mathbf{P}_{i'j'}$$

By developing each term and exploiting that for any $P \in P_n$, $P1_n = P^{\top}1_n = \frac{1}{n}1_n$, we derive

$$f(\mathbf{T}) = \frac{1}{n^2} \sum_{i,j=1}^{n} -\langle \mathbf{x}_i, \mathbf{x}_j \rangle \cdot \langle \mathbf{t}_i, \mathbf{t}_j \rangle - \min_{\mathbf{P} \in P_n} \frac{1}{n^2} \sum_{i,j,i',j'=1}^{n} -\langle \mathbf{x}_i, \mathbf{x}_{i'} \rangle \cdot \langle \mathbf{t}_j, \mathbf{t}_{j'} \rangle \mathbf{P}_{ij} \mathbf{P}_{i'j'}$$

$$= \max_{\mathbf{P} \in P_n} \frac{1}{n^2} \sum_{i,j,i',j'=1}^{n} \langle \mathbf{x}_i, \mathbf{x}_{i'} \rangle \cdot \langle \mathbf{t}_j, \mathbf{t}_{j'} \rangle \mathbf{P}_{ij} \mathbf{P}_{i'j'} - \frac{1}{n^2} \sum_{i,j=1}^{n} \langle \mathbf{x}_i, \mathbf{x}_j \rangle \cdot \langle \mathbf{t}_i, \mathbf{t}_j \rangle$$

$$= \max_{\mathbf{P} \in P_n} \langle \frac{1}{n^2} \mathbf{P}^\top \mathbf{X} \mathbf{X}^\top \mathbf{P}, \mathbf{T} \mathbf{T}^\top \rangle - \langle \frac{1}{n^2} \mathbf{X} \mathbf{X}^\top, \mathbf{T} \mathbf{T}^\top \rangle$$

$$= \max_{\mathbf{P} \in P_n} \langle \frac{1}{n^2} (\mathbf{P}^\top \mathbf{X} \mathbf{X}^\top \mathbf{P} - \mathbf{X} \mathbf{X}^\top), \mathbf{T} \mathbf{T} \rangle$$

$$= \max_{\mathbf{P} \in P_n} \langle \mathbf{A}_{\mathbf{X}, \mathbf{P}} \mathbf{T}, \mathbf{T} \rangle$$

$$= \max_{\mathbf{P} \in P_n} \langle \mathbf{A}_{\mathbf{X}, \mathbf{P}} \mathbf{T}, \mathbf{T} \rangle$$

where we define $\mathbf{A}_{\mathbf{X},\mathbf{P}} := \frac{1}{n^2} (\mathbf{P}^\top \mathbf{X} \mathbf{X}^\top \mathbf{P} - \mathbf{X} \mathbf{X}^\top) \in \mathbb{R}^{n \times n}$. To study the convexity of this matrix input function, we vectorize it. From (Petersen and Pedersen, 2008, Eq. (520)), we note that, for any $\mathbf{M} \in \mathbb{R}^{n \times n}$

$$\langle \mathbf{MT}, \mathbf{T} \rangle = \mathbf{vec}(\mathbf{T})^{\top} \mathbf{vec}(\mathbf{MT}) = \mathbf{vec}(\mathbf{T})^{\top} (\mathbf{M} \otimes I_n) \mathbf{vec}(\mathbf{T})$$

where **vec** is the vectorization operator, raveling a matrix along its rows, and \otimes is the Kronecker product. Applying this identity, we reformulate:

$$f(\mathbf{T}) = \max_{\mathbf{P} \in U_n} \mathbf{vec}(\mathbf{T})^{\top} (\mathbf{A}_{\mathbf{X}, \mathbf{P}} \otimes I_n) \mathbf{vec}(\mathbf{T})$$
(22)

To study the convexity of r, we study the convexity of each $r_{\mathbf{A}_{\mathbf{X},\mathbf{P}}}(\mathbf{T}) := \mathbf{vec}(\mathbf{T})^{\top}(\mathbf{A}_{\mathbf{X},\mathbf{P}} \otimes I_n)\mathbf{vec}(\mathbf{T})$, which are quadratic forms induced by the $\mathbf{A}_{\mathbf{X},\mathbf{P}} \otimes I_n$. This remains to study the (semi-) positive definiteness of the matrices $\mathbf{A}_{\mathbf{X},\mathbf{P}} \otimes I_n$. As each $\mathbf{A}_{\mathbf{X},\mathbf{P}} \in \mathbb{R}^{n \times n}$ is symmetric and square, $\mathbf{A}_{\mathbf{X},\mathbf{P}} \otimes I_n$ is also symmetric and from (Petersen and Pedersen, 2008, Eq. (519)) its eigenvalues are the outer products of the eigenvalues of $\mathbf{A}_{\mathbf{X},\mathbf{P}}$ and I_n , namely

$$\mathbf{eig}(\mathbf{A}_{\mathbf{X},\mathbf{P}} \otimes I_n) = \{\lambda_i(\mathbf{A}_{\mathbf{X},\mathbf{P}}) \cdot \lambda_j(I_n)\}_{1 \leq i,j \leq n} \\ = \{\underbrace{\lambda_1(\mathbf{A}_{\mathbf{X},\mathbf{P}}), \dots, \lambda_1(\mathbf{A}_{\mathbf{X},\mathbf{P}})}_{\text{n times}}, \dots, \underbrace{\lambda_n(\mathbf{A}_{\mathbf{X},\mathbf{P}}), \dots, \lambda_n(\mathbf{A}_{\mathbf{X},\mathbf{P}})}_{\text{n times}}\}$$
(23)

It follows that the minimal eigenvalue of $\mathbf{A}_{\mathbf{X},\mathbf{P}} \otimes I_n$ is $\lambda_{\min}(\mathbf{A}_{\mathbf{X},\mathbf{P}} \otimes I_n) = \lambda_{\min}(\mathbf{A}_{\mathbf{X},\mathbf{P}})$. Utilizing the expression of $\mathbf{A}_{\mathbf{X},\mathbf{P}}$

$$\lambda_{\min}(\mathbf{A}_{\mathbf{X},\mathbf{P}}) = \frac{1}{n^2} \lambda_{\min}(\mathbf{P}^{\top} \mathbf{X} \mathbf{X}^{\top} \mathbf{P} - \mathbf{X} \mathbf{X}^{\top})$$

$$\geq \frac{1}{n^2} (\lambda_{\min}(\mathbf{P}^{\top} \mathbf{X} \mathbf{X}^{\top} \mathbf{P}) + \lambda_{\min}(-\mathbf{X} \mathbf{X}^{\top}))$$

$$= \frac{1}{n^2} (\lambda_{\min}(\mathbf{P}^{\top} \mathbf{X} \mathbf{X}^{\top} \mathbf{P}) - \lambda_{\max}(\mathbf{X} \mathbf{X}^{\top}))$$
(24)

Reminding that $\mathbf{P} \in U_n$, one has $\mathbf{P}^{\top} = \mathbf{P}^{-1}$, so $\mathbf{P}^{\top} \mathbf{X} \mathbf{X}^{\top}$ and $\mathbf{X} \mathbf{X}^{\top}$ are similar, and they have the same eigenvalues. In particular $\lambda_{\min}(\mathbf{P}^{\top} \mathbf{X} \mathbf{X}^{\top} \mathbf{P}) = \lambda_{\min}(\mathbf{X} \mathbf{X}^{\top})$. Combining these results, it follows that

$$\lambda_{\min}(\mathbf{A}_{\mathbf{X},\mathbf{P}} \otimes I_n) = \lambda_{\min}(\mathbf{A}_{\mathbf{X},\mathbf{P}}) \ge \frac{1}{n^2} (\lambda_{\min}(\mathbf{X}\mathbf{X}^\top) - \lambda_{\max}(\mathbf{X}\mathbf{X}^\top))$$
(25)

We then remind that each $r_{\mathbf{A}\mathbf{X},\mathbf{P}}$ is the quadratic form defined by $\mathbf{A}_{\mathbf{X},\mathbf{P}}\otimes I_n$, so by applying Prop. B.5, it is $\mathbf{A}_{\mathbf{X},\mathbf{P}}\otimes I_n$ -weakly convex, and hence $\frac{1}{n^2}(\lambda_{\max}(\mathbf{X}\mathbf{X}^\top)-\lambda_{\min}(\mathbf{X}\mathbf{X}^\top))$ -weakly convex. Therefore, applying Prop. (B.6), r is $\frac{1}{n^2}(\lambda_{\max}(\mathbf{X}\mathbf{X}^\top)-\lambda_{\min}(\mathbf{X}\mathbf{X}^\top))$ -weakly convex, in \mathbb{R}^d . Reminding that $\gamma_{\mathrm{inner}}=\frac{1}{n}(\lambda_{\max}(\mathbf{X}\mathbf{X}^\top)-\lambda_{\min}(\mathbf{X}\mathbf{X}^\top))$, r is $\frac{1}{n}\gamma_{\mathrm{inner}}$ weakly convex. This implies that $\mathbf{T}\mapsto f(\mathbf{T})+\frac{1}{n}\gamma_{\mathrm{inner}}\|\mathbf{T}\|_2^2$ is convex. By reminding that \mathbf{T} stores the $T(\mathbf{x}_i)$ as rows, $\frac{1}{n}\|\mathbf{T}\|_2^2=\|T\|_{L_2(r_n)}$. Consequently, $\mathrm{GMG}_{r_n}^{\langle\cdot,\cdot\rangle}$ is γ_{inner} in $L_2(r_n)$.

We then study the convexity of $GMG_{r_n}^2$. We follow exactly the same approach. One has:

$$GMG_{r_n}^2(T) = \frac{1}{n^2} \sum_{i,j=1}^n \frac{1}{2} |\|\mathbf{x}_i - \mathbf{x}_j\|_2^2 - \|T(\mathbf{x}_i) - T(\mathbf{x}_j)\|_2^2|^2$$
$$- \frac{1}{n^2} \min_{\mathbf{P} \in P_n} \sum_{i,j,i',j'=1}^n \frac{1}{2} |\|\mathbf{x}_i - \mathbf{x}_j\|_2^2 - \|T(\mathbf{x}_i) - T(\mathbf{x}_j)\|_2^2|^2|^2 \mathbf{P}_{ij} \mathbf{P}_{i'j'}$$

Similarly, studying the convexity of $\mathrm{GMG}^2_{r_n}(T)$ remains to study the convexity of the matrix input function:

$$g(\mathbf{T}) := \frac{1}{n^2} \sum_{i,j=1}^n \frac{1}{2} |\|\mathbf{x}_i - \mathbf{x}_j\|_2^2 - \|\mathbf{t}_i - \mathbf{t}_j\|_2^2|^2$$
$$- \frac{1}{n^2} \min_{\mathbf{P} \in P_n} \sum_{i,j,i',j'=1}^n \frac{1}{2} |\|\mathbf{x}_i - \mathbf{x}_j\|_2^2 - \|\mathbf{t}_i - \mathbf{t}_j\|_2^2|^2 \mathbf{P}_{ij} \mathbf{P}_{i'j'}$$

As before, by developing each term, one has:

$$g(\mathbf{T}) = \max_{\mathbf{P} \in P_n} \frac{1}{n^2} \sum_{i,j,i',j'=1}^n \langle \mathbf{x}_i, \mathbf{x}_{i'} \rangle \cdot \langle \mathbf{t}_j, \mathbf{t}_{j'} \rangle \mathbf{P}_{ij} \mathbf{P}_{i'j'} + \frac{1}{2n} \sum_{i,j=1}^n \mathbf{P}_{ij} \|\mathbf{x}_i\|_2^2 \|\mathbf{t}_i\|_2^2$$
$$- \left(\frac{1}{n^2} \sum_{i,j=1}^n \langle \mathbf{x}_i, \mathbf{x}_j \rangle \cdot \langle \mathbf{t}_i, \mathbf{t}_j \rangle + \frac{1}{2n} \sum_{i,j=1}^n \|\mathbf{x}_i\|_2^2 \|\mathbf{t}_i\|_2^2 \right)$$

The quadratic terms in **P** can be factorized as before using $\mathbf{A}_{\mathbf{X},\mathbf{P}}$. For the new terms w.r.t. the inner product case, we introduce $\mathbf{D}_{\mathbf{X}} := \operatorname{diag}(\|\mathbf{x}_1\|_2^2, \dots, \|\mathbf{x}_n\|_2^2)$, and remark that we can rewrite:

$$\frac{1}{2n} \sum_{i,j=1}^{n} \mathbf{P}_{ij} \|\mathbf{x}_i\|_2^2 \|\mathbf{t}_i\|_2^2 - \frac{1}{2n} \sum_{i,j=1}^{n} \|\mathbf{x}_i\|_2^2 \|\mathbf{t}_i\|_2^2 = \mathbf{vec}(T)^\top \left(\frac{1}{2n} (\mathbf{P}^\top - I_n) \otimes \mathbf{D}_{\mathbf{X}}\right) \mathbf{vec}(T)$$

As we can always symetrize the matrix when considering its associated quadratic form, we have:

$$\frac{1}{2n} \sum_{i,j=1}^{n} \mathbf{P}_{ij} \|\mathbf{x}_i\|_2^2 \|\mathbf{t}_i\|_2^2 - \frac{1}{2n} \sum_{i,j=1}^{n} \|\mathbf{x}_i\|_2^2 \|\mathbf{t}_i\|_2^2 = \mathbf{vec}(T)^\top \left(\frac{1}{2} \left(\frac{1}{2n} (\mathbf{P}^\top + \mathbf{P}) - I_n\right) \otimes \mathbf{D}_{\mathbf{X}}\right) \mathbf{vec}(T)$$

As a result, we denote $\mathbf{B}_{\mathbf{X},\mathbf{P}} = \frac{1}{n} (\frac{1}{2} (\mathbf{P}^\top + \mathbf{P}) - I_n) \otimes \mathbf{D}_{\mathbf{X}}$ and finally get:

$$g(\mathbf{T}) = \max_{\mathbf{P} \in P_n} \mathbf{vec}(T)^{\top} (\mathbf{A}_{\mathbf{X},\mathbf{P}} \otimes I_n + \mathbf{B}_{\mathbf{X},\mathbf{P}}) \mathbf{vec}(T)$$

As we did for f, studying the weak convexity of f remains to lower bound the minimal eigenvalue of $\mathbf{A}_{\mathbf{X},\mathbf{P}}\otimes I_n+\mathbf{B}_{\mathbf{X},\mathbf{P}}$. First, one remark that:

$$\lambda_{\min}(\mathbf{A}_{\mathbf{X},\mathbf{P}} \otimes I_n + \mathbf{B}_{\mathbf{X},\mathbf{P}}) \ge \lambda_{\min}(\mathbf{A}_{\mathbf{X},\mathbf{P}} \otimes I_n) + \lambda_{\min}(\mathbf{B}_{\mathbf{X},\mathbf{P}})$$

As we we have already lower bounded $\lambda_{\min}(\mathbf{A}_{\mathbf{X},\mathbf{P}}\otimes I_n) \geq \frac{1}{n^2}(\lambda_{\min}(\mathbf{X}\mathbf{X}^\top) - \lambda_{\max}(\mathbf{X}\mathbf{X}^\top))$, we focus on the RHS. Similarly, one has:

$$\lambda_{\min}(\mathbf{B}_{\mathbf{X},\mathbf{P}}) = \lambda_{\min} \left(\frac{1}{2n} (\frac{1}{2} (\mathbf{P}^{\top} + \mathbf{P}) - I_n) \otimes \mathbf{D}_{\mathbf{X}} \right)$$

$$\geq \lambda_{\min} \left(\frac{1}{4n} (\mathbf{P}^{\top} + \mathbf{P}) \otimes \mathbf{D}_{\mathbf{X}} \right) + \lambda_{\min} \left(-\frac{1}{2n} I_n \otimes \mathbf{D}_{\mathbf{X}} \right)$$

$$\geq \lambda_{\min} \left(\frac{1}{4n} (\mathbf{P}^{\top} + \mathbf{P}) \otimes \mathbf{D}_{\mathbf{X}} \right) - \lambda_{\max} \left(\frac{1}{2n} I_n \otimes \mathbf{D}_{\mathbf{X}} \right)$$
(26)

For both terms, we apply again (Petersen and Pedersen, 2008, Eq. (519)). For the LHS, one has:

$$\operatorname{eig}\left(\frac{1}{4n}(\mathbf{P}^{\top} + \mathbf{P}) \otimes \mathbf{D}_{\mathbf{X}}\right) = \left\{\lambda_{i}\left(\frac{1}{4n}(\mathbf{P}^{\top} + \mathbf{P})\right)\lambda_{j}(\mathbf{D}_{\mathbf{X}}\right)\right\}_{1 \leq i, j \leq n}$$
(27)

We remark that $\frac{1}{2}(\mathbf{P}^{\top}+\mathbf{P})$ is a symetric bi-stochastic matrix, so $\lambda_{\min}(\frac{1}{2}(\mathbf{P}^{\top}+\mathbf{P})) \geq -1$. Therefore, $\lambda_{\min}(\frac{1}{4n}(\mathbf{P}^{\top}+\mathbf{P})) \geq -\frac{1}{2n}$. As a result, since the eigenvalues of $\mathbf{D}_{\mathbf{X}}$ are the $\|\mathbf{x}_i\|_2^2$, this yields:

$$\lambda_{\min} \left(\frac{1}{4n} (\mathbf{P}^{\top} + \mathbf{P}) \otimes \mathbf{D}_{\mathbf{X}} \right) \ge -\frac{1}{2n} \max_{i=1,\dots,n} \|\mathbf{x}_i\|_2^2$$

Similarly, we have:

$$-\lambda_{\max}\left(\frac{1}{2n}I_n\otimes\mathbf{D_X}\right)\geq -\frac{1}{2n}\max_{i=1,\dots,n}\|\mathbf{x}_i\|_2^2$$

from which we deduce that:

$$\lambda_{\min}(\mathbf{B}_{\mathbf{X},\mathbf{P}}) \ge -\frac{1}{n} \max_{i=1,\dots,n} \|\mathbf{x}_i\|_2^2$$

We can then lower bound:

$$\lambda_{\min}(\mathbf{A}_{\mathbf{X},\mathbf{P}} \otimes I_n + \mathbf{B}_{\mathbf{X},\mathbf{P}}) \ge \frac{1}{n^2} (\lambda_{\min}(\mathbf{X}\mathbf{X}^{\top}) - \lambda_{\max}(\mathbf{X}\mathbf{X}^{\top})) - \frac{1}{n} \max_{i=1,\dots,n} \|\mathbf{x}_i\|_2^2$$

$$= -\frac{1}{n} \gamma_{2,n}$$
(28)

which yields the $\frac{1}{n}\gamma_{2,n}$ -weak convexity of g, and finally the $\gamma_{2,n}$ -weak convexity of $\mathcal{GM}_{r_n}^2$.

Asymptotic. For any T, we note that, almost surely, $||T||_{L_2(r_n)}^2 \to ||T||_{L_2(r)}^2$. As a result, since convexity is preserved under pointwise convergence and by virtue of Prop. (B.3), we study the (almost sure) convergence of $\gamma_{\text{inner},n}$ and $\gamma_{2,n}$.

We start by $\gamma_{\text{inner},n}$. We first remark that $\lambda_{\max}(\frac{1}{n}\mathbf{X}\mathbf{X}^{\top}) = \lambda_{\max}(\frac{1}{n}\mathbf{X}^{\top}\mathbf{X})$. Moreover, as $\mathbf{A} \in S_d^+(\mathbb{R}) \mapsto \lambda_{\max}(\mathbf{A})$ is continuous and $\frac{1}{n}\mathbf{X}^{\top}\mathbf{X} \to \mathbb{E}_{\mathbf{x} \sim r}[\mathbf{x}\mathbf{x}^{\top}]$ almost surely, one has $\lambda_{\max}(\frac{1}{n}\mathbf{X}\mathbf{X}^{\top}) \to \lambda_{\max}(\mathbb{E}_{\mathbf{x} \sim r}[\mathbf{x}\mathbf{x}^{\top}])$ almost surely. Moreover, for any n > d, $\lambda_{\min}(\frac{1}{n}\mathbf{X}\mathbf{X}^{\top}) = 0$. As a result, $\gamma_{\text{inner},n} \to \lambda_{\max}(\mathbb{E}_{\mathbf{x} \sim r}[\mathbf{x}\mathbf{x}^{\top}])$ almost surely, which provides the desired asymptotic result.

We continue with $\gamma_{2,n}$. We first remark that $\max_{i=1,\dots,n} \|\mathbf{x}_i\|_2^2 \leq \sup_{\mathbf{x} \in \operatorname{Spt}(r)} \|\mathbf{x}\|_2^2$. As a result, by defining $\tilde{\gamma}_{2,n} = \gamma_{\operatorname{inner},n} + \max_{\mathbf{x} \in \operatorname{Spt}(r)} \|\mathbf{x}\|_2^2$, $\operatorname{GMG}_{r_n}^2$ is also $\tilde{\gamma}_{2,n}$ -weakly convex. Moreover, $\max_{\mathbf{x} \in \operatorname{Spt}(r)} \|\mathbf{x}\|_2^2$ does not depends on n, $\tilde{\gamma}_{2,n} \to \lambda_{\max}(\mathbb{E}_{\mathbf{x} \sim r}[\mathbf{x}\mathbf{x}^\top]) + \max_{\mathbf{x} \in \operatorname{Spt}(r)} \|\mathbf{x}\|_2^2$ almost surely, which also provides the desired asymptotic result.

C EXPERIMENTAL DETAILS

All our experiments build on python and the jax-framework (Babuschkin et al., 2020), alongside ott-jax for optimal transport utilities.

Method	Parameter	Values
β -VAE	β	[2, 4, 6, 8, 10, 16]
β -TCVAE	β	[2, 4, 6, 8, 10, 16]
+ HFS	γ	[1, 10]
+ DST	λ	[0.1, 1, 5, 10, 20]
+ GMG	λ	[0.1, 1, 5, 10, 20]
+ Jac	$ $ λ	[0.1, 1, 5, 10, 20]

C.1 DETAILS ON DISENTANGLEMENT BENCHMARK

To effectively conduct comprehensive and representative research on disentangled representation learning, we convert the public PyTorch framework proposed in Roth et al. (2023) to an equivalent jax variant. We verify our implementation through replications of baseline and HFS results in Roth et al. (2023), mainting relative performance orderings and close absolute disentanglement scores (as measured using DCI-D, whose implementation directly follows from Locatello et al. (2019a) and leverages gradient boosted tree implementations from scikit-learn).

For exact and fair comparison, we utilize standard hyperparamater choices from Roth et al. (2023) (which leverages hyerparameters directly from Locatello et al. (2019a), Locatello et al. (2020) and https://github.com/google-research/disentanglement_lib). Consequently, the base VAE architecture utilized across all experiment is the same as the one utilized in Roth et al. (2023) and Locatello et al. (2020): With image input sizes of $64 \times 64 \times N_c$ (with N_c the number of input image channels, usually 3). The latent dimensionality, if not otherwise specified, is set to 10. The exact VAE model architecture is as follows:

- **Encoder**: [conv(32, 4 × 4, stride 2) + ReLU] × 2, [conv(64, 4 × 4, stride 2) + ReLU] × 2, MLP(256), MLP(2 × 10)
- **Decoder**: MLP(256), [upconv(64, 4×4 , stride 2) + ReLU] \times 2, [upconv(32, 4×4 , stride 2) + ReLU], [upconv(n_c , 4×4 , stride 2) + ReLU]

Similar, we retain all training hyperparameters from (Roth et al., 2023) and (Locatello et al., 2020): Using an Adam optimizer ((Kingma and Ba, 2014), $\beta_1 = 0.9$, $\beta_2 = 0.999$, $\epsilon = 10^{-8}$) and a learning rate of 10^{-4} . Following Locatello et al. (2020); Roth et al. (2023) we utilize a batch-size of 64, for which we also ablate all baseline methods. The total number of training steps is set to 300000.

As commonly done for this setting (Locatello et al., 2019a; 2020; Roth et al., 2023), we also perform a small grid search over all the hyperparameters. We report the full details in Tab. 3.

For λ_e and λ_d , we set $\lambda_e = 0$ for the Decoder setting and $\lambda_d = 0$ for the Encoder setting while altering the weighting for the other λ . All experiments run on a single RTX 2080TI GPU.

C.2 STABILITY ANALYSIS

For the gradient stability analysis experiment, we repeat the following experiment for each of the four image datasets \mathcal{D} that we consider. We first considered a fixed neural map T_{θ} , which we choose to be a randomly initialized neural network, consisting of encoder and decoder, before any training. For $k=1,\ldots,5$, we sample a batch $\mathbf{x}_1^k,\ldots,\mathbf{x}_n^k\sim\mathcal{D}$ and let $r_n^k=\frac{1}{n}\sum_{i=1}^n\delta_{\mathbf{x}_i^k}$. We report the pariwise cosine similarity between the gradients of the DST and the GMG. Formally, we compute $\mathrm{cos\text{-}sim}(\nabla_{\theta}\mathrm{DST}_{r_n^k}(T_{\theta}),\nabla_{\theta}\mathrm{DST}_{r_n^l}(T_{\theta}))$, and $\mathrm{cos\text{-}sim}(\nabla_{\theta}\mathrm{GMG}_{r_n^k}(T_{\theta}),\nabla_{\theta}\mathrm{GMG}_{r_n^l}(T_{\theta}))$, for $k,l=1,\ldots,5$.

D ADDITIONAL EMPIRICAL RESULTS

In this section, we report additional empirical results revolving around the regularization of encoder. First in D.1, we conduct the stability analysis when regularizing the encoder. Then, we report further

results of encoder and decoder regularization on DSprites. Lastly, we take a first exploratory step towards decoder-free disentangled representation learning in D.3.

D.1 ENCODER STABILITY ANALYSIS

We repeat the following experiment for each of the four image datasets \mathcal{D} that we consider. We first considered a fixed neural map T_{θ} . For $i=1,\ldots,5$, we sample a batch $\mathbf{x}_1^i,\ldots,\mathbf{x}_n^i\sim\mathcal{D}$ and let $r_n^i=\frac{1}{n}\sum_{i=1}^n\delta_{\mathbf{x}_n^i}$. We report the pariwise cosine similarity between the gradients of the DST. Namely, for $i,j=1,\ldots,5$, we compute $\operatorname{cos-sim}(\nabla_{\theta}\mathrm{DST}_{r_n^i}(T_{\theta}),\nabla_{\theta}\mathrm{DST}_{r_n^j}(T_{\theta}))$, and the GMG, $\operatorname{cos-sim}(\nabla_{\theta}\mathrm{GMG}_{r_n^i}(T_{\theta}),\nabla_{\theta}\mathrm{GMG}_{r_n^j}(T_{\theta}))$.




Figure 4: Gradient stability analysis on the DST and GMG as Encoder regularizations. The cosine similarity is computed between all pairs of gradients ∇_{θ} obtained through 5 randomly sampled batches and a fixed network T_{θ} for each dataset.

D.2 ENCODER ANALYSIS ON DSPRITES

Table 4: Disentanglement of regularizing the Encoder and the Encoder and Decoder as measured by **DCI-D** on DSprites. We highlight **best**, <u>second best</u>, and *third best* results for each method and dataset.

DCI-D	β -VAE	β -TCVAE	β -VAE + HFS	β -TCVAE + HFS
DSprites Higgins et al. (2017)				
Base	27.6 ± 13.4	36.0 ± 5.3	38.7 ± 15.7	48.1 ± 10.8
+ Enc-DST	32.8 ± 15.0	36.5 ± 5.9	33.9 ± 15.9	48.9 ± 11.1
+ Enc-GMG	27.5 ± 14.3	37.4 ± 5.8	31.0 ± 14.3	45.9 ± 10.9
+ Dec-DST	28.6 ± 19.3	32.4 ± 8.5	39.3 ± 18.1	49.0 ± 11.2
+ Dec-GMG	39.5 ± 15.2	42.2 ± 3.6	46.7 ± 2.0	50.1 ± 8.5

D.3 TOWARDS DECODER-FREE DISENTANGLEMENT

Recently, works such as Burns et al. (2021); von Kügelgen et al. (2021); Eastwood et al. (2023); Matthes et al. (2023); Aitchison and Ganev (2024) have shown the possibility of disentanglement through self-supervised, contrastive learning objectives in an effort to align with the scalability of encoder-only representation learning (Chen et al., 2020b; Zbontar et al., 2021; Bardes et al., 2022; Garrido et al., 2023). However, these encoder-only approaches still require weak supervision or access to multiple views of an image to learn meaningful representations of the data samples.

As the goal of geometry preservation connects the data manifold and the latent domain through a minimal distortion objective and is applicable to both the encoder and decoder of a VAE (§3, Table 4), we posit that its application may provide sufficient training signal to learn meaningful representations and encourage disentanglement, eliminating the need for a reconstruction loss and decoder. Table 5 shows preliminary results on unsuper-

Table 5: Disentanglement (DCI-D) without a decoder trained with various regularizations on Shapes3D (Burgess and Kim, 2018).

Decoder-free	β-VAE	β -TCVAE	
Base	0.0 ± 0.0	0.0 ± 0.0	
$\mathbf{L2^2}: \ \cdot - \cdot\ _2^2$			
+ DST	38.2 ± 0.8	42.7 ± 1.6	
+ GMG	$\textbf{13.9} \pm 0.4$	20.5 ± 0.5	
$\mathbf{ScL2^2}: \alpha \ \cdot -\cdot \ _2^2, \alpha > 0$ learnable			
+ DST	45.6 ±1.2	53.5 ±1.0	
+ GMG	15.2 ± 0.3	25.2 ± 0.6	
$\mathbf{Cos}: cos\text{-}sim(\cdot,\cdot)$			
+ DST	37.0 ± 0.4	46.1 ± 1.5	
+ GMG	37.0 ± 0.9	38.8 ± 1.1	

vised decoder-free disentangled representation learning on the Shapes3D benchmark, where the decoder and associated reconstruction objective have been removed.

Standard approaches such as β -VAE or β -TCVAE collapse and do not achieve measurable disentanglement (DCI-D of 0.0). However, the inclusion of either DST or GMG significantly raises achievable disentanglement and, combined with the β -TCVAE matching objective, can achieve DCI-D scores of up to 53.5 without needing any decoder or reconstruction loss. While these are preliminary insights, we believe they offer promise for more scalable approaches to unsupervised disentangled representation learning and potential bridges to popular and scalable self-supervised representation learning approaches. Note, that here the distortion loss significantly outperforms the GMG. This is expected due to the nature of the GMG, as the distortion loss offers a more restrictive and, thus, stronger signal for learning representations, which is necessary in the absence of a reconstruction objective. This highlights that while in most scenarios (§ 2.1, Figure 2), the GMG is preferable over the distortion loss, there also exist settings where a more restrictive optimization signal is desirable.

E PYTHON CODE FOR THE COMPUTATION OF THE GROMOV-MONGE GAP

```
import jax
import jax.numpy as jnp
from ott.geometry import costs, geometry
from ott.solvers.quadratic import gromov_wasserstein
from ott.problems.quadratic import quadratic_problem
def gromov_monge_gap_from_samples(
   source: jax.Array,
   target: jax.Array,
   cost_fn: costs.CostFn = costs.Cosine(),
   epsilon: float = 0.1,
   stat_fn: Callable, # usually computes the mean of the cost matrix
    **kwargs,
) -> float:
    """Gromov Monge gap regularizer on samples."""
    # define source and target geometries
   cost_matrix_x = cost_fn.all_pairs(x=source, y=source)
   scale_cost_x = stat_fn(scale_cost_x)
   cost_matrix_x = cost_matrix_x / jax.lax.stop_gradient(scale_cost_x)
   geom_xx = geometry.Geometry(cost_matrix=cost_matrix_x)
   cost_matrix_y = cost_fn.all_pairs(x=target, y=target)
   scale_cost_y = stat_fn(cost_matrix_y)
   cost_matrix_y = cost_matrix_y / jax.lax.stop_gradient(scale_cost_y)
   geom_yy = geometry.Geometry(cost_matrix=cost_matrix_y)
    # define and solve entropic GW problem
   prob = quadratic_problem.QuadraticProblem(geom_xx, geom_yy)
   solver = gromov_wasserstein.GromovWasserstein(
       epsilon=epsilon, **kwargs
   out = solver(prob)
    # compute the distortion induced by the map
   distortion_cost = jnp.nanmean(
        (geom_xx.cost_matrix - geom_yy.cost_matrix) **2
    # compute optimal (entropic) gromov-monge displacement
   reg_gw_cost = out.reg_gw_cost
   ent_req_gw_cost = req_gw_cost - 2 * epsilon * jnp.log(len(source))
    # compute gromov-monge gap
   loss = distortion_cost - ent_reg_gw_cost
   return loss * jax.lax.stop_gradient(scale_cost_x * scale_cost_y)
```