

Molecular Dynamics and Machine Learning Unlock Possibilities in Beauty Design—A Perspective

Yuzhi Xu,^{1,2,3} Haowei Ni,¹ Qinhui Gao,⁴ Ziqing Zhao,¹ Chia-Hua Chang,¹ Yanran Huo,¹ Fanyu Zhao,³ Shiyu Hu,³ Wei Xia,³ Yike Zhang,⁵ Radu Grovu,⁶ Min He,⁷ John Z. H. Zhang,^{3,2} and Yuanqing Wang^{1,2,8, a)}

¹⁾Simons Center for Computational Physical Chemistry, New York, N.Y. 10003

²⁾Department of Chemistry, New York University, New York, N.Y. 10003

³⁾Shanghai Frontiers Science Center of Artificial Intelligence and Deep Learning and NYU-ECNU Center for Computational Chemistry, NYU Shanghai, Shanghai 200062, P.R.China

⁴⁾Department of Digital Humanities, King's College London, Strand, London WC2R 2LS, U.K.

⁵⁾Division of Cardiology, The First Affiliated Hospital of Nanjing Medical University, Nanjing 210029, P.R.China

⁶⁾Internal Medicine Department, Rhode Island Hospital, Brown University Health, Providence, R.I., 02903

⁷⁾Xbiome, Inc., Cambridge, Mass. 01451

⁸⁾Center for Data Science, New York University, New York, N.Y. 10004

Computational molecular design—the endeavor to design molecules, with various missions, aided by machine learning and molecular dynamics approaches, has been widely applied to create valuable new molecular entities, from small molecule therapeutics to protein biologics. In the small data regime, physics-based approaches model the interaction between the molecule being designed and proteins of key physiological functions, providing structural insights into the mechanism. When abundant data has been collected, a quantitative structure-activity relationship (QSAR) can be more directly constructed from experimental data, from which machine learning can distill key insights to guide the design of the next round of experiment design. Machine learning methodologies can also facilitate physical modeling, from improving the accuracy of force fields and extending them to unseen chemical spaces, to more directly enhancing the sampling on the conformational spaces. We argue that these techniques are mature enough to be applied to not just extend the *longevity* of life, but the *beauty* it manifests. In this perspective, we review the current frontiers in the research & development of skin care products, as well as the statistical and physical toolbox applicable to addressing the challenges in this industry. Feasible interdisciplinary research projects are proposed to harness the power of machine learning tools to design innovative, effective, and inexpensive skin care products.

I. Introduction: Beauty design and computer-aided molecular discovery.

The human race has been using cosmetics and skin-care products for more than 8000 years¹—the history of which arguably predates that of medical practice. Through these 8 millenniums, while medical and medicinal sciences have been transformed into quantitative, mechanistic, rigorous, and digitalized disciplines, the design of cosmetics and skin-care products, referred to as *beauty design* henceforth, remains largely empirical, experimental, and trial-and-error-based.

The *in silico* approaches to design new molecule entities, namely therapeutics, using computer simulation and modeling are collectively termed *computer-aided* molecular discovery. Roughly, they can be characterize into two categories, *physics-inspired* and *data-driven*, which correspond to *ligand-based* and *structure-based* drug discovery in medicinal chemistry, respectively. The former models the *conformational* landscape of a biomolecular system \mathcal{G} , which adopts a Boltzmann-type distribution:

$$p(\mathbf{x}|\mathcal{G}) \propto \exp(-u(\mathbf{x};\mathcal{G})), \quad (1)$$

based on which ensemble properties of the system can be estimated using the samples drawn from this distribution \mathbf{x}_i :

$$\langle \mathcal{O}(\mathbf{x}) \rangle = \int d\mathbf{x} p(\mathbf{x}|\mathcal{G}) \mathcal{O}(\mathbf{x}) \approx \sum_{\mathbf{x}_i} \mathcal{O}(\mathbf{x}_i). \quad (2)$$

And molecular dynamics (MD) simulations can be employed to generate such samples. If one only wishes to draw insights from the *minima* of Equation 1, they can further simplify Equation 2 to be:

$$\hat{\mathcal{O}}(\mathbf{x}) \approx \mathcal{O}(\operatorname{argmax}_{\mathbf{x}} p(\mathbf{x};\mathcal{G})), \quad (3)$$

where the optima are found by structural methods such as docking² or protein folded pose prediction^{3,4}, although this can be a very crude approximation neglecting the dynamic nature of biomolecular systems.

On the other hand, *data-driven* methods seek insights directly from data, most typically the molecule-measurement pairs. Quantitative structure-activity relationship (QSAR) models are constructed to model the mapping $f: \mathcal{G} \rightarrow \mathbb{R}$ from the molecular topological graphs \mathcal{G} to scalar physical, chemical, or physiological properties. This is where machine learning has become an indispensable workhorse for designing new molecular entities⁵⁻⁷. In principle, this school of molecular modeling characterizes the *posterior predictive distribution*

$$p(\hat{y}_{\mathcal{G}}|\mathcal{G}, \mathcal{M}) = \int d\Theta p(\hat{y}_{\mathcal{G}}|\mathcal{G}, \Theta, \mathcal{M}) p(\Theta|\mathcal{M}, \mathcal{D}), \quad (4)$$

where the property of a molecule \mathcal{G} given a model \mathcal{M} can be written as that dependent upon the parameters to the model Θ , marginalized across the parameter distribution conditioned on the existing data $\mathcal{D} = \{\mathcal{G}_i, y_i\}$. This principled Bayesian approach is usually replaced with a single point maximum-likelihood estimate (MLE) on the parameter space.

While traditional fingerprint⁸-based methods are still used, small molecules are now more ubiquitously modeled as

^{a)}Correspondence email address: wangyq@wangyq.net

graphs, and processed through a graph neural network (GNN)^{9–14} to form useful representations for downstream prediction tasks. Once the backbone of this mapping is constructed, it can also be used to quantify the uncertainty of its predictions in a Bayesian setting^{15,16}, for example with variational inference¹⁷, or in an active learning setting to guide the design of the next round of experiments¹⁸.

Additionally, machine learning techniques are also applied in a *structure-based* molecular discovery setting, where the accuracy or efficiency of traditional physical models are revolutionized by machine learning methodologies. On the one hand, machine learning provides new functional forms^{4,19–24} and parametrization methods^{25–29} for force fields to be used in molecular dynamics (MD) simulations. On the other hand, it has also been applied to directly predict the structure or conformational distributions³⁰. When it comes to proteins, in particular, the determination of the folded conformation thereof has been dramatically accelerated using machine learning-based models^{3,4,31}, and with the predicted structures one can readily *design*³² new proteins to fulfill desired properties or functions.

In this perspective, we first review the overarching goals and specific challenges in beauty design. Next, we identify the physical (MD-based) and machine learning toolboxes useful in modeling and optimizing the properties of interest in beauty design. Finally, we also envision beauty design areas where MD simulation and ML modeling can be of immediate utility, and propose feasible projects.

II. Objectives in skincare and cosmetics

In this section, we review the primary goals in beauty design, the general taxonomy, and the physical laws to be considered in each axis.

A. Coloration.

Cosmetic colorants can be classified into two major categories based on their chemical composition: inorganic and organic colorants³³. The global annual production of inorganic colorants is approximately 10 million tons, accounting for 80% of the total colorant production³⁴. Key inorganic colorants include titanium dioxide, iron oxides, carbon black, zinc oxide, ultramarine pigments, and mica-based pearlescent pigments, most of which are derived from natural minerals^{35,36}—these pigments are widely used in facial, eye, lip, and nail products. Due to their stable crystalline structure, inorganic pigments exhibit high stability, excellent opacity from their high refractive index, and natural coloration³⁷. Nevertheless, they also have certain limitations, such as a narrow range of light absorption, making it difficult to cover a broader visible spectrum, high surface energy leading to aggregation, and large particle size affecting skin permeability³⁸.

As to organic colorants, they can be further divided into natural and synthetic colorants³⁹. Natural colorants are typically derived from plant and insect pigments, such as carmine,

carotenoids, anthraquinones, betalains, and flavonoids. These colorants are popular among consumers due to their natural origin⁴⁰, although they also face challenges such as color instability and susceptibility to environmental degradation. In contrast, synthetic colorants generally offer excellent color stability, a wide range of spectral options, and superior light and heat resistance⁴¹. This is mainly due to their molecular structures, which often contain a large number of conjugated double bonds (e.g., azo and bonds, triphenylmethane structures). These conjugated systems extend electron delocalization, enabling absorption over a broader wavelength range of visible light and producing vivid colors³⁴. The highly conjugated structures not only enhance molecular stability and reduce structural breakdown under light or heat exposure but also enable optimal electronic transitions, effectively absorbing specific wavelengths of light to improve color vibrancy and stability⁴². Synthetic colorants encompass a wide variety of types, including azo dyes, triphenylmethane dyes, flavonoid dyes, and anthraquinone dyes. Through optimized molecular design, these different types of dyes ensure color performance and stability across various application environments. Moreover, the consistency achieved through batch synthesis results in highly uniform molecular structures and purity, further enhancing the clarity and stability of the colors. However, due to environmental and health concerns, the design of synthetic colorants requires more stringent molecular design criteria to minimize potential risks⁴³.

B. Moisturization.

The moisturizing efficacy of skincare products is typically assessed through Transepidermal Water Loss (TEWL), which measures the rate at which water diffuses from the stratum corneum into the atmosphere⁴⁴, where an evaporimeter is used to quantify the amount of water lost per square meter of skin surface per hour, providing insights into the skin’s water kinetics. Since TEWL primarily reflects the process of water diffusing from the deeper layers of the skin through the epidermis to the external environment, this process can be approximately described by Fick’s First Law of Diffusion⁴⁵:

$$J = -D \frac{dC}{dx} \quad (5)$$

where J is the water flux D is the diffusion coefficient of water in the stratum corneum and dC/dx is the concentration gradient of water.

Based upon this relationship, different types of moisturizing products regulate TEWL by adjusting the diffusion coefficient D or the concentration gradient dC/dx , thus achieving a moisturizing effect. Typically, moisturizing ingredients in skincare products fall into one or more of the following three categories: humectants, occlusives, and emollients³⁶. Humectants are generally compounds containing multiple hydroxyl groups ($-OH$), such as glycerin, hyaluronic acid, α Hydroxy Acids, and propylene glycol. These molecules form stable hydrogen-bond networks with water, increasing the water content in the stratum corneum and lowering the chemical




FIG. 1. Semantic overview.

potential of water. By increasing the water concentration C within the stratum corneum, humectants reduce the concentration gradient dC/dx . According to Fick's first law, when humectants increase the water concentration in the stratum corneum, the concentration gradient dC/dx decreases, leading to a reduction in the water flux J and consequently reducing the rate of water evaporation. Occlusives, such as petroleum jelly, mineral oil, and silicone oils, are typically hydrophobic compounds that form a physical barrier on the skin surface⁴⁶. This barrier prevents water evaporation and thereby reduces the rate of water flux J from the skin, effectively minimizing transepidermal water loss. Emollients work by softening and smoothing the skin, filling the gaps between skin cells, many of which possess both occlusive and emollient properties.

C. Skin Lightening

To achieve skin lightening typically requires intervention in the processes of melanin production, transport, and metabolism in the skin⁴⁷. Inhibiting melanin production is primarily achieved by targeting tyrosinase, a copper-containing metalloenzyme with two copper ions (CuA and CuB) in its active site, which are responsible for substrate binding and catalysis^{48,49}. There are several strategies for inhibiting tyrosinase activity. The most direct approach involves inhibiting its active site. For example, arbutin effectively reduces tyrosinase activity through competitive inhibition, thereby decreasing melanin production⁵⁰. Another approach is to chelate the copper ions (CuA and CuB) at the active site, preventing the enzyme from functioning properly. Kojic acid, for instance, can form chelates with the copper ions in tyrosinase, significantly inhibiting its catalytic function⁴⁹. Additionally, retinoic acid can suppress TYR gene promoter activity, reducing tyrosinase transcription and consequently lowering melanin synthesis at the source⁵¹. Tyrosinase activity is also influenced by glycosylation modifica-

tions, and inhibitors like mannose can reduce tyrosinase glycosylation, thereby decreasing its enzymatic activity⁵². Moreover, linoleic acid and β -sitosterol promote tyrosinase ubiquitination and proteasomal degradation, reducing the amount of active enzyme in the cell, and further inhibiting melanin production^{53,54}.

Blocking melanin transport is also an essential strategy for achieving skin whitening. By interfering with the transfer of melanosomes from melanocytes to surrounding keratinocytes, the distribution of melanin in the epidermis can be reduced. Niacinamide, for example, can inhibit protease-activated receptor-2 (PAR-2) on keratinocytes, thereby reducing melanosome transfer⁴⁷. Similarly, licorice extract interferes with the melanosome transport process by inhibiting signaling pathways such as PI3K/Akt and MAPK, further reducing melanin deposition in the epidermis⁵⁵.

D. Skin Penetration.

An important function in cosmetics is to activate collagen production and enhance antioxidant effects at the deeper layers of the skin^{56,57}. The skin serves as a protective barrier against external environmental factors. Therefore, for cosmetic molecules to be effective, they must penetrate this barrier to reach the underlying layers. Common skin-penetrating ingredients in cosmetic products can be broadly categorized into small molecules, peptides and liposomes.

In particular, small molecules in cosmetic products, such as Vitamin C, Vitamin A, Vitamin E, salicylic acid, and niacinamide, play crucial roles in processes such as amino acid hydroxylation in collagen, activation of signaling pathways, and antioxidation⁵⁸⁻⁶³. Due to their molecular weights ranging from 122 to 430, these small molecules can penetrate the deeper layers of the skin through aqueous channels or the lipid bilayer. Despite their favorable permeability, the rate of penetration is influenced by several factors, including formula-

tion, concentration, temperature, and pH. For example, Vitamin C requires a water-based formulation with a pH lower than its pKa (pKa 4.2) to reduce the charge density on the molecule⁶⁴. Similarly, the skin permeability of salicylates exhibits a parabolic relationship with the partition coefficient, with optimal permeability typically occurring at a log P value of around 2.5. At this pH, salicylates have sufficient solubility in the lipid layer to facilitate diffusion through the stratum corneum while maintaining enough hydrophilicity to penetrate the viable epidermal tissue⁶⁵. To enhance the skin penetration of these small molecules while preserving their efficacy, various optimization strategies can be employed. These include molecular group modification, chemical potential optimization, eutectic systems, and coordination compounds.

In addition to their direct effects on the skin, small molecules also play a role in facilitating the penetration of peptides and proteins into the body. Peptides and larger protein molecules are particularly effective in wrinkle reduction, skin vitality maintenance, and collagen repair, as well as in sensitive factor attachment protein receptor (SNARE) regulation^{66,67}. However, due to their larger molecular weight and hydrophilic nature, these molecules face significant challenges in penetrating the hydrophobic lipid matrix of the stratum corneum⁶⁸. To address these permeability issues, various strategies are being developed, such as the use of small-molecule chemical penetration enhancers (CPEs)⁶⁹. These include azone derivatives, fatty acids, alcohols, esters, sulfonides, pyrrolidones, glycols, surfactants, and terpenes^{70,71}. These chemicals can enhance the permeability of peptides and proteins by increasing the diffusion rate within the skin, inducing lipid fluidization, optimizing thermodynamic activity, influencing partition coefficients, and enhancing the release of drugs from formulations into the upper layers of the skin. However, while CPEs significantly improve permeability, they may also cause skin irritation and inflammation, and so far, no ideal chemical penetration enhancer has been identified⁷².

In the way of searching for more effective, non-invasive, non-toxic, and non-irritating skin penetration enhancers, short-sequence peptides have been identified as safer alternatives to small molecule CPEs^{73,74}. By designing and synthesizing specific short-chain peptides, it is possible to facilitate the delivery of other substances to the deeper layers of the skin. These short peptides have small molecular weights, making them more likely to penetrate the skin. Among them, cell-penetrating peptides (CPPs) and skin-penetrating peptides (SPPs) are recognized as valuable delivery tools due to their high transduction efficiency and low cytotoxicity⁷⁵⁻⁷⁸. A notable example is the TAT peptide (Trans-Activator of Transcription), a short peptide sequence (amino acids 47-57) that has been extensively studied and applied in cell penetration and drug delivery. When combined with the tripeptide GKH, the resulting TAT-GKH complex significantly enhances GKH's skin penetration efficiency and is utilized in lipolytic cosmetic products⁷⁹. Additionally, Chen et al. reported the use of the skin-penetrating peptide SPACE to enhance the topical delivery of high molecular weight hyaluronic acid (HA, molecular weight: 200-325 kDa)⁸⁰. Carmichael and

colleagues demonstrated that botulinum toxin type A (BoNT-A) could be administered subcutaneously or topically with a novel transdermal delivery peptide to reduce inflammation by inhibiting the activation of nociceptors in the skin⁸¹. Similar to small molecules, modifying the side chains of these peptides can also enhance their permeability. For example, the carrier peptide GHK can bind to Cu(II) and Mn ions in the skin⁸²⁻⁸⁴. Through chemical modification, GHK can be linked to a palmitoyl group (a long-chain fatty acid), increasing its hydrophobicity and enhancing its interaction with the lipid bilayer, thereby improving its ability to penetrate the skin^{85,86}.

Besides, liposomes are utilized to encapsulate peptides and proteins, enhancing their ability to penetrate the skin barrier more effectively^{87,88}. Liposomes offer several advantages for the transdermal delivery of peptides and proteins, including improved permeability, protection of active ingredients, and controlled release. With the advent of novel liposome technologies, such as elastic liposomes, cationic liposomes, and pH-sensitive liposomes, delivery efficiency has been further enhanced^{89,90}.

III. Frontiers in molecular discovery for beauty design

In §II, we have outlined the general guidelines of beauty design. In this section, we focus on two directions particularly popular in beauty design—collagen formation and microhyaluronic acid application.

A. The odyssey towards collagen

Collagen, the most common type of protein found in skin as well as tendons and bones, assists fibroblasts forming in the dermis, promoting the growth of new cells and replacing dead skin cells. Oral supplement of collagen has been linked to improved firmness, suppleness, moisture, smoothness, and elasticity of skin⁹¹⁻⁹³, although larger-scale study is needed to determine whether commercially available products are useful for long-term use. Used topically, certain mixtures of amino acids, such as arginine and ornithine, have been proven to promote the disposition of collagen in wounds⁹⁴⁻⁹⁶. Topically, the application of amino acids have also shown to be useful to boost collagen formation and skin moisture⁹⁷.

B. Micro hyaluronic acids: potency and risks

Micro hyaluronic acids with low molecular weight are well known for their ability to penetrate the skin effectively. These tiny molecules can go through the skins' protective barrier and provide moisture directly to the deeper layers of the skin improving its elasticity and minimizing noticeable aging signs. This property is crucial in cosmetic dermatology, where hydration and skin suppleness are key to a youthful appearance. The effectiveness of micro hyaluronic acids in promoting skin hydration and reducing wrinkles has been substantiated

by various studies, which highlight its efficacy as a humectant, with the ability to hold water up to a thousand times their weight and facilitate better absorption compared to their higher molecular counterparts^{98,99}. Advancements in skincare technology have leveraged the properties of micro hyaluronic acids to develop sophisticated delivery systems that maximize skin benefits. Techniques such as encapsulation in liposomes or binding to nanoparticles have been explored to further enhance the penetration and longevity of hyaluronic acids in skin layers. These innovations not only improve the immediate hydrating effects but also contribute to the long-term anti-aging benefits by maintaining higher levels of skin hydration over extended periods⁹⁹.

Despite their benefits, micro hyaluronic acids come with potential risks that must be managed carefully. Topical applications are generally safe but can occasionally cause skin irritation or mild allergic reactions in sensitive individuals. More invasive applications, such as injections, carry greater risks including infection, tissue necrosis, or adverse reactions at the injection site. It is vital that these treatments are administered by qualified healthcare professionals to minimize risks and ensure safety. Additionally, Bravo *et al.*⁹⁸ conducted a comprehensive review and noted that extremely low molecular weights of hyaluronic acid may trigger inflammatory responses in the skin. It is necessary to have precise control over the molecular size used in skin care formulations.

Because of the growing usage of hyaluronic acid in the field of aesthetics, notably in injectable skin rejuvenation treatments, it has become essential to build a solid regulatory framework in order to protect the efficacy and safety of this substance. Regulatory scrutiny ensures that hyaluronic acid products, classified as medical devices, meet stringent safety standards and are marketed accurately regarding their benefits and potential side effects. The framework also highlights the importance of the administration by licensed practitioners and the need for ongoing research to support the claims made by manufacturers regarding the cellular effects of these injections¹⁰⁰. Hyaluronic acid usage regulations are made to protect the general public's health as well as satisfy the increasing market for anti-aging goods. This involves guaranteeing the chemicals' moral provenance and scientific backing. Strict regulatory measures are required as demand for these therapies rises in order to guard against abuse and guarantee that the goods live up to expectations without sacrificing safety.

IV. *In silico* beauty design: recent advances.

a. Efforts using machine learning to design peptides for skin care. Current trends in cosmetic molecule development closely align with those in medicinal chemistry^{101,102}. In recent years, machine learning has played a significant role in peptide design and optimization, greatly improving the development efficiency of cosmetic peptides. Specifically, Yue *et al.*¹⁰³ reported using machine learning to predict the antimicrobial activity and hemolytic properties of peptides, leading to the design of a novel β -defensin-based antimicrobial peptide, IK-16-1. Additionally, Hsueh *et al.* design a mul-

tifunctional peptides which can efficiently penetrate the cells and combined with melanin¹⁰⁴. In the area of skin penetration, De Oliveira *et al.* reported the development of BChemRF-CPPred, which extracts descriptors from peptide structures and sequences to distinguish cell-penetrating peptides (CPPs) from non-CPPs¹⁰⁵. The design of short peptides has become a prominent strategy in molecular discovery, primarily due to their strong targeting capabilities, high efficacy, stability, and low allergenicity¹⁰⁶. Although substantial progress has been made utilizing scientists' domain knowledge, the *de novo* design of short peptides remains relatively limited^{107,108}.

b. Natural molecular discovery for skin care. Natural molecular discovery is emerging as a new trend in molecular design. This shift is largely driven by the perception that natural molecules are inherently safer and exhibit lower toxicity, aligning with consumer preferences and regulatory standards for safer cosmetic and therapeutic products. Consequently, most animal and plant extracts are widely recognized as safe, leading to the screening of many cosmetic molecules using chemical libraries derived from natural sources¹⁰⁹. For example, Bournez *et al.* employed homology modeling and molecular docking to screen 4,534 natural products for potential SIK2 inhibitors, aiming to enhance melanin production and achieve tanning effects without UV-induced damage¹¹⁰. Besides, Su *et al.* compiled a database of 5,616 natural compounds and, through XP docking and molecular dynamics simulations, identified five compounds with promising dynamic binding properties. Subsequent *in vivo* experiments on zebrafish revealed that two components, Sennoside B (SB) and Sennoside C (SC), exhibited inhibitory effects on melanin production¹¹¹.

c. Structural modification for cosmetic molecules. Structure modification is a primary method for developing new cosmetic molecules¹¹². Typically, chemical structures are altered to enhance their biological activity, stability, and Absorption, Distribution, Metabolism, Excretion, and Toxicity (ADMET) properties. In addition to these direct approaches, derivatives can also be generated based on known targets and established cosmetic molecules. Specifically, a typical example is the molecular modification of vitamins, to address the inherent instability and challenges with the topical application of vitamin C, formulation chemists have engineered derivatives such as ascorbyl 2-phosphate (SAP and MAP) and ascorbyl 6-palmitate (AA-Pal)⁶⁴. These derivatives stabilize vitamin C within cosmetic formulations and enhance skin penetration by maintaining stability at a neutral pH and protecting the molecule from oxidation, facilitating their inclusion in various skincare products. Similarly, many researchers have modified natural products to enhance their bioactivity. Kumari *et al.* demonstrated that isoliquiritigenin and its derivatives, such as 5'-formylisoliquiritigenin and isoliquiritigenin dimers, exhibit higher binding affinity and more potent tyrosinase inhibition compared to isoliquiritigenin alone¹¹³. Molecular docking studies revealed that these derivatives have lower binding energies with tyrosinase, indicating their potential as more effective tyrosinase inhibitors for the prevention and treatment of melanogenesis and related hyperpigmentation disorders.

In addition to these methods, cosmetic formulations following molecular editing are further evaluated through their physical properties. Our article focuses on the use of chemical computational tools for prediction and combination. Therefore, in the subsequent Physical Toolbox section, we briefly introduce the relevant prediction tools, which will not be reiterated here.

V. The physical toolbox: molecular dynamics (MD) simulation.

Characterizing Equation 1 by drawing samples while integrating Newton’s second motion of law, MD simulation finds use in all schools of molecular modeling. There have been a wide variety of mature and wide used pipelines for modeling the energy landscape of biomolecular systems of therapeutical interest, which can be seamlessly adopted to model those of cosmeceutical significance.

a. Modeling of cosmeceutical molecule-target binding. For beauty design, MD simulations^{23,114–116} can be employed to model the interactions between the cosmeceutical molecules and their targets as well as provide detailed insights into the hydrogen bonding interactions that are critical for binding specificity and stability^{117,118}. Quantitatively, the binding free energy between the small molecule and the target protein can be reliably predicted¹¹⁹. For crude predictions, popular methods include MM-GBSA (Molecular Mechanics Generalized Born Surface Area) and MM-PBSA (Molecular Mechanics Poisson-Boltzmann Surface Area)^{120,121}. Alanine scanning methods^{122,123} can be used to further deduce the *contribution* of each residue towards the binding free energy. More elaborately, free energy calculations based on thermodynamic integration (TI)¹²⁴ and free energy perturbation (FEP)^{125–127} have shown best correlation with experimental data, and therefore can be used to reliably predict the binding free energy between the small cosmeceutical molecule and the target.

b. Physical property prediction Usually, Cosmetic compounds necessitate a thorough evaluation of ADMET profiles to guarantee their suitability for topical application¹⁰⁹. This comprehensive assessment ensures that active ingredients are effectively absorbed through the skin barrier, distributed appropriately within the skin layers, metabolized without generating harmful by-products, and excreted without accumulation that could lead to adverse effects. Traditionally, ADMET evaluation has relied primarily on experimental methods to determine these key pharmacokinetic and toxicological properties of drugs or active molecules. Recently, the application of machine learning (ML) and deep learning (DL) techniques in ADMET prediction has emerged as a complementary approach to experimental methods, offering advantages in terms of speed and accuracy^{128–130}. Besides, Physical property prediction is required in the product’s fluid properties when applied to the skin. These properties include viscosity, rheological behavior, surface tension, diffusion coefficients, and absorption kinetics^{131,132}. These factors collectively determine a cosmetic product’s ease of application,

spreadability, and stability. Advanced analytical techniques such as atomic force microscopy and nuclear magnetic resonance spectroscopy provide molecular-level structural and dynamic information, further aiding in the understanding of how molecular interactions influence macroscopic physical properties. Employing predictive methods for the design of cosmetic products offers a cheap method to predict the cosmetic molecule’s behavior. Computational method such as computational fluid dynamics, molecular simulations, machine learning enable the forecasting of a product’s behavior under various conditions, facilitating the optimization of formulations before physical prototypes are developed.

c. Spectroscopy prediction During the product development stage, spectroscopic techniques such as near-infrared (NIR), Raman, and UV-visible spectroscopy are employed to characterize raw materials and analyze formulations¹³³. Spectroscopic prediction serves as a valuable alternative to costly experimental procedures in the cosmetics industry, as it allows the prediction of product performance and properties using existing spectral data and mathematical models, either before or during experimentation. By analyzing the spectral characteristics of raw materials, spectroscopic prediction can anticipate their behavior and efficacy within a formulation. Drawing on physical theories regarding how molecular structures influence macroscopic properties, researchers can logically deduce the interactions between different components, thereby rapidly identifying the most suitable ingredients¹³⁴. For instance, by predicting the spectral properties of blended components and leveraging the physical characteristics of light absorption, reflection, and scattering, it is possible to optimize the formulations of products such as lipsticks and foundations, which require precise color differentiation¹³⁵. Moreover, spectroscopic prediction can be utilized to forecast product stability¹³⁶. For example, in sunscreen products, spectroscopic prediction models can quickly estimate SPF values by leveraging the physical relationship between UV absorption and sunscreen efficacy, thereby reducing the need for extensive in vitro or in vivo testing¹³⁷.

VI. The machine learning toolbox

a. Graph-based machine learning. Graph neural networks^{9–14} have become the *de facto* work horse of all schools of molecular modeling. A graph neural network can be most generally defined as one adopting a layer-wise updating scheme that aggregates representations from a node’s neighborhood and updates its embedding:

$$\mathbf{X}'_v = \phi(\mathbf{X}_v, \rho(\mathbf{X}_u, \hat{A}_{uv}, u \in \mathcal{N}(v))), \quad (6)$$

where ϕ, ρ are the *update* and *aggregate* function, respectively. Omitting the nonlinear transformation step ϕ ubiquitous in all neural models, and assuming a convolutional *aggregate* function, $\rho = \text{SUM}$ or $\rho = \text{MEAN}$, a graph neural network layer is characterized by the aggregation/convolution operation that pools representations from neighboring nodes, forming an intermediary representation \mathbf{X}' , which on a global

level, with activation function σ and weights W , can be written as: $\mathbf{X}' = \sigma(\hat{A}\mathbf{X}W)$. The primary difference among architectures amounts to distinct effective adjacency matrix \hat{A} . The most classical examples include: graph convolutional networks⁹ (GCN), which normalizes \hat{A} by the in-degree of nodes $D_{ii} = \sum_j A_{ij}$ and graph attention networks² (GAT), which takes \hat{A} as the attention score;

$$\hat{A}_{\text{GCN}} = D^{-\frac{1}{2}}AD^{-\frac{1}{2}}; \hat{A}_{\text{GAT},ij} = \text{Softmax}(\sigma(\text{NN}(\mathbf{x}_i|\mathbf{x}_j))); \quad (7)$$

The resulting representation is rich and naturally permutation-invariant, providing a strong starting point for downstream tasks.

b. Generative modeling. Once we have a reliable model for the prediction of activities given the structure of the graph, we can couple it with a generative model for the true *de novo* design of molecules and peptides. This can be achieved through a principled, probabilistic manner, where the conditional distribution in Equation 4 is replaced by the distribution of the chemical graphs without measurements:

$$p(\mathcal{G}|\mathcal{M}) = \int d\Theta p(\mathcal{G}, \Theta|\mathcal{M})p(\Theta|\mathcal{M}), \quad (8)$$

and the model is typically trained by maximizing the (lower bound of) the likelihood of data $p(\mathcal{G}|\mathcal{M})$. For the generation of molecular structures, common approaches include variational autoencoder (VAE)-based models, where molecules are generated as strings¹³⁸, graphs¹³⁹, or a Markov-decision process to construct from fragments¹⁴⁰; flow-based models¹⁴¹; and diffusion-based models¹⁴². The conformations of molecules can also be simultaneously or conditionally generated in the 3-dimensional space¹⁴³.

c. Active learning. With some over-simplification, molecular discovery can be abstracted as a sequential active learning process. Given a finite pool of candidate chemical compounds, represented by chemical graphs $\mathcal{F} = \{\mathcal{G}_i, i = 1, 2, \dots, N\}$, each associated with a *potency* $y(\mathcal{G}_i) \in \mathbb{R}$, of which a noisy observation is expensive to acquire, we are interested in finding the compound with the highest potency $\mathcal{G}^* = \text{argmax}_i y(\mathcal{G}_i)$ with fewest function evaluations. This can be achieved via *active learning*, starting with an empty portfolio $\mathcal{P} = \emptyset$, and in each round, choosing a compound from the candidate pool, \mathcal{F} , based on a decision informed by the current portfolio, $\hat{\mathcal{G}} = D(\mathcal{F}|\mathcal{P}), \hat{\mathcal{G}} \in \mathcal{F}$. Subsequently, we subtract this compound from the candidate pool to add it to the portfolio, $\mathcal{F} \leftarrow \mathcal{F} \setminus \{\hat{\mathcal{G}}\}, \mathcal{P} \leftarrow \mathcal{P} \cup \{\hat{\mathcal{G}}\}$.

Bayesian optimization (BO) is a powerful approach in active learning where the decision $D(\mathcal{F}|\mathcal{P})$ is constructed in a modular way. Firstly, we extract fixed D -dimensional features from the compounds to form a *representation*: $H = h(\mathcal{G}) \in \mathbb{R}^D$; secondly, this representation is used to construct a *predictive posterior distribution* (Equation 4) with *uncertainty quantification*: $p(y|\mathcal{G};\Theta), \Theta = \theta(H)$, whose parameters Θ is mapped from the graph representations. Lastly, an *acquisition* function, α , is applied on this predictive posterior to form a score, on the basis of which a decision is made: $\hat{\mathcal{G}} = D(\mathcal{F}|\mathcal{P}) = \text{argmax}_{\mathcal{G} \in \mathcal{F}} \alpha(p(y|\mathcal{G}))$. This process can be executed iteratively until a satisfactory compound has been identified.

d. Large-language models. Large language models (LLMs)—neural networks composed primarily of transformer¹⁴⁴—have shown surprising promise^{145–149} in representing and generating text-structured data and gained popularity very quickly in recent years. There have been a plethora of LLMs for chemical tasks¹⁵⁰, the most popular ones include Yu *et al.*¹⁵¹, Zhang *et al.*¹⁵², which have been fine-tuned on instruction datasets¹⁵³. LLMs can be integrated into various workflows discussed here, from planning experiments to be carried out in an active learning setting, to writing code to prepare biomolecular systems for simulations.

VII. Roadmap: Potential use cases

Having reviewed the goals and challenges in the beauty design industry and the powerful methodologies in our physical and statistical toolboxes, in this section, we propose three projects that are immediately feasible, to showcase the potential power of combining MD and ML approaches in beauty design.

a. The molecular modification of arbutin via multiobjective active learning. Inhibiting the activity of tyrosinase, an essential enzyme for melanin synthesis in skin, arbutin has been ubiquitously used as skin-lightening agents targeting the hyperpigmentation issues. Usually extracted from the leaves of plants, such as the bearberry, the efficacy of this natural product is bottlenecked by its inability to penetrate skin to reach the melanocytes¹⁵⁴. We propose to develop arbutin derivatives displaying similar efficacy while showing significantly greater lipophilicity for better skin penetration.

First, docking and molecular dynamics simulations will be carried out to better understand the binding mechanism of arbutin. With these structural insights, we will carry out molecular modification, for example using fragment-based tools such as Zhang *et al.*¹⁵⁵, to replace functional groups on arbutin. This can be done in an ultra-high throughput manner, generating a large library of arbutin-derivatives. Multitask Bayesian optimization algorithms can then design experiments to be carried out sequentially, to optimize for both lipophilicity and melanocytes-inhibiting activity, which can be accessed either experimentally or using free energy calculations.

b. Optimizing the ADMET profile of hyaluronic acid. The extremely small molecular weight of hyaluronic acid is both a blessing and a curse—it enables the easy penetration of the skin while also triggers inflammatory reactions⁹⁸. The chemical modification of hyaluronic acid¹⁵⁶ has been proven to be an effective avenue towards efficient drug delivery. Using the modification strategies outlined in Bokaty, Dubashynskaya, and Skorik¹⁵⁶, we can chemically decorate the backbone with inactive functional groups to moderately increase the molecular weight of hyaluronic acid, while not altering its ability to hold water. These modifications can be proposed either by generative models or fragment-based structure enumeration and library generation tools such as Polishchuk¹⁵⁷, while the physical properties and the ADMET profile can be predicted using a machine learning-enabled QSAR model.

c. Cosmetic peptide design. The majority of peptide design in the current beauty and skincare market still relies heavily on traditional, resource-intensive methods that are both time-consuming and laborious. These conventional approaches often result in peptides that lack the necessary modifications to enhance their skin penetration and bioactivity. However, with the rapid advancements in deep learning, particularly the application of diffusion models, peptide sequence prediction and optimization have become more efficient and precise. By training diffusion models on extensive datasets of known cosmeceutical peptides, researchers can capture the complex patterns in their sequence distribution and functional properties, enabling the generation of novel peptides with desired characteristics. Specifically, advanced peptide design using diffusion models can utilize existing cosmetic peptides as input and, through conditional generation, produce new sequences that exhibit multiple skincare benefits. For instance, these generated peptides can be designed to possess not only antioxidant and anti-inflammatory properties but also promote collagen synthesis and provide skin-whitening effects. Moreover, through multiple iterations of model evaluation and screening, the efficacy and safety of the generated peptides can be validated, ensuring their reliability in practical applications. During the peptide design process, targeted chemical modifications can be incorporated, such as cyclization, lipid chain conjugation, or glycosylation of terminal groups, to enhance the peptides' penetration ability and stability. These modifications not only improve the efficiency of peptide translocation across the skin barrier but also increase their bioactivity at the target site, thereby ensuring the effective realization of multiple cosmetic benefits. Through iterative evaluation and selection, researchers can continuously optimize and identify the most promising peptide sequences, eliminating ineffective or potentially harmful molecules, thus significantly enhancing the quality and efficacy of the final products.

d. Towards precision beauty design: personalized mixture for cosmetics and skin care. Drawing inspirations from the rich literature of *drug combinations*¹⁵⁸, which characterizes the synergistic effect among multiple small molecule drugs on multi-factorial diseases, we propose to design a system to design formulae of skin-care and cosmetic mixtures that cater to personal conditions and requirements. For instance, such a system can take a picture of a person's face, and propose various solutions for lipstick mixtures, which contain waxes, oils, pigments, moisturizers, and antioxidants, and simulate their effects based upon techniques such as spectra prediction.

VIII. Conclusions

From this perspective, we discuss the contributions of computational tools, including molecular dynamics and machine learning, to the discovery and development of cosmetic molecules. Cosmetic molecules exhibit a high degree of similarity to drug molecules, and the development of corresponding cosmetic molecules is still in its early stages, offering significant potential for further growth. At the end of this paper,

we present a roadmap outlining several feasible approaches. These approaches represent the future of small molecules, proteins, and physics-based comprehensive product development, respectively.

Disclosures.

YW has limited financial interests in Flagship Pioneering, Inc. and its subsidiaries.

Funding.

YW acknowledges support from the Schmidt Science Fellowship, in partnership with the Rhodes Trust, and the Simons Center for Computational Physical Chemistry at New York University. Computational resources from the Washington Square and Abu Dhabi campuses of New York University, as well as from Memorial Sloan Kettering Cancer Center, have been used in this study.

¹“Why Did We Start Wearing Makeup? | Britannica — britannica.com,” [Accessed 07-10-2024].

²X.-Y. Meng, H.-X. Zhang, M. Mezei, and M. Cui, “Molecular docking: a powerful approach for structure-based drug discovery,” *Current computer-aided drug design* **7**, 146–157 (2011).

³J. Jumper, R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates, A. Žídek, A. Potapenko, *et al.*, “Highly accurate protein structure prediction with alphafold,” *nature* **596**, 583–589 (2021).

⁴J. Abramson, J. Adler, J. Dunger, R. Evans, T. Green, A. Pritzel, O. Ronneberger, L. Willmore, A. J. Ballard, J. Bambrick, *et al.*, “Accurate structure prediction of biomolecular interactions with alphafold 3,” *Nature*, 1–3 (2024).

⁵M. Retchin, Y. Wang, K. Takaba, and J. D. Chodera, “Druggym: A testbed for the economics of autonomous drug discovery,” *bioRxiv*, 2024–05 (2024).

⁶S. Dara, S. Dhamecherla, S. S. Jadav, C. M. Babu, and M. J. Ahsan, “Machine learning in drug discovery: a review,” *Artif. Intell. Rev.* **55**, 1947–1999 (2022).

⁷H. Wang, T. Fu, Y. Du, W. Gao, K. Huang, Z. Liu, P. Chandak, S. Liu, P. Van Katwyk, A. Deac, *et al.*, “Scientific discovery in the age of artificial intelligence,” *Nature* **620**, 47–60 (2023).

⁸D. S. Wigh, J. M. Goodman, and A. A. Lapkin, “A review of molecular representation in the age of machine learning,” *WIREs Comput. Mol. Sci.* **12**, e1603 (2022).

⁹T. N. Kipf and M. Welling, “Semi-supervised classification with graph convolutional networks,” *CoRR* **abs/1609.02907** (2016), arXiv:1609.02907.

¹⁰K. Xu, W. Hu, J. Leskovec, and S. Jegelka, “How powerful are graph neural networks?” arXiv preprint arXiv:1810.00826 (2018).

¹¹J. Gilmer, S. S. Schoenholz, P. F. Riley, O. Vinyals, and G. E. Dahl, “Neural message passing for quantum chemistry,” arXiv preprint arXiv:1704.01212 (2017).

¹²W. Hamilton, Z. Ying, and J. Leskovec, “Inductive representation learning on large graphs,” in *Advances in neural information processing systems* (2017) pp. 1024–1034.

¹³P. W. Battaglia, J. B. Hamrick, V. Bapst, A. Sanchez-Gonzalez, V. Zambaldi, M. Malininowski, A. Tacchetti, D. Raposo, A. Santoro, R. Faulkner, *et al.*, “Relational inductive biases, deep learning, and graph networks,” arXiv preprint arXiv:1806.01261 (2018).

¹⁴Y. Wang and K. Cho, “Non-convolutional graph neural networks,” (2024), arXiv:2408.00165 [cs.LG].

¹⁵Y. Wang and T. Karalatsos, “Stochastic aggregation in graph neural networks,” (2021), arXiv:2102.12648 [stat.ML].

- ¹⁶Y. Wang, M. Nguyen, M. Retchin, J. D. Chodera, and T. Karalatsos, "Bayesian active drug discovery," in https://realworldml.github.io/files/cr/48_BADD_paper_workshop.pdf.
- ¹⁷D. M. Blei, A. Kucukelbir, and J. D. McAuliffe, "Variational inference: A review for statisticians," *J. Am. Stat. Assoc.* **112**, 859–877 (2017).
- ¹⁸A. Kristiadi, F. Strieth-Kalthoff, M. Skreta, P. Poupart, A. Aspuru-Guzik, and G. Pleiss, "A sober look at llms for material discovery: Are they actually good for bayesian optimization over molecules?" (2024), arXiv:2402.05015 [cs.LG].
- ¹⁹J. S. Smith, O. Isayev, and A. E. Roitberg, "Ani-1: an extensible neural network potential with dft accuracy at force field computational cost," *Chem. Sci.* **8**, 3192–3203 (2017).
- ²⁰K. T. Schütt, O. T. Unke, and M. Gastegger, "Equivariant message passing for the prediction of tensorial properties and molecular spectra," (2021), arXiv:2102.03150 [cs.LG].
- ²¹P. Thölkke and G. D. Fabritius, "Torchmd-net: Equivariant transformers for neural network based molecular potentials," (2022), arXiv:2202.02541 [cs.LG].
- ²²O. T. Unke and M. Meuwly, "Physnet: A neural network for predicting energies, forces, dipole moments, and partial charges," *J. Chem. Theory Comput.* **15**, 3678–3693 (2019).
- ²³P. Eastman, R. Galvelis, R. P. Peláez, C. R. Abreu, S. E. Farr, E. Gallicchio, A. Gorenko, M. M. Henry, F. Hu, J. Huang, *et al.*, "Openmm 8: molecular dynamics simulation with machine learning potentials," *J. Phys. Chem. B* **128**, 109–116 (2023).
- ²⁴Y. Wang and J. D. Chodera, "Spatial attention kinetic networks with e(n)-equivariance," in *ICLR 2023* (2023).
- ²⁵Y. Wang, J. Fass, B. Kaminow, J. E. Herr, D. Rufa, I. Zhang, I. Pulido, M. Henry, H. E. B. Macdonald, K. Takaba, *et al.*, "End-to-end differentiable construction of molecular mechanics force fields," *Chem. Sci.* **13**, 12016–12033 (2022).
- ²⁶K. Takaba, A. J. Friedman, C. E. Cavender, P. K. Behara, I. Pulido, M. M. Henry, H. MacDermott-Opeskin, C. R. Iacovella, A. M. Nagle, A. M. Payne, *et al.*, "Machine-learned molecular mechanics force fields from large-scale quantum chemical data," *Chem. Sci.* (2024).
- ²⁷L. Wang, P. K. Behara, M. W. Thompson, T. Gokey, Y. Wang, J. R. Wagner, D. J. Cole, M. K. Gilson, M. R. Shirts, and D. L. Mobley, "The open force field initiative: Open software and open science for molecular modeling," *J. Phys. Chem. B* (2024).
- ²⁸Y. Wang, I. Pulido, K. Takaba, B. Kaminow, J. Scheen, L. Wang, and J. D. Chodera, "Espalomacharge: Machine learning-enabled ultrafast partial charge assignment," *J. Phys. Chem. A* **128**, 4160–4167 (2024).
- ²⁹Y. Wang, K. Takaba, M. S. Chen, M. Wieder, Y. Xu, T. Zhu, J. Z. H. Zhang, A. Nagle, K. Yu, X. Wang, D. J. Cole, J. A. Rackers, K. Cho, J. G. Greener, P. Eastman, S. Martiniani, and M. E. Tuckerman, "On the design space between molecular mechanics and machine learning force fields," (2024), arXiv:2409.01931 [physics.chem-ph].
- ³⁰F. Noé, S. Olsson, J. Köhler, and H. Wu, "Boltzmann generators – sampling equilibrium states of many-body systems with deep learning," (2019), arXiv:1812.01729 [stat.ML].
- ³¹G. Ahdriz, N. Bouatta, C. Floristean, S. Kadyan, Q. Xia, W. Gerecke, T. J. O'Donnell, D. Berenberg, I. Fisk, N. Zanichelli, *et al.*, "Openfold: Retraining alphafold2 yields new insights into its learning mechanisms and capacity for generalization," *Nat. Methods*, 1–11 (2024).
- ³²B. Kuhlman, G. Dantas, G. C. Ireton, G. Varani, B. L. Stoddard, and D. Baker, "Design of a novel globular protein fold with atomic-level accuracy," *science* **302**, 1364–1368 (2003).
- ³³R. Hefford, "Colourants and dyes for the cosmetics industry," in *Handbook of textile and industrial dyeing* (Elsevier, 2011) pp. 175–203.
- ³⁴P. A. Lewis, "Colorants: organic and inorganic pigments," in *AZimuth*, Vol. 1 (Elsevier, 1998) pp. 283–312.
- ³⁵G. Giancola and M. L. Schlossman, "Decorative cosmetics," *Cosmeceuticals and Active Cosmetics* **191** (2015).
- ³⁶P. Neytal, N. Maji, and S. Maji, "Advances in cosmetic products towards a new future," in *Preserving Health, Preserving Earth: The Path to Sustainable Healthcare* (Springer, 2024) pp. 193–214.
- ³⁷H. Lam, "Factors enhancing adhesion of color cosmetic products to skin: the role of pigments and fillers," *Surf. Sci. Adhes. Cosmet.*, 487–541 (2021).
- ³⁸A. Gürses, M. Açıkıldız, K. Güneş, M. S. Gürses, A. Gürses, M. Açıkıldız, K. Güneş, and M. S. Gürses, "Colorants in health and environmental aspects," *Dyes Pigments*, 69–83 (2016).
- ³⁹H. Zollinger, *Color chemistry: syntheses, properties, and applications of organic dyes and pigments* (John Wiley & Sons, 2003).
- ⁴⁰G. T. Sigurdson, P. Tang, and M. M. Giusti, "Natural colorants: Food colorants from natural sources," *Annu. Rev. Food Sci. Technol.* **8**, 261–280 (2017).
- ⁴¹G. Lanzendörfer-Yu, "Colorants in cosmetic applications," *Phys. Sci. Rev.* **6**, 835–877 (2021).
- ⁴²J. A. Kiernan, "Classification and naming of dyes, stains and fluorochromes," *Biotech. Histochem.* **76**, 261–278 (2001).
- ⁴³A. A. Renita, T. K. Gajaria, S. Sathish, J. A. Kumar, D. S. Lakshmi, J. Kujawa, and W. Kujawski, "Progress and prospective of the industrial development and applications of eco-friendly colorants: an insight into environmental impact and sustainability issues," *Foods* **12**, 1521 (2023).
- ⁴⁴V. Rogiers, "Eemco guidance for the assessment of transepidermal water loss in cosmetic sciences," *Skin Pharmacol. Physiol.* **14**, 117–128 (2001).
- ⁴⁵L. Rodrigues, P. Pinto, N. Galego, P. A. Da Silva, and L. M. Pereira, "Transepidermal water loss kinetic modeling approach for the parameterization of skin water dynamics," *Skin Res. Technol.* **5**, 72–82 (1999).
- ⁴⁶H. de Clermont-Gallerande, "Functional roles of lipids in make-up products," *OCL* **27**, 33 (2020).
- ⁴⁷N. Smit, J. Vicanova, and S. Pavel, "The hunt for natural skin whitening agents," *Int. J. Mol. Sci.* **10**, 5326–5349 (2009).
- ⁴⁸X. Lai, H. J. Wichers, M. Soler-Lopez, and B. W. Dijkstra, "Structure and function of human tyrosinase and tyrosinase-related proteins," *Chem. Eur. J.* **24**, 47–55 (2018).
- ⁴⁹T.-S. Chang, "An updated review of tyrosinase inhibitors," *Int. J. Mol. Sci.* **10**, 2440–2475 (2009).
- ⁵⁰Y. C. Boo, "Arbutin as a skin depigmenting agent with antimelanogenic and antioxidant properties," *Antioxidants* **10**, 1129 (2021).
- ⁵¹K. U. Zaidi, S. A. Ali, A. Ali, and I. Naaz, "Natural tyrosinase inhibitors: Role of herbals in the treatment of hyperpigmentary disorders," *Mini Rev. Med. Chem.* **19**, 796–808 (2019).
- ⁵²S. M. Petrescu, A.-J. Petrescu, H. N. Titu, R. A. Dwek, and F. M. Platt, "Inhibition of n-glycan processing in b16 melanoma cells results in inactivation of tyrosinase but does not prevent its transport to the melanosome," *J. Biol. Chem.* **272**, 15796–15803 (1997).
- ⁵³R. Logesh, S. R. Prasad, S. Chipurupalli, N. Robinson, and S. K. Mohankumar, "Natural tyrosinase enzyme inhibitors: A path from melanin to melanoma and its reported pharmacological activities," *Biochim. Biophys. Acta Rev. Cancer*, 188968 (2023).
- ⁵⁴J. Shi, Y. Guo, H. Wang, Y. Xiao, W. Liu, and L. Lyu, "The ubiquitin-proteasome system in melanin metabolism," *J. Cosmet. Dermatol.* **21**, 6661–6668 (2022).
- ⁵⁵J.-y. Wang, X.-y. Xie, Y. Deng, H.-q. Yang, X.-s. Du, P. Liu, and Y. Du, "Licorice zinc suppresses melanogenesis via inhibiting the activation of p38mapk and jnk signaling pathway in balb/c mice skin," *Acta Cir. Bras.* **37**, e371002 (2022).
- ⁵⁶I. Kusumawati and G. Indrayanto, "Natural antioxidants in cosmetics," *Stud. Nat. Prod. Chem.* **40**, 485–505 (2013).
- ⁵⁷R. Costa and L. Santos, "Delivery systems for cosmetics-from manufacturing to the skin of natural antioxidants," *Powder Technol.* **322**, 402–416 (2017).
- ⁵⁸S. Murad, D. Grove, K. Lindberg, G. Reynolds, A. Sivarajah, and S. Pinnell, "Regulation of collagen synthesis by ascorbic acid," *Proc. Natl. Acad. Sci.* **78**, 2879–2882 (1981).
- ⁵⁹A. K. Mohiuddin, "Skin aging & modern age anti-aging strategies," *Int. J. Clin. Dermatol. Res.* **7**, 209–240 (2019).
- ⁶⁰L. Talakoub, I. M. Neuhaus, and S. S. Yu, "Antiaging cosmeceuticals," *Int. Textb. Aesthet. Surg.*, 1183–1209 (2016).
- ⁶¹K. V. Sajna, L. D. Gottumukkala, R. K. Sukumaran, and A. Pandey, "White biotechnology in cosmetics," in *Industrial biorefineries & white biotechnology* (Elsevier, 2015) pp. 607–652.
- ⁶²M. G. Traber and J. Atkinson, "Vitamin e, antioxidant and nothing more," *Free radical biology and medicine* **43**, 4–15 (2007).
- ⁶³C. A. Dinarello, "Anti-inflammatory agents: present and future," *Cell* **140**, 935–950 (2010).

- ⁶⁴N. P. Stamford, "Stability, transdermal penetration, and cutaneous effects of ascorbic acid and its derivatives," *J. Cosmet. Dermatol.* **11**, 310–317 (2012).
- ⁶⁵H. A. Benson, "Transdermal drug delivery: penetration enhancement techniques," *Curr. Drug Deliv.* **2**, 23–33 (2005).
- ⁶⁶E. Rostkowska, E. Poleszak, K. Wojciechowska, and K. Dos Santos Szewczyk, "Dermatological management of aged skin," *Cosmetics* **10**, 55 (2023).
- ⁶⁷S. H. Lim, Y. Sun, T. Thiruvallur Madanagopal, V. Rosa, and L. Kang, "Enhanced skin permeation of anti-wrinkle peptides via molecular modification," *Sci. Rep.* **8**, 1596 (2018).
- ⁶⁸S. M. Mortazavi and H. R. Moghimi, "Skin permeability, a dismissed necessity for anti-wrinkle peptide performance," *Int. J. Cosmet. Sci.* **44**, 232–248 (2022).
- ⁶⁹A. Ahad, M. Aqil, K. Kohli, H. Chaudhary, Y. Sultana, M. Mujeeb, and S. Talegaonkar, "Chemical penetration enhancers: a patent review," *Expert Opin. Ther. Pat.* **19**, 969–988 (2009).
- ⁷⁰M. Kumar and R. Ghirmikar, "Proteins and peptides in personal care," in *Biotechnology in Personal Care* (CRC Press, 2016) pp. 81–110.
- ⁷¹P. Karande, A. Jain, K. Ergun, V. Kispersky, and S. Mitragotri, "Design principles of chemical penetration enhancers for transdermal drug delivery," *Proc. Natl. Acad. Sci.* **102**, 4688–4693 (2005).
- ⁷²L. Zeng, F. Huang, Q. Zhang, J. Liu, D. Quan, and W. Song, "Molecular perspective of efficiency and safety problems of chemical enhancers: bottlenecks and recent advances," *Drug Deliv. Transl. Res.* , 1–19 (2022).
- ⁷³S. Kumar, M. Zakrewsky, M. Chen, S. Menegatti, J. A. Muraski, and S. Mitragotri, "Peptides as skin penetration enhancers: mechanisms of action," *J. Control. Release* **199**, 168–178 (2015).
- ⁷⁴V. Hmingthansanga, N. Singh, S. Banerjee, S. Manickam, R. Velayutham, and S. Natesan, "Improved topical drug delivery: Role of permeation enhancers and advanced approaches," *Pharmaceutics* **14**, 2818 (2022).
- ⁷⁵D. M. Copolovici, K. Langel, E. Eriste, and U. Langel, "Cell-penetrating peptides: design, synthesis, and applications," *ACS Nano* **8**, 1972–1994 (2014).
- ⁷⁶G. Guidotti, L. Brambilla, and D. Rossi, "Cell-penetrating peptides: from basic research to clinics," *Trends Pharmacol. Sci.* **38**, 406–424 (2017).
- ⁷⁷C. G. Gennari, S. Franzè, S. Pellegrino, E. Corsini, G. Vistoli, L. Montanari, P. Minghetti, and F. Cilurzo, "Skin penetrating peptide as a tool to enhance the permeation of heparin through human epidermis," *Biomacromolecules* **17**, 46–55 (2016).
- ⁷⁸T. Hsu and S. Mitragotri, "Delivery of sirna and other macromolecules into skin and cells using a peptide enhancer," *Proc. Natl. Acad. Sci.* **108**, 15816–15821 (2011).
- ⁷⁹J. M. Lim, M. Y. Chang, S. G. Park, N. G. Kang, Y. S. Song, Y. H. Lee, Y. C. Yoo, W. G. Cho, S. Y. Choi, and S. H. Kang, "Penetration enhancement in mouse skin and lipolysis in adipocytes by tat-gkh, a new cosmetic ingredient," *J. Cosmet. Sci.* **54**, 483–492 (2003).
- ⁸⁰M. Chen, V. Gupta, A. C. Anselmo, J. A. Muraski, and S. Mitragotri, "Topical delivery of hyaluronic acid into skin using space-peptide carriers," *J. Control. Release* **173**, 67–74 (2014).
- ⁸¹N. M. Carmichael, J. O. Dostrovsky, and M. P. Charlton, "Peptide-mediated transdermal delivery of botulinum neurotoxin type a reduces neurogenic inflammation in the skin," *Pain* **149**, 316–324 (2010).
- ⁸²G.-H. Hur, S.-C. Han, A.-R. Ryu, Y. Eom, J.-W. Kim, and M.-Y. Lee, "Effect of oligoarginine conjugation on the antiwrinkle activity and transdermal delivery of ghk peptide," *J. Pept. Sci.* **26**, e3234 (2020).
- ⁸³M. Hussain and D. J. Goldberg, "Topical manganese peptide in the treatment of photodamaged skin," *J. Cosmet. Laser Ther.* **9**, 232–236 (2007).
- ⁸⁴F. Errante, P. Ledwoń, R. Latajka, P. Rovero, and A. M. Papini, "Cosmeceutical peptides in the framework of sustainable wellness economy," *Front. Chem.* **8**, 572923 (2020).
- ⁸⁵V. Arul, D. Gopinath, K. Gomathi, and R. Jayakumar, "Biotinylated ghk peptide incorporated collagenous matrix: A novel biomaterial for dermal wound healing in rats," *J. Biomed. Mater. Res. B Appl. Biomater.* **73**, 383–391 (2005).
- ⁸⁶S. Jeong, S. Yoon, S. Kim, J. Jung, M. Kor, K. Shin, C. Lim, H. S. Han, H. Lee, K.-Y. Park, *et al.*, "Anti-wrinkle benefits of peptides complex stimulating skin basement membrane proteins expression," *Int. J. Mol. Sci.* **21**, 73 (2019).
- ⁸⁷G. M. El Maghraby, B. W. Barry, and A. Williams, "Liposomes and skin: from drug delivery to model membranes," *Eur. J. Pharm. Sci.* **34**, 203–222 (2008).
- ⁸⁸M. F. Peralta, M. L. Guzmán, A. Pérez, G. A. Apezteguia, M. L. Fórmica, E. L. Romero, M. E. Olivera, and D. C. Carrer, "Liposomes can both enhance or reduce drugs penetration through the skin," *Sci. Rep.* **8**, 13253 (2018).
- ⁸⁹S. Rai, V. Pandey, and G. Rai, "Transfersomes as versatile and flexible nano-vesicular carriers in skin cancer therapy: The state of the art," *Nano Rev. Exp.* **8**, 1325708 (2017).
- ⁹⁰J. Bouwstra and P. Honeywell-Nguyen, "Skin structure and mode of action of vesicles," *Adv. Drug Deliv. Rev.* **54**, S41–S55 (2002).
- ⁹¹S.-Y. Pu, Y.-L. Huang, C.-M. Pu, Y.-N. Kang, K. D. Hoang, K.-H. Chen, and C. Chen, "Effects of oral collagen for skin anti-aging: A systematic review and meta-analysis," *Nutrients* **15**, 2080 (2023).
- ⁹²N. Inoue, F. Sugihara, and X. Wang, "Ingestion of bioactive collagen hydrolysates enhance facial skin moisture and elasticity and reduce facial ageing signs in a randomised double-blind placebo-controlled clinical study," *Journal of the Science of Food and Agriculture* **96**, 4077–4081 (2016).
- ⁹³F. Sugihara, N. Inoue, and X. Wang, "Clinical effects of ingesting collagen hydrolysate on facial skin properties," *Jpn Pharmacol Ther* **43**, 67–70 (2015).
- ⁹⁴J. Z. Williams, N. Abumrad, and A. Barbul, "Effect of a specialized amino acid mixture on human collagen deposition," *Annals of surgery* **236**, 369–375 (2002).
- ⁹⁵E. Arribas-López, N. Zand, O. Ojo, M. J. Snowden, and T. Kochhar, "The effect of amino acids on wound healing: a systematic review and meta-analysis on arginine and glutamine," *Nutrients* **13**, 2498 (2021).
- ⁹⁶V. L. Albaugh, K. Mukherjee, and A. Barbul, "Proline precursors and collagen synthesis: biochemical challenges of nutrient supplementation and wound healing," *The Journal of nutrition* **147**, 2011–2017 (2017).
- ⁹⁷N. S. Trookman, R. L. Rizer, R. Ford, E. Ho, and V. Gotz, "Immediate and long-term clinical benefits of a topical treatment for facial lines and wrinkles," *The Journal of clinical and aesthetic dermatology* **2**, 38 (2009).
- ⁹⁸B. Bravo, P. Correia, J. Gonçalves Junior, B. Sant'Anna, and D. Kerob, "Benefits of topical hyaluronic acid for skin quality and signs of skin aging: From literature review to clinical evidence," *Dermatol. Ther.* **35**, e15903 (2022).
- ⁹⁹M. Juhaščík, A. Kováčik, and G. Huerta-Ángeles, "Recent advances of hyaluronan for skin delivery: From structure to fabrication strategies and applications," *Polymers* **14** (2022), 10.3390/polym14224833.
- ¹⁰⁰J. Allen and K. Dodou, "Current knowledge and regulatory framework on the use of hyaluronic acid for aesthetic injectable skin rejuvenation treatments," *Cosmetics* **11** (2024), 10.3390/cosmetics11020054.
- ¹⁰¹B. David, J.-L. Wolfender, and D. A. Dias, "The pharmaceutical industry and natural products: historical status and new trends," *Phytochem. Rev.* **14**, 299–315 (2015).
- ¹⁰²T. Pillaiyar, M. Manickam, and V. Namasivayam, "Skin whitening agents: Medicinal chemistry perspective of tyrosinase inhibitors," *J. Enzyme Inhib. Med. Chem.* **32**, 403–425 (2017).
- ¹⁰³L. Yue, L. Song, S. Zhu, X. Fu, X. Li, C. He, and J. Li, "Machine learning assisted rational design of antimicrobial peptides based on human endogenous proteins and their applications for cosmetic preservative system optimization," *Sci. Rep.* **14**, 947 (2024).
- ¹⁰⁴H. T. Hsueh, R. T. Chou, U. Rai, W. Liyanage, Y. C. Kim, M. B. Appell, J. Pejavar, K. T. Leo, C. Davison, P. Kolodziejski, *et al.*, "Machine learning-driven multifunctional peptide engineering for sustained ocular drug delivery," *Nat. Commun.* **14**, 2509 (2023).
- ¹⁰⁵E. C. L. de Oliveira, K. Santana, L. Josino, A. H. Lima e Lima, and C. de Souza de Sales Júnior, "Predicting cell-penetrating peptides using machine learning algorithms and navigating in their chemical space," *Sci. Rep.* **11**, 7628 (2021).
- ¹⁰⁶V. Apostolopoulos, J. Bojarska, T.-T. Chai, S. Elnagdy, K. Kaczmarek, J. Matsoukas, R. New, K. Parang, O. P. Lopez, H. Parhiz, *et al.*, "A global review on short peptides: frontiers and perspectives," *Molecules* **26**, 430 (2021).
- ¹⁰⁷R. R. Aruan, H. Hutabarat, A. A. Widodo, M. T. C. C. Firdiyono, C. Wirawanty, and L. Fransiska, "Double-blind, randomized trial on the effectiveness of acetylhexapeptide-3 cream and palmitoyl pentapeptide-4

- cream for crow's feet," *J. Clin. Aesthet. Dermatol.* **16**, 37 (2023).
- ¹⁰⁸R. Ganceviciene, A. I. Liakou, A. Theodoridis, E. Makrantonaki, and C. C. Zouboulis, "Skin anti-aging strategies," *Dermato-endocrinology* **4**, 308–319 (2012).
- ¹⁰⁹A. Martins, H. Vieira, H. Gaspar, and S. Santos, "Marketed marine natural products in the pharmaceutical and cosmeceutical industries: Tips for success," *Mar. Drugs* **12**, 1066–1101 (2014).
- ¹¹⁰C. Bournez, J.-M. Gally, S. Aci-Sèche, P. Bernard, and P. Bonnet, "Virtual screening of natural products to enhance melanogenesis," *Mol. Inform.*, e202300335 (2024).
- ¹¹¹Y. Su, S. Xu, X. Hu, R. Wang, M. Dong, Y. Wang, S. Wang, Y. Zhang, Q. Tian, and L. Han, "Rapid discovery of natural skin-lightening ingredients based on an integrated screening strategy based on molecular docking and zebrafish model," *J. Cosmet. Dermatol.* (2024).
- ¹¹²N. Ai, W. J. Welsh, U. Santhanam, H. Hu, and J. Lyga, "Novel virtual screening approach for the discovery of human tyrosinase inhibitors," *PLoS one* **9**, e112788 (2014).
- ¹¹³A. Kumari, R. kumar, G. Sulabh, P. Singh, J. Kumar, V. K. Singh, and K. K. Ojha, "In silico admet, molecular docking and molecular simulation-based study of glabridin's natural and semisynthetic derivatives as potential tyrosinase inhibitors," *Advances in Traditional Medicine* **23**, 733–751 (2023).
- ¹¹⁴D. A. Case, H. M. Aktulga, K. Belfon, I. Ben-Shalom, S. R. Brozell, D. S. Cerutti, T. E. Cheatham III, V. W. D. Cruzeiro, T. A. Darden, R. E. Duke, *et al.*, *Amber 2021* (University of California, San Francisco, 2021).
- ¹¹⁵J. Lee, X. Cheng, S. Jo, A. D. MacKerell, J. B. Klauda, and W. Im, "Charmm-gui input generator for namd, gromacs, amber, openmm, and charmm/openmm simulations using the charmm36 additive force field," *Biophys. J.* **110**, 641a (2016).
- ¹¹⁶M. J. Abraham, T. Murtola, R. Schulz, S. Páll, J. C. Smith, B. Hess, and E. Lindahl, "Gromacs: High performance molecular simulations through multi-level parallelism from laptops to supercomputers," *SoftwareX* **1**, 19–25 (2015).
- ¹¹⁷S. Decherchi and A. Cavalli, "Thermodynamics and kinetics of drug-target binding by molecular simulation," *Chem. Rev.* **120**, 12788–12833 (2020).
- ¹¹⁸P. Agu, C. Afiukwa, O. Orji, E. Ezeh, I. Ofoke, C. Ogbu, E. Ugwuja, and P. Aja, "Molecular docking as a tool for the discovery of molecular targets of nutraceuticals in diseases management," *Sci. Rep.* **13**, 13398 (2023).
- ¹¹⁹D. L. Mobley and M. K. Gilson, "Predicting binding free energies: frontiers and benchmarks," *Annual review of biophysics* **46**, 531–558 (2017).
- ¹²⁰E. Wang, H. Sun, J. Wang, Z. Wang, H. Liu, J. Z. Zhang, and T. Hou, "End-point binding free energy calculation with mm/pbsa and mm/gbsa: strategies and applications in drug design," *Chem. Rev.* **119**, 9478–9508 (2019).
- ¹²¹S. Genheden and U. Ryde, "The mm/pbsa and mm/gbsa methods to estimate ligand-binding affinities," *Expert Opin. Drug Discov.* **10**, 449–461 (2015).
- ¹²²X. Liu, L. Peng, Y. Zhou, Y. Zhang, and J. Z. Zhang, "Computational alanine scanning with interaction entropy for protein–ligand binding free energies," *J. Chem. Theory Comput.* **14**, 1772–1780 (2018).
- ¹²³Y. Yan, M. Yang, C. G. Ji, and J. Z. Zhang, "Interaction entropy for computational alanine scanning," *J. Chem. Inf. Model.* **57**, 1112–1122 (2017).
- ¹²⁴S. Bruckner and S. Boresch, "Efficiency of alchemical free energy simulations. ii. improvements for thermodynamic integration," *Journal of computational chemistry* **32**, 1320–1333 (2011).
- ¹²⁵W. L. Jorgensen and L. L. Thomas, "Perspective on free-energy perturbation calculations for chemical equilibria," *Journal of chemical theory and computation* **4**, 869–876 (2008).
- ¹²⁶A. S. Mey, B. K. Allen, H. E. B. Macdonald, J. D. Chodera, D. F. Hahn, M. Kuhn, J. Michel, D. L. Mobley, L. N. Naden, S. Prasad, *et al.*, "Best practices for alchemical free energy calculations [article v1. 0]," *Living journal of computational molecular science* **2** (2020).
- ¹²⁷M. R. Shirts, D. L. Mobley, and J. D. Chodera, "Alchemical free energy calculations: ready for prime time?" *Annual reports in computational chemistry* **3**, 41–59 (2007).
- ¹²⁸M. T. Cronin, S. J. Enoch, J. C. Madden, J. F. Rathman, A.-N. Richarz, and C. Yang, "A review of in silico toxicology approaches to support the safety assessment of cosmetics-related materials," *Comput. Toxicol.* **21**, 100213 (2022).
- ¹²⁹H. Xin, A. S. Virk, S. S. Virk, F. Akin-Ige, and S. Amin, "Applications of artificial intelligence and machine learning on critical materials used in cosmetics and personal care formulation design," *Curr. Opin. Colloid Interface Sci.*, 101847 (2024).
- ¹³⁰A. Wilm, J. Kühnl, and J. Kirchmair, "Computational approaches for skin sensitization prediction," *Crit. Rev. Toxicol.* **48**, 738–760 (2018).
- ¹³¹R. Lochhead *et al.*, "The use of polymers in cosmetic products," *Cosmet. Sci. Technol.* **13**, 171–221 (2017).
- ¹³²R. Lochhead, "Basic physical sciences for the formulation of cosmetic products," *Cosmet. Sci. Technol.*, 39–76 (2017).
- ¹³³A. Paudel, D. Rajjada, and J. Rantanen, "Raman spectroscopy in pharmaceutical product design," *Adv. Drug Deliv. Rev.* **89**, 3–20 (2015).
- ¹³⁴D. Kowalczyk, A. Gładysz, M. Pitucha, D. M. Kamiński, A. Barańska, and B. Drop, "Spectroscopic study of the molecular structure of the new hybrid with a potential two-way antibacterial effect," *Molecules* **26**, 1442 (2021).
- ¹³⁵F. Rico, A. Mazabel, G. Egurrola, J. Pulido, N. Barrios, R. Marquez, and J. García, "Meta-analysis and analytical methods in cosmetics formulation: A review," *Cosmetics* **11**, 1 (2023).
- ¹³⁶N. Prieto, R. Roehle, P. Lavín, G. Batten, and S. Andrés, "Application of near infrared reflectance spectroscopy to predict meat and meat products quality: A review," *Meat Sci.* **83**, 175–186 (2009).
- ¹³⁷C. Cole, J. Silverman, and M. Bonitatibus, "Evaluating sunscreen ultraviolet protection using a polychromatic diffuse reflectance device," *Photo-dermatol. Photoimmunol. Photomed.* **35**, 436–441 (2019).
- ¹³⁸R. Gómez-Bombarelli, J. N. Wei, D. Duvenaud, J. M. Hernández-Lobato, B. Sánchez-Lengeling, D. Sheberla, J. Aguilera-Iparraguirre, T. D. Hirzel, R. P. Adams, and A. Aspuru-Guzik, "Automatic chemical design using a data-driven continuous representation of molecules," *ACS central science* **4**, 268–276 (2018).
- ¹³⁹T. N. Kipf and M. Welling, "Variational graph auto-encoders," (2016), arXiv:1611.07308 [stat.ML].
- ¹⁴⁰W. Jin, R. Barzilay, and T. Jaakkola, "Junction tree variational auto-encoder for molecular graph generation," (2019), arXiv:1802.04364 [cs.LG].
- ¹⁴¹C. Zang and F. Wang, "Moflow: An invertible flow model for generating molecular graphs," in *Proceedings of the 26th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining, KDD '20* (ACM, 2020).
- ¹⁴²C. Liu, W. Fan, Y. Liu, J. Li, H. Li, H. Liu, J. Tang, and Q. Li, "Generative diffusion models on graphs: Methods and applications," (2023), arXiv:2302.02591 [cs.LG].
- ¹⁴³E. Hoogeboom, V. G. Satorras, C. Vignac, and M. Welling, "Equivariant diffusion for molecule generation in 3d," (2022), arXiv:2203.17003 [cs.LG].
- ¹⁴⁴A. Vaswani, N. Shazeer, N. Parmar, J. Uszkoreit, L. Jones, A. N. Gomez, L. Kaiser, and I. Polosukhin, "Attention is all you need," (2023), arXiv:1706.03762 [cs.CL].
- ¹⁴⁵H. Touvron, T. Lavril, G. Izacard, X. Martinet, M.-A. Lachaux, T. Lacroix, B. Rozière, N. Goyal, E. Hambro, F. Azhar, A. Rodriguez, A. Joulin, E. Grave, and G. Lample, "Llama: Open and efficient foundation language models," (2023), arXiv:2302.13971 [cs.CL].
- ¹⁴⁶OpenAI, "Chatgpt," <https://chat.openai.com> (2023), accessed: 2024-09-25.
- ¹⁴⁷T. B. Brown, B. Mann, N. Ryder, M. Subbiah, J. Kaplan, P. Dhariwal, A. Neelakantan, P. Shyam, G. Sastry, A. Askell, S. Agarwal, A. Herbert-Voss, G. Krueger, T. Henighan, R. Child, A. Ramesh, D. M. Ziegler, J. Wu, C. Winter, C. Hesse, M. Chen, E. Sigler, M. Litwin, S. Gray, B. Chess, J. Clark, C. Berner, S. McCandlish, A. Radford, I. Sutskever, and D. Amodei, "Language models are few-shot learners," (2020), arXiv:2005.14165 [cs.CL].
- ¹⁴⁸E. M. Bender, T. Gebru, A. McMillan-Major, and S. Shmitchell, "On the dangers of stochastic parrots: Can language models be too big?" in *Proceedings of the 2021 ACM conference on fairness, accountability, and transparency* (2021) pp. 610–623.
- ¹⁴⁹H. Ni, S. Meng, X. Chen, Z. Zhao, A. Chen, P. Li, S. Zhang, Q. Yin, Y. Wang, and Y. Chan, "Harnessing earnings reports for stock predictions: A qlora-enhanced llm approach," (2024), arXiv:2408.06634 [q-fin.CP].
- ¹⁵⁰K. M. Jablonka, P. Schwaller, A. Ortega-Guerrero, and B. Smit, "Is gpt-3 all you need for low-data discovery in chemistry?" (2023).

- ¹⁵¹B. Yu, F. N. Baker, Z. Chen, X. Ning, and H. Sun, "Llasmol: Advancing large language models for chemistry with a large-scale, comprehensive, high-quality instruction tuning dataset," (2024), arXiv:2402.09391 [cs.AI].
- ¹⁵²D. Zhang, W. Liu, Q. Tan, J. Chen, H. Yan, Y. Yan, J. Li, W. Huang, X. Yue, W. Ouyang, D. Zhou, S. Zhang, M. Su, H.-S. Zhong, and Y. Li, "Chemllm: A chemical large language model," (2024), arXiv:2402.06852 [cs.AI].
- ¹⁵³S. Zhang, L. Dong, X. Li, S. Zhang, X. Sun, S. Wang, J. Li, R. Hu, T. Zhang, F. Wu, and G. Wang, "Instruction tuning for large language models: A survey," (2024), arXiv:2308.10792 [cs.CL].
- ¹⁵⁴A. Masyita, E. Salim, R. M. Asri, F. Nainu, A. Hori, R. Yulianty, M. Hatta, Y. Rifai, and T. Kuraishi, "Molecular modeling and phenoloxidase inhibitory activity of arbutin and arbutin undecylenic acid ester," *Biochem. Biophys. Res. Commun.* **547**, 75–81 (2021).
- ¹⁵⁵Y. Zhang, Z. Zhang, D. Ke, X. Pan, X. Wang, X. Xiao, and C. Ji, "Frag-grow: A web server for structure-based drug design by fragment growing within constraints," *J. Chem. Inf. Model.* (2024).
- ¹⁵⁶A. N. Bokaty, N. V. Dubashynskaya, and Y. A. Skorik, "Chemical modification of hyaluronic acid as a strategy for the development of advanced drug delivery systems," *Carbohydrate Polymers*, 122145 (2024).
- ¹⁵⁷P. Polishchuk, "Crem: chemically reasonable mutations framework for structure generation," *Journal of Cheminformatics* **12**, 28 (2020).
- ¹⁵⁸P. Jaaks, E. A. Coker, D. J. Vis, O. Edwards, E. F. Carpenter, S. M. Leto, L. Dwane, F. Sassi, H. Lightfoot, S. Barthorpe, *et al.*, "Effective drug combinations in breast, colon and pancreatic cancer cells," *Nature* **603**, 166–173 (2022).