

Table S3. Serotype distribution of the 10 overall most common serotypes in NIPP and in IPD (2009-2011).

NIPP			IPD*		
Serotype†	n (%)	Cumulative n (%)	Serotype†	n (%)	Cumulative n (%)
3	49 (16.3)	49 (16.3)	3	160 (12.6)	160 (12.6)
19A	24 (8.0)	73 (24.3)	7F	126 (10.0)	286 (22.6)
11A	21 (7.0)	94 (31.3)	19A	115 (9.1)	401 (31.7)
6C	17 (5.7)	111 (37.0)	14	106 (8.4)	507 (40.1)
19F	17 (5.7)	128 (42.7)	1	87 (6.9)	594 (47.0)
22F	16 (5.3)	144 (48.0)	8	79 (6.2)	673 (53.2)
23B	14 (4.7)	158 (52.7)	22F	55 (4.3)	728 (57.5)
9N	12 (4.0)	170 (56.7)	4	43 (3.4)	771 (60.9)
7F	10 (3.3)	180 (60.0)	11A	39 (3.1)	810 (64.0)
8	10 (3.3)	190 (63.3)	6C	36 (2.8)	846 (66.0)

* Information published previously (19)

†Serotypes in bold were associated with NIPP or IPD ($P<0.05$)