



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 3, 2024 – 08:21 AM EST

PDB ID : 1EM2  
Title : Star-related lipid transport domain of MLN64  
Authors : Tsujishita, Y.; Hurley, J.H.  
Deposited on : 2000-03-14  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	TAR	A	500	X	-	-	-

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1891 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called MLN64 PROTEIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	214	1716	1090	307	313	3	3	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	388	MSE	PHE	engineered mutation	UNP Q14849

- Molecule 2 is D(-)-TARTARIC ACID (three-letter code: TAR) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	10	4	6	0	0


- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	165	Total 165	O 165	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MLN64 PROTEIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.42Å 83.42Å 81.95Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.00 – 2.20 25.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (25.00-2.20) 96.6 (25.00-2.20)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.15 (at 2.19Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.207 , 0.264 0.207 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.3	Xtrriage
Anisotropy	0.179	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 47.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.032 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	1891	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TAR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.90	1/1751 (0.1%)	1.03	10/2367 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	351	ARG	CA-CB	-16.31	1.18	1.53

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	352	ARG	N-CA-CB	13.10	134.18	110.60
1	A	351	ARG	CB-CA-C	-11.79	86.81	110.40
1	A	351	ARG	N-CA-CB	10.17	128.90	110.60
1	A	357	ARG	NE-CZ-NH2	8.13	124.36	120.30
1	A	361	LEU	CA-CB-CG	7.87	133.41	115.30
1	A	352	ARG	N-CA-C	-6.75	92.78	111.00
1	A	351	ARG	NE-CZ-NH2	6.64	123.62	120.30
1	A	435	ARG	NE-CZ-NH2	-5.53	117.53	120.30
1	A	236	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	A	277	PRO	N-CA-C	5.01	125.13	112.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	351	ARG	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1716	0	1714	59	0
2	A	10	0	4	0	0
3	A	165	0	0	21	3
All	All	1891	0	1718	59	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:351:ARG:NH1	3:A:692:HOH:O	1.89	1.06
1:A:406:LEU:HD23	1:A:427:MSE:HE2	1.38	1.01
1:A:432:PHE:O	3:A:717:HOH:O	1.85	0.92
1:A:351:ARG:O	1:A:351:ARG:HG2	1.68	0.91
1:A:422:SER:HB3	3:A:702:HOH:O	1.71	0.91
1:A:435:ARG:N	3:A:717:HOH:O	2.08	0.86
1:A:303:GLN:HG3	3:A:695:HOH:O	1.80	0.82
1:A:262:GLU:HG3	1:A:273:THR:HG23	1.60	0.81
1:A:415:PRO:HG3	3:A:707:HOH:O	1.83	0.78
1:A:283:PHE:HB2	1:A:427:MSE:HE1	1.62	0.78
1:A:283:PHE:CB	1:A:427:MSE:HE1	2.15	0.76
1:A:232:SER:O	1:A:236:ARG:HG3	1.91	0.70
1:A:382:GLU:OE1	3:A:669:HOH:O	2.11	0.68
1:A:262:GLU:CG	1:A:273:THR:HG23	2.24	0.68
1:A:432:PHE:C	3:A:717:HOH:O	2.27	0.67
1:A:351:ARG:CZ	3:A:692:HOH:O	2.38	0.67
1:A:306:ARG:HG2	3:A:652:HOH:O	1.95	0.67
1:A:435:ARG:HG3	3:A:711:HOH:O	1.98	0.63

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:306:ARG:HB3	3:A:642:HOH:O	1.97	0.63
1:A:358:ASP:HA	1:A:392:LYS:HD3	1.80	0.62
1:A:323:VAL:O	1:A:324:GLU:HG2	1.99	0.62
1:A:240:ARG:O	1:A:244:GLU:HG3	1.99	0.61
1:A:406:LEU:HD23	1:A:427:MSE:CE	2.22	0.61
1:A:411:LYS:HA	3:A:630:HOH:O	2.03	0.59
1:A:411:LYS:HD3	3:A:630:HOH:O	2.02	0.58
1:A:251:GLN:HG2	1:A:255:GLN:HE22	1.66	0.58
1:A:369:HIS:HD1	1:A:371:ALA:H	1.52	0.57
1:A:309:LEU:HD12	3:A:642:HOH:O	2.04	0.56
1:A:297:TYR:OH	1:A:355:ARG:NE	2.40	0.55
1:A:406:LEU:CD2	1:A:427:MSE:HE2	2.26	0.55
1:A:318:GLN:HG3	1:A:331:TYR:HB3	1.91	0.52
1:A:425:ALA:O	1:A:429:GLU:HG3	2.09	0.52
1:A:279:HIS:HD2	3:A:725:HOH:O	1.92	0.51
1:A:416:ARG:HG2	1:A:420:HIS:CE1	2.46	0.51
1:A:377:LYS:HE3	1:A:378:TYR:CE1	2.47	0.50
1:A:433:HIS:C	3:A:717:HOH:O	2.48	0.50
1:A:417:TYR:O	1:A:421:GLN:HG3	2.11	0.49
1:A:422:SER:CB	3:A:702:HOH:O	2.42	0.49
1:A:307:MSE:HG3	1:A:310:TRP:CZ2	2.48	0.49
1:A:301:ILE:HG23	1:A:362:SER:HB2	1.94	0.48
1:A:304:PRO:HA	1:A:307:MSE:CE	2.43	0.48
1:A:303:GLN:NE2	1:A:306:ARG:HG3	2.28	0.48
1:A:279:HIS:CD2	1:A:385:PRO:HD3	2.49	0.48
1:A:297:TYR:CE2	1:A:302:LEU:HD11	2.50	0.47
1:A:376:HIS:HB3	3:A:611:HOH:O	2.16	0.45
1:A:323:VAL:O	1:A:324:GLU:CG	2.64	0.45
1:A:324:GLU:HG3	1:A:326:ASN:H	1.81	0.45
1:A:303:GLN:HB3	1:A:306:ARG:CG	2.47	0.44
1:A:396:ASN:HD21	1:A:398:ARG:HB2	1.82	0.44
1:A:329:ILE:HD13	1:A:350:VAL:HA	2.00	0.44
1:A:323:VAL:O	1:A:324:GLU:C	2.57	0.43
1:A:287:THR:CB	3:A:753:HOH:O	2.66	0.43
1:A:351:ARG:HB2	1:A:352:ARG:H	1.54	0.43
1:A:307:MSE:HE2	1:A:307:MSE:HB2	1.93	0.42
1:A:311:ASN:ND2	1:A:313:THR:H	2.17	0.42
1:A:418:LEU:HD11	3:A:699:HOH:O	2.18	0.42
1:A:377:LYS:HE3	1:A:378:TYR:HE1	1.84	0.42
1:A:321:GLN:NE2	1:A:369:HIS:NE2	2.69	0.41
1:A:304:PRO:HA	1:A:307:MSE:HE2	2.02	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:672:HOH:O	3:A:672:HOH:O[4_556]	2.08	0.12
3:A:600:HOH:O	3:A:611:HOH:O[4_556]	2.14	0.06
3:A:674:HOH:O	3:A:713:HOH:O[4_556]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	212/229 (93%)	205 (97%)	5 (2%)	2 (1%)	<b>14</b> <b>14</b>

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	324	GLU
1	A	277	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	185/194 (95%)	173 (94%)	12 (6%)	<b>14</b> <b>16</b>

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	255	GLN
1	A	260	LYS
1	A	277	PRO
1	A	306	ARG
1	A	307	MSE
1	A	311	ASN
1	A	325	ASP
1	A	327	THR
1	A	351	ARG
1	A	396	ASN
1	A	407	ASN
1	A	441	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	255	GLN
1	A	258	ASN
1	A	303	GLN
1	A	311	ASN
1	A	321	GLN
1	A	396	ASN
1	A	407	ASN
1	A	420	HIS
1	A	433	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	TAR	A	500	-	9,9,9	1.43	3 (33%)	12,12,12	1.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TAR	A	500	-	2/2/4/4	0/12/12/12	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	TAR	C3-C4	2.22	1.55	1.52
2	A	500	TAR	O11-C1	-2.18	1.23	1.30
2	A	500	TAR	O41-C4	-2.02	1.24	1.30

There are no bond angle outliers.

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	500	TAR	C2
2	A	500	TAR	C3

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	211/229 (92%)	0.17	16 (7%) <b>21</b> <b>19</b>	11, 24, 45, 55	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	443	ALA	3.6
1	A	306	ARG	3.3
1	A	303	GLN	3.2
1	A	279	HIS	2.9
1	A	278	PHE	2.5
1	A	325	ASP	2.4
1	A	338	ALA	2.4
1	A	304	PRO	2.4
1	A	277	PRO	2.3
1	A	337	ALA	2.3
1	A	230	SER	2.2
1	A	376	HIS	2.2
1	A	351	ARG	2.1
1	A	339	GLY	2.1
1	A	255	GLN	2.1
1	A	440	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	TAR	A	500	10/10	0.92	0.09	22,26,28,29	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.